

10-10-00

A/Sege

10/05/00
jc929 U.S. PTO


Please type a plus sign (+) inside this box ☐

Attorney Docket **P1776R2**
PATENT

CERTIFICATION UNDER 37 CFR 1.10

EL142012682US: Express Mail Number **October 5, 2000:** Date of Deposit

I hereby certify that this Non-provisional Application Transmittal and the documents referred to as enclosed therein are being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Box PATENT APPLICATION, Director of Patents and Trademarks, Washington, D.C. 20231.


Gloria L. Tabuena

jc922 U.S. PTO
09/684458
10/05/00

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

BOX PATENT APPLICATION
Director of Patents and Trademarks
Washington, DC 20231

NON-PROVISIONAL APPLICATION TRANSMITTAL UNDER 37 CFR 1.53(b)

Transmitted herewith for filing is a non-provisional patent application:

Inventor(s) (or Application "Identifier"):

Mary E. Gerritsen, San Mateo, California
Audrey Goddard, San Francisco, California
J. Christopher Grimaldi, San Francisco, California
Fuad Mehraban, Trumbull, Connecticut

Title: **NOVEL POLYPEPTIDES, THEIR NUCLEIC ACIDS, AND METHODS FOR THEIR USE IN ANGIOGENESIS AND VASCULARIZATION**

1. Type of Application

- ☒ This application is for an original, non-provisional application.
- ☐ This is a non-provisional application claiming priority to provisional application no. __, filed __, the entire disclosure of which is hereby incorporated by reference.
- ☐ This is a ☐ continuation-in-part ☐ continuation ☐ divisional application claiming priority to application Serial Number __, filed __, the entire disclosure of which is hereby incorporated by reference.

2. Papers Enclosed Which Are Required For Filing Date Under 37 CFR 1.53(b) (Non-provisional)

164 pages of specification

13 pages of claims

1 page(s) of abstract

0 sheet(s) of drawings

☐ formal ☐ informal

09/684458 10/05/00

3. Declaration or Oath

(for new and CIP applications; also for Cont./Div. where inventor(s) are being added)

☒ An executed declaration of the inventor(s) ☐ is enclosed ☒ will follow.

(for Cont./Div. where inventorship is the same or inventor(s) being deleted)

☐ A copy of the executed declaration/oath filed in the prior application is enclosed (37 CFR 1.63(d)).

(for Cont./Div. where inventor(s) being deleted)

☐ A signed statement is attached deleting inventor(s) named in the prior application (see 37 CFR 1.63(d)(2) and 1.33(b)).

4. Assignment

(for new and CIP applications)

☒ An Assignment of the invention to GENENTECH, INC. ☐ is enclosed with attached Recordation Form Cover Sheet ☒ will follow.

(for cont./div.)

☐ The prior application is assigned of record to Genentech, Inc.

5. Amendments (for continuation and divisional applications)

☐ Cancel in this application original claims __ of the prior application before calculating the filing fee. (At least one original independent claim must be retained for filing purposes.)

☐ A preliminary amendment is enclosed. (Claims added by this amendment have been properly numbered consecutively beginning with the number next following the highest numbered original claim in the prior application.)

Relate Back -- 35 U.S.C. 120 or 35 U.S.C. 119

☒ Amend the specification by inserting before the first line the sentence:

--This is a

☒ non-provisional application
☐ continuation
☐ divisional
☐ continuation-in-part

of co-pending application(s)

☐ Serial No. _ filed on __, which application(s) is(are) incorporated herein by reference and to which application(s) priority is claimed under 35 USC §120. --

☐ International Application _ filed on __ which designated the U.S., which application(s) is(are) incorporated herein by reference and to which application(s) priority is claimed under 35 USC §120.--

X provisional application Nos. 60/158,587 and 60/162,611 filed October 7, 1999 and October 28, 1999, the entire disclosure of which is hereby incorporated by reference and to which application(s) priority is claimed under 35 USC §119.--.

6. Fee Calculation (37 CFR 1.16)

The fee has been calculated as follows:

CLAIMS FOR FEE CALCULATION					
Number Filed		Number Extra		Rate	Basic Fee 37 CFR 1.16(a)
					\$710.00
Total Claims	112	- 20 =	92	X \$18.00	\$1,656.00
Independent Claims	42	- 3 =	39	X \$80.00	\$3,042.00
Multiple dependent claim(s), if any				+ \$270.00	\$0.00
Filing Fee Calculation					\$5,486.00

7. Method of Payment of Fees

The Commissioner is hereby authorized to charge Deposit Account No. 07-0630 in the amount of **\$5,486.00**. **A duplicate copy of this transmittal is enclosed.**

8. Authorization to Charge Additional Fees

The Commissioner is hereby authorized to charge any additional fees required under 37 CFR §1.16 and 1.17, or credit overpayment to Deposit Account No. 07-0630. **A duplicate copy of this sheet is enclosed.**

9. Additional Papers Enclosed

- ☐ Information Disclosure Statement (37 CFR §1.98) w/ PTO-1449 and citations
- ☒ Submission of "Sequence Listing", computer readable copy, certificate re: sequence listing, and/or amendment pertaining thereto for biological invention containing nucleotide and/or amino acid sequence.
- ☐ A new Power of Attorney or authorization of agent.
- ☐ Other:

10. Maintenance of Copendency of Prior Application (for continuation and divisional applications)

*[This item **must** be completed and the necessary papers filed in the prior application if the period set in the prior application has run]*

- _____ A petition, fee and/or response has been filed to extend the term in the pending prior application until _____
- _____ A copy of the petition for extension of time in the **prior** application is attached.


11. Correspondence Address:

X Address all future communications to:

GENENTECH, INC.
Attn: Timothy R. Schwartz, Ph.D.
1 DNA Way
South San Francisco, California 94080-4990
(650) 225-7467

Respectfully submitted,
GENENTECH, INC.

Date: October 5, 2000

By: 
Timothy R. Schwartz, Ph.D.
Reg. No. 32,171

1 DNA Way
South San Francisco, CA 94080-4990
Phone: (650) 225-7467
Fax: (650) 952-9881

005007 3643650

**NOVEL POLYPEPTIDES, THEIR NUCLEIC ACIDS, AND METHODS FOR THEIR USE IN
ANGIOGENESIS AND VASCULARIZATION**

FIELD OF THE INVENTION

The present invention relates generally to the identification and isolation of novel DNA and their encoded intracellular polypeptides designated herein as "PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72" polypeptides, whose gene expression is modulated in cells undergoing angiogenesis and/or vascularization. Accordingly, the present invention further relates to compositions and methods useful for promoting or inhibiting angiogenesis and/or neo- or cardio-vascularization in mammals in need of such biological effect. This includes the diagnosis and treatment of cardiovascular disorders as well as oncological disorders.

BACKGROUND OF THE INVENTION

Intracellular proteins play important roles in, among other things, the formation, differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration, differentiation, or interaction with other cells, is typically governed by information received from other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and hormones) which are, in turn, recognized by and activate diverse cell receptors or membrane-bound proteins. Each activation signal initiates a specific, signal transduction pathway composed of intracellular proteins (e.g., protein kinases, DNA-binding regulatory proteins, protein processing proteins, proteases, glycosidases) resulting in the modulation, either up- or down-regulation, of the activity, expression, or amount of other intracellular proteins involved in or necessary for the cell's fate in response to the signal. For example, detectable changes in the RNA or protein levels of intracellular proteins necessary for cell growth or differentiation in response to appropriate transduction of signals can be controlled in part by receptor-mediated phosphorylation of signal-induction-pathway related intracellular proteins.

Intracellular proteins and their gene sequences have various industrial applications, including as drug targets for pharmaceuticals, diagnostics, pharmaceuticals, biosensors, and bioreactors. While most protein drugs available at present are secreted cytokines or their antibody mimics, most targets of small molecule, peptide, or antisense drugs are intracellular proteins or the intracellular genes that encode them. For example, such drugs can interact with an intracellular protein target to block its activity and disrupt the related signal transduction pathway, thereby stopping (or modulating) the cell's response or activity controlled by that pathway. Both industry and academia are undertaking efforts to identify new, native intracellular proteins and their genes, the signal transduction pathways in which they function, and the proteins or genes they modulate. Classically, such genes and their proteins are discovered by binary comparison studies in which a differential analysis is made of RNA or protein upon a cell or tissue response to a certain stimulus.

One consequence of cellular response is the formation of new blood vessels, which can occur by two

related mechanisms: angiogenesis--the growth of new vessels from pre-existing vessels--and vasculogenesis--the formation of vessels through aggregation of endothelial cells. All blood vessel inner surfaces are lined with endothelial cells. Vascular endothelial cells, at the interface between blood and extravascular space, play prominent roles in maintaining cardiovascular homeostasis and mediate pathophysiologic responses to injury.

5 For example, angiogenesis occurs in the adult during events such as wound healing and ovulation. During angiogenesis, endothelial cells responding to environmental stimuli undergo a number of cellular alterations and responses, resulting in a complex series of steps, which involve degradation of the basement membrane by cellular proteases, penetration and migration of endothelial cells into the extracellular matrix, endothelial proliferation, and the formation of interconnected vascular networks. This formation of new vessels takes place
10 in distinct phases that entails and relies upon modulation or expression of a variety of intracellular proteins, extracellular matrix components, proteases and protease inhibitors, inflammatory molecules, chemokines, and molecules involved in cell division and proliferation, cytoskeletal rearrangement, adhesion molecules and also apoptosis of certain endothelial cell populations.

Endothelial cells also undergo angiogenesis during the neovascularization associated with tumor
15 growth and metastasis and a variety of non-neoplastic diseases or disorders. In the case of tumor growth, angiogenesis appears to be crucial for the transition from hyperplasia to neoplasia, and for providing nourishment to the growing solid tumor (Folkman, *et al.*, *Nature* 339:58 (1989)). Angiogenesis allows tumors to be in contact with the vascular bed of the host, which provides a route for metastasis of the tumor cells. In fact, the progression of solid tumor growth and metastasis depends on angiogenesis, as supported for example,
20 by studies showing a correlation between the number and density of microvessels in histologic sections of invasive human breast carcinoma and actual presence of distant metastases (Weidner, *et al.*, *New Engl. J. Med.*, 324:1 (1991)). Recent data suggests that blocking new blood vessel growth can slow tumor growth by cutting off the supply of oxygen and nutrients; without a new blood supply tumors cannot grow more than about 1-2 mm in diameter. Thus new angiostatic therapies to treat cancer are desired.

25 There exists a need for additional products, methods and assays that provide a means to control signal transduction pathways and thereby modulate cellular and tissue response and activity. Such products, methods and assays will provide benefit in numerous medical conditions and procedures.

In view of the role of vascular endothelial cell growth and angiogenesis in many diseases and disorders, it is desirable to have a means of modulating one or more of the biological effects causing these processes, in
30 order to provide benefits such as enhancing repair or maintenance of blood vessels and reducing or inhibiting cancer and tumor progression. It is also desirable to have a means of assaying for the presence of pathogenic polypeptides in normal and diseased conditions, and especially cancer. Further, as there is no generally applicable therapy for the treatment of cardiac hypertrophy, the identification of factors that can prevent or reduce cardiac myocyte hypertrophy is of primary importance in the development of new therapeutic strategies
35 to inhibit pathophysiological cardiac growth.

While there are several treatment modalities for various cardiovascular and oncologic disorders, there is still a need for additional therapeutic approaches. As a further means to address these existing needs, the identification and characterization of novel intracellular polypeptides designated herein as "PRO-C-MG.2, PRO-

C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72" polypeptides are provided.

SUMMARY OF THE INVENTION

cDNA clones, designated herein as DNA-C-MG.2-1776, DNA-C-MG.12-1776, DNA-C-MG.45-1776, DNA-C-MG.64-1776 or DNA-C-MG.72-1776, that encode a novel polypeptide designated in the present application as "PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72", respectively, have been identified, whose DNA-C-MG.2-1776, DNA-C-MG.12-1776, DNA-C-MG.45-1776, DNA-C-MG.64-1776 or DNA-C-MG.72-1776 RNA is modulated in cells undergoing tube formation by endothelial cells, which is a necessary step in the development of a blood vessel during angiogenesis and vasculogenesis. Differential cDNA screening, GeneCalling™ technology, was applied to human umbilical cord endothelial cells (HUVECS) undergoing tube formation in collagen gels in the presence of growth factors, mimicking the angiogenic environment of endothelial cells in vivo. The three dimensional gel is pre-requisite for the differentiation and fusion of endothelial cells into tubes; HUVECS grown on the surface of gelatin or on plastic do not undergo tube-formation.

Accordingly, the present invention concerns compositions and methods for promoting or inhibiting angiogenesis and/or vascularization, preferably neo- or cardio-vascularization in mammals, and for identifying additional molecules providing that benefit. The molecules of the present invention are believed to be useful drugs for the diagnosis and/or treatment (including prevention) of disorders where such effects are desired, such as the promotion or inhibition of angiogenesis, inhibition or stimulation of vascular endothelial cell growth, stimulation of growth or proliferation of vascular endothelial cells, inhibition of tumor growth, inhibition of angiogenesis-dependent tissue growth, stimulation of angiogenesis-dependent tissue growth, inhibition of cardiac hypertrophy and stimulation of cardiac hypertrophy, *e.g.*, for the treatment of congestive heart failure. The present invention provides methods for promoting or inhibiting angiogenesis by supplying to endothelial tissue an effective amount of a compound of the invention. Also provided are methods for treating a tumor, reducing the size of a tumor, reducing the vasculature supporting a tumor or reducing the tumor burden of a mammal by administering an effective amount of a compound of the invention.

In one embodiment, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence that encodes a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

In one embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% sequence identity, with increasing preference for each one percent increase in sequence identity, to at least about 99% sequence identity to (a) a DNA molecule encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide having the sequence of amino acid residues from about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively, or (b) the complement of the DNA molecule of (a), or a DNA molecule encoding the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide encoded by the ATCC-deposited DNA of the invention as described herein.

In another embodiment, the isolated nucleic acid molecule comprises (a) a nucleotide sequence encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide having the sequence of amino acid residues from about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively, or (b) the complement of the nucleotide sequence of (a), or the ATCC-deposited DNA encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

In other embodiments, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% sequence identity, with increasing preference for each one percent increase in sequence identity, to at least about 99% sequence identity to (a) a DNA molecule having the sequence of nucleotides from about 66 to about 1796 of SEQ ID NO:1, about 465 to about 1886 of SEQ ID NO:3, about 271 to about 1788 of SEQ ID NO:17, about 267 to about 1298 of SEQ ID NO:15, or about 71 to about 2059 of SEQ ID NO:13, respectively, (b) the complement of the DNA molecule of (a), or the ATCC-deposited DNA encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

In another embodiment, the isolated nucleic acid molecule comprises (a) the nucleotide sequence of from about 66 to about 1796 of SEQ ID NO:1, about 465 to about 1886 of SEQ ID NO:3, about 271 to about 1788 of SEQ ID NO:17, about 267 to about 1298 of SEQ ID NO:15, or about 71 to about 2059 of SEQ ID NO:13, respectively, or (b) the complement of the nucleotide sequence of (a), or the deposited DNA encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

In another embodiment, the invention concerns an isolated nucleic acid molecule which encodes an active PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide as defined herein comprising a nucleotide sequence that hybridizes to the complement of a nucleic acid sequence that encodes amino acids about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively. Preferably, hybridization occurs under stringent hybridization and wash conditions.

In yet another embodiment, the invention concerns an isolated nucleic acid molecule which encodes an active PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide as defined herein comprising a nucleotide sequence that hybridizes to the complement of the nucleic acid sequence between about 66 to about 1796 of SEQ ID NO:1, about 465 to about 1886 of SEQ ID NO:3, about 271 to about 1788 of SEQ ID NO:17, about 267 to about 1298 of SEQ ID NO:15, or about 71 to about 2059 of SEQ ID NO:13, respectively. Preferably, hybridization occurs under stringent hybridization and wash conditions.

In a further embodiment, the invention concerns an isolated nucleic acid molecule which is produced by hybridizing a test DNA molecule under stringent conditions with (a) a DNA molecule encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide having the sequence of amino acid residues from about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively, or (b) the complement of the DNA molecule of (a), and, if the test DNA molecule has at least about an 80% sequence identity, with increasing preference for each one percent increase in sequence identity,

to at least about 99% sequence identity to (a) or (b), and isolating the test DNA molecule. Such a molecule, hybridizable to the DNA molecule encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide having the sequence of amino acid residues from about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively, have at least about 596 and 1535 nucleotides, respectively.

In another embodiment, the invention concerns an isolated PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 nucleic acid molecule comprising (a) a nucleotide sequence encoding a polypeptide scoring at least about 80% positives, with increasing preference for each one percent increase in positives, to at least about 99% positives when compared with the amino acid sequence of residues about 1 to about 577 of SEQ ID NO:2 or about 1 to about 474 of SEQ ID NO:4, respectively, or (b) the complement of the nucleotide sequence of (a).

Another embodiment is directed to fragments of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide coding sequence that can find use as, for example, hybridization probes or for encoding fragments of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide that can optionally encode a polypeptide comprising a binding site for an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 binding target, preferably an antibody, a natural PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 intracellular binding target, or a nonnatural binding agent. Such nucleic acid fragments are usually at least about 20 nucleotides in length with increasing preference to at least about 1000 nucleotides in length, wherein in this context the term "about" means the referenced nucleotide sequence length plus or minus 10% of that referenced length. In a preferred embodiment, the nucleotide sequence fragment is derived from any coding region of the nucleotide sequence shown in SEQ ID NO:1 or SEQ ID NO:3. Also contemplated are the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide fragments encoded by these nucleotide molecule fragments, preferably those PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide fragments that comprise a binding site for an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 binding target, preferably an antibody, a natural PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 intracellular binding target, or a nonnatural binding agent.

In another embodiment, the invention provides a vector comprising a nucleotide sequence encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 or variants thereof. The vector can comprise any of the isolated nucleic acid molecules identified herein.

A host cell comprising such a vector is also provided. The host cells can be vertebrate, mammalian, fungal, plant, or bacterial cells. Preferred are yeast cells, CHO cells, *E. coli*, yeast, human or mouse cells. A process for producing PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides is further provided and comprises culturing host cells under conditions suitable for expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 in order to produce the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide. In a further

embodiment, the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide can be recovered from the cell culture. As used throughout, "cell culture" includes the cells or cell medium.

In another embodiment, the invention provides isolated PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide encoded by any of the isolated nucleic acid sequences identified
5 herein.

In a specific embodiment, the invention provides isolated native sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, which in certain embodiments, includes an amino acid sequence comprising residues from about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to
10 about 633 of SEQ ID NO: 14, respectively.

In another embodiment, the invention concerns an isolated PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, comprising an amino acid sequence having at least about 80% sequence identity, with increasing preference for each one percent increase in sequence identity, to at least about 99% sequence identity to the sequence of amino acid residues from about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ
15 ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively.

In a further embodiment, the invention concerns an isolated PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide comprising an amino acid sequence having at least about 80% sequence identity, with increasing preference for each one percent increase in sequence identity, to at least about 99% sequence identity to an amino acid sequence encoded by the human protein cDNA deposited with the ATCC as described herein.
20

In a further embodiment, the invention concerns an isolated PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide comprising an amino acid sequence scoring at least about 80% positives, with increasing preference for each one percent increase in positives, to at least about 99% positives when compared with the amino acid sequence of residues from about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ
25 ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively.

In yet another embodiment, the invention concerns an isolated PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, comprising the sequence of amino acid residues from about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively, or a fragment thereof which is biologically active or sufficient to provide a binding site for an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 binding target, preferably an antibody, a natural PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 intracellular binding
30 target, or a nonnatural binding agent, wherein the identification of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide fragments that possess biological activity or provide the binding site can be accomplished in a routine manner using techniques which are well known in the art.
35 Preferably, the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 fragment

retains a qualitative biological activity of a native PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

In a still further embodiment, the invention provides a polypeptide produced by (i) hybridizing a test DNA molecule under stringent conditions with (a) a DNA molecule encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide having the sequence of amino acid residues from about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively, or (b) the complement of the DNA molecule of (a), and if the test DNA molecule has at least about an 80% sequence identity, with increasing preference for each one percent increase in sequence identity, to at least about 99% sequence identity to (a) or (b), (ii) culturing a host cell comprising the test DNA molecule under conditions suitable for expression of the polypeptide in order to produce the polypeptide, and then optionally (iii) recovering the polypeptide from the cell culture.

In another embodiment, the invention provides chimeric molecules comprising a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide fused to a heterologous polypeptide or amino acid sequence, wherein the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide can comprise any PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, variant or fragment thereof as described herein. An example of such a chimeric molecule comprises a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide fused to an epitope tag sequence, a Fc region of an immunoglobulin, or a secretion signal peptide.

In one embodiment, the present invention provides a composition comprising a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide in admixture with a pharmaceutically acceptable carrier. In one aspect, the composition comprises a therapeutically effective amount of the polypeptide. In another aspect, the composition comprises a further active ingredient, namely, a cardiovascular, endothelial or angiogenic agent or an angiostatic agent, preferably an angiogenic or angiostatic agent. Preferably, the composition is sterile.

In a further embodiment, the present invention provides a method for preparing such a composition useful for the treatment of a cardiovascular, endothelial or angiogenic disorder comprising admixing a therapeutically effective amount of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide with a pharmaceutically acceptable carrier.

In another embodiment, the invention provides an antibody as defined herein which specifically binds to a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide as described herein. Optionally, the antibody is a monoclonal antibody, an antibody fragment or a single chain antibody.

In yet another embodiment, the invention concerns agonists and antagonists of a native PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide as defined herein. Preferably, the agonist or antagonist is a molecule that modulates PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 biological activity by acting at the post-translational, translational, transcriptional, or translocational level. In a particular embodiment the agonist or antagonist is an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody, an antigene molecule (sense or antisense),

a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 gene (e.g. for gene therapy) or a small molecule.

In one such embodiment are therapeutic PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 nucleic acids that are used to modulate cellular expression or intracellular concentration or availability of active PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. These nucleic acids include antigene compounds, more typically antisense: single-stranded sequences comprising complements of the disclosed PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 nucleic acids, and also include nucleic acid expressing PRO-C-MG.2 and PRO-C-MG.12 for gene therapy. Antigene modulation of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 expression can employ PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antisense nucleic acids operably linked to gene regulatory sequences. Cell are transfected with a vector comprising an PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 sequence with a promoter sequence oriented such that transcription of the gene yields an antisense transcript capable of binding to endogenous PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 encoding mRNA. Transcription of the antisense nucleic acid may be constitutive or inducible and the vector may provide for stable extrachromosomal maintenance or integration. In yet another embodiment, single-stranded antigene nucleic acids that bind to genomic DNA or RNA encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 are administered to the target cell, in or temporarily isolated from a host, at a concentration that results in a substantial reduction in PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 expression.

In one embodiment provided are PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 compounds that have one or more PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-specific binding affinities, including the ability to specifically bind at least one natural human intracellular PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-specific binding target or a binding agent such as a anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-specific antibody or agent identified in assays as described herein. Natural PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 binding targets are readily identified by screening cells, membranes and cellular extracts and fractions with the disclosed materials and methods. For example, two-hybrid screening using PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 fragments are used to identify intracellular targets which specifically bind such fragments.

In another embodiment, the present invention provides a composition comprising an agonist or antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide in admixture with a pharmaceutically acceptable carrier. In one aspect, the composition comprises a therapeutically effective amount of the agonist or antagonist. In another aspect, the composition comprises a further active ingredient, namely, a cardiovascular, endothelial or angiogenic agent or an angiostatic agent, preferably an angiogenic or angiostatic agent.

In a further embodiment, the present invention provides a method for preparing such a composition useful for the treatment of a cardiovascular, endothelial or angiogenic disorder comprising admixing a therapeutically

effective amount of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide agonist or antagonist with a pharmaceutically acceptable carrier.

In one embodiment, the invention provides efficient methods of identifying compounds active at the level of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 modulatable cellular function. Generally, these screening methods involve assaying for compounds which modulate a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 interaction with a natural PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 binding target. The methods are amenable to automated, cost-effective high throughput screening of chemical libraries for lead compounds. Assays for binding agents are provided including protein-protein binding assays, immunoassays, and cell based assays. A preferred assay is a high-through put cell-based or in vitro binding assay. For example, the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 compositions can be part of a fusion product with another peptide or polypeptide, e.g. a polypeptide that is capable of providing or enhancing protein-protein binding, stability under assay conditions, or a tag for detection or anchoring. The assay mixtures can contain a natural intracellular PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 binding target, or active portion thereof. The assay mixture can also contain a candidate pharmacological agent. The resultant mixture is incubated under conditions where, but for the presence of the candidate pharmacological agent, the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 specifically binds the cellular binding target, portion or analog with a reference binding affinity. A detected difference in the binding affinity of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 protein to the target in the absence of the agent as compared with the binding affinity in the presence of the agent indicates that the agent modulates the binding of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 protein to the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 binding target. Analogously, in a cell-based transcription assay, a difference in the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 transcriptional induction in the presence and absence of an agent indicates the agent modulates PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-induced transcription. A difference, as used herein, is statistically significant and preferably represents at least a 50%, more preferably at least a 90% difference.

In a further embodiment, the invention concerns a method of identifying agonists or antagonists to a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide which comprises contacting either the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, a cell comprising the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 nucleic acid with a candidate molecule and monitoring the specific binding to the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or nucleic acid and/or monitoring a biological activity mediated by the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide. Preferably, the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide is a native PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

In another embodiment, the present invention provides a method for identifying an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide comprising: (a) contacting target cells and a test compound to be screened under conditions suitable for the induction, stimulation or dependence of a cellular response normally induced by, stimulated by or dependent on a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide; and (b) determining the induction, stimulation or dependence of the cellular response to determine if the test compound is an effective agonist, wherein the induction or enhancement of the cellular response is indicative of the test compound being an effective agonist. In a preferred embodiment, the target cells have been engineered or treated to prevent expressing endogenous PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 during the test period. The cellular response is preferably cell proliferation or tube formation.

In another embodiment, the present invention provides a method for identifying an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide comprising: (a) contacting target cells and a test compound to be screened under conditions suitable for the induction, stimulation or dependence of a cellular response normally induced by, stimulated by or dependent on a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide; and (b) determining the induction, stimulation or dependence of the cellular response to determine if the test compound is an effective agonist, wherein the induction or enhancement of the cellular response is indicative of the test compound being an effective agonist. In a preferred embodiment, the target cells have been engineered or treated to prevent expressing endogenous PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 during the test period. The cellular response is preferably cell proliferation or tube formation.

In another embodiment, the invention provides a method for identifying a compound that inhibits the activity of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide comprising contacting a test compound with a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide under conditions and for a time sufficient to allow the test compound and polypeptide to interact and determining whether the activity of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide is inhibited. In a specific preferred embodiment, either the test compound or the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide is immobilized on a solid support. In another preferred aspect, the non-immobilized component carries a detectable label. In a preferred aspect, this method comprises the steps of: (a) contacting cells and a test compound to be screened in the presence of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide under conditions suitable for the induction, stimulation, or dependence of a cellular response on a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide; and (b) determining the induction, stimulation or dependence of the cellular response to determine if the test compound is an effective antagonist. In another preferred aspect, this process comprises the steps of: (a) contacting cells and a test compound to be screened in the presence of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide under conditions suitable for the stimulation or dependence of cell proliferation or tube formation on a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide; and (b) measuring the cell proliferation or tube formation to determine if

the test compound is an effective antagonist.

In another embodiment, the invention provides a method for identifying a compound that inhibits the expression of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide in cells that normally expresses the polypeptide, wherein the method comprises contacting the cells with a test compound and determining whether the expression of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide is inhibited. In a preferred aspect, this method comprises the steps of: (a) contacting cells and a test compound to be screened under conditions suitable for allowing expression of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide; and (b) determining the inhibition of expression of said polypeptide.

In a still further embodiment, the invention provides a compound that inhibits the expression of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, such as a compound that is identified by the methods set forth above.

Another aspect of the present invention is directed to an agonist or an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide which can optionally be identified by the methods described above.

The invention also provides a microarray that comprises a polynucleotide sequence encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, optionally with a portion of the 5' or 3' untranslated sequence of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 gene or mRNA.

In a still further embodiment, the invention concerns a composition of matter comprising a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, nucleic acid, or agonist or antagonist thereof as herein described, preferably an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody or a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene molecule, in combination with a carrier. Optionally, the carrier is a pharmaceutically acceptable carrier.

Another embodiment of the present invention is directed to the use of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or nucleic acid or an agonist or antagonist thereof as herein described, preferably an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody or a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene molecule, for the preparation of a medicament useful in the treatment of a condition which is responsive to the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, nucleic acid, agonist or antagonist.

In another aspect, the present invention provides an article of manufacture comprising: (a) a composition of matter comprising a therapeutically effective dosage of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or nucleic acid or an agonist or antagonist thereof as herein described, preferably an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody or a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene molecule; (b) a container containing said composition; and optionally, (c) a label affixed to said container, or a

package insert included in said pharmaceutical product referring to the use of the compound in the treatment of a cardiovascular, endothelial or angiogenic disorder.

In a still further aspect, the present invention provides a method for diagnosing a disease or susceptibility to a disease which is related to a mutation in a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide-encoding nucleic acid sequence comprising: (a) isolating or amplifying a nucleic acid sequence encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide from a sample derived from a host; and (b) determining the presence or absence of said mutation in the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide nucleic acid sequence, wherein the presence or absence of said mutation is indicative of the presence of said disease or susceptibility to said disease.

In a still further aspect, the invention provides a method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal which comprises analyzing the level of expression of a gene encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide (a) in a test sample of tissue cells obtained from said mammal, and (b) in a control sample of known normal tissue cells of the same cell type, wherein a higher or lower expression level in the test sample as compared to the control sample is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in said mammal. The expression of a gene encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide may optionally be accomplished by measuring the level of mRNA or polypeptide in the test sample as compared to the control sample.

In a still further aspect, the present invention provides a method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal which comprises detecting the presence or absence of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide in a test sample of tissue cells obtained from said mammal, wherein the presence or absence of said PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide in said test sample is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in said mammal.

In a still further embodiment, the invention provides a method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal comprising (a) contacting an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody with a test sample of tissue cells obtained from the mammal, and (b) detecting the formation of a complex between the antibody and the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide in the test sample, wherein the formation of said complex is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in the mammal. The detection may be qualitative or quantitative, and may be performed in comparison with monitoring the complex formation in a control sample of known normal tissue cells of the same cell type. A larger or smaller quantity of complexes formed in the test sample indicates the presence of a cardiovascular, endothelial or angiogenic dysfunction in the mammal from which the test tissue cells were obtained. The antibody preferably carries a detectable label.

In another embodiment, the invention provides a method for determining the presence of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide in a sample comprising

exposing a sample suspected of containing the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide to an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody and determining binding of said antibody to a component of said sample.

In further aspects, the invention provides a cardiovascular, endothelial or angiogenic disorder diagnostic kit comprising an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody or nucleic acid, and a carrier, in suitable packaging. Preferably, such kit further comprises instructions for using said antibody or nucleic acid to detect the presence of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or nucleic acid. Preferably, the carrier is a buffer, for example. Preferably, the cardiovascular, endothelial or angiogenic disorder is cancer.

In a further embodiment, the invention provides an article of manufacture, comprising: a container; a composition comprising a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide contained within the container; and optionally, a label on the container, wherein the label on the container indicates that the composition can be used for treating cardiovascular, endothelial or angiogenic disorders.

In a further embodiment, the invention provides an article of manufacture, comprising: a container; a composition comprising a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide agonist or antagonist contained within the container; and optionally a label on the container; wherein the label on the container indicates that the composition can be used for treating cardiovascular, endothelial or angiogenic disorders.

In a further embodiment, the invention provides an article of manufacture, comprising: a container; a composition comprising an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody or antigene compound contained within the container; and optionally a label on the container; wherein the label on the container indicates that the composition can be used for treating cardiovascular, endothelial or angiogenic disorders.

In yet another embodiment, the present invention provides a method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal an effective amount of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide. Preferably, the disorder is cardiac hypertrophy, vascular trauma such as with wounds, burns, or surgery, or a type of cancer. In a further aspect, the mammal is further exposed to angioplasty or a drug that treats cardiovascular, endothelial or angiogenic disorders such as ACE inhibitors or chemotherapeutic agents if the cardiovascular, endothelial or angiogenic disorder is a type of cancer. Preferably, the mammal is human. Preferably it is one who is at risk of developing cardiac hypertrophy and more preferably has suffered myocardial infarction.

In another preferred embodiment, the cardiovascular, endothelial or angiogenic disorder is a cancer and the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide is administered in combination with a chemotherapeutic agent, a growth inhibitory agent or a cytotoxic agent.

In a further embodiment, the invention concerns a method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal an effective amount of an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide. Preferably,

the cardiovascular, endothelial or angiogenic disorder is cardiac hypertrophy or vascular trauma. Also preferred is where the mammal is human, and where an effective amount of an angiogenic agent is administered in conjunction with the agonist.

In a further embodiment, the invention concerns a method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal an effective amount of an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide. Preferably, the cardiovascular, endothelial or angiogenic disorder is a cancer or age-related macular degeneration. Also preferred is where the mammal is human, and where an effective amount of an angiostatic agent is administered in conjunction with the antagonist.

In a further embodiment, the invention concerns a method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal an effective amount of an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody or antigene compound. Preferably, the cardiovascular, endothelial or angiogenic disorder is cardiac hypertrophy, vascular trauma, a cancer, or age-related macular degeneration. Also preferred is where the mammal is human. Also preferred here is when an effective amount of an angiogenic or angiostatic agent is administered in conjunction with the antibody.

In still further embodiments, the invention provides a method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal that suffers therefrom comprising administering to the mammal a nucleic acid molecule that is an antigene compound or that codes for either (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide (b) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or (c) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, wherein said agonist or antagonist is preferably an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody. In a preferred embodiment the antigene compound is an antisense oligonucleotide, and more preferably a sense or antisense peptide nucleic acid. In a preferred embodiment, the mammal is human. In another preferred embodiment, the gene is administered via *ex vivo* gene therapy. In a further preferred embodiment, the gene is comprised within a vector, more preferably an adenoviral, adeno-associated viral, lentiviral, or retroviral vector.

In yet another aspect, the invention provides a recombinant retroviral particle comprising a retroviral vector consisting essentially of a promoter, a nucleic acid encoding (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) an agonist polypeptide of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (c) an antagonist polypeptide of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, and a signal sequence for cellular secretion of the polypeptide, wherein the retroviral vector is in association with retroviral structural proteins. Preferably, the signal sequence is from a mammal, such as from a native PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

In a still further embodiment, the invention supplies an *ex vivo* producer cell comprising a nucleic acid construct that expresses retroviral structural proteins and also comprises a retroviral vector consisting essentially of a promoter, nucleic acid encoding (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or

PRO-C-MG.72 polypeptide, (b) an agonist polypeptide of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or (c) an antagonist polypeptide of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, and a signal sequence for cellular secretion of the polypeptide, wherein said producer cell packages the retroviral vector in association with the structural proteins to produce recombinant retroviral particles.

In yet another embodiment, the invention provides a method for inhibiting endothelial cell growth in a mammal comprising administering to the mammal (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or (b) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, where the antagonist is preferably a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compound or a small molecule, and wherein endothelial cell growth in said mammal is inhibited. Preferably, the mammal is human, and the endothelial cell growth is associated with a tumor.

In yet another embodiment, the invention provides a method for stimulating endothelial cell growth in a mammal comprising administering to the mammal (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or (b) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, where preferably the agonist is a preferably a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compound, such as a PNA, or a small molecule, and wherein endothelial cell growth in the mammal is stimulated. Preferably, the mammal is human.

In yet another embodiment, the invention provides a method for inhibiting tube formation in a mammal comprising administering to the mammal (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or (b) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, wherein preferably the antagonist is a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compound or small molecule, and wherein tube formation in said mammal is inhibited.

In yet another embodiment, the invention provides a method for stimulating tube formation in a mammal comprising administering to the mammal (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or (b) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, where the agonist is preferably a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compound or small molecule, and wherein tube formation in said mammal is stimulated.

In yet another embodiment, the invention provides a method for inhibiting angiogenesis induced by, enhanced by or dependent on a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide in a mammal comprising administering to the mammal (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or (b) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, wherein preferably the antagonist is a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compound or small molecule, and wherein angiogenesis in the mammal is inhibited. Preferably, the mammal is a human, and more

preferably the mammal has a tumor.

In yet another embodiment, the invention provides a method for stimulating angiogenesis induced by, enhanced by, or dependent on a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide in a mammal comprising administering to the mammal (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or (b) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, where the agonist is preferably a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compound or small molecule, and wherein angiogenesis in the mammal is stimulated. Preferably, the mammal is a human.

BRIEF DESCRIPTION OF THE DRAWINGS

10 None.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

I. Definitions

The terms "PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide", "PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 protein" and "PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72" when used herein encompass native sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 and PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide variants (which are further defined herein). The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide can be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant and/or synthetic methods.

A "native sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72" comprises a polypeptide having the same amino acid sequence as a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 derived from nature. Such native sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be isolated from nature or can be produced by recombinant and/or synthetic means. The term "native sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72" specifically encompasses naturally-occurring truncated or secreted forms (*e.g.*, an extracellular domain sequence), naturally-occurring variant forms (*e.g.*, alternatively spliced forms) and naturally-occurring allelic variants of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. In one embodiment of the invention, the native sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 is a mature or full-length native sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 comprising amino acids about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively. Also, while the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide disclosed in SEQ ID NO:2 or SEQ ID NO:4, respectively, is shown to begin with the methionine residue designated herein as amino acid position 1, it is conceivable and possible that another methionine residue encoded by a start codon located either upstream or downstream from the codon of amino acid position 1 in SEQ ID NO:1 or SEQ ID NO:3 can be employed as the starting amino acid residue for the PRO-C-MG.2, PRO-C-MG.12, PRO-C-

MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

"PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 variant polypeptide"

means an active PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide as defined herein having at least about 80% amino acid sequence identity with the amino acid sequence of (a)

residues about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14,

respectively, or (b) another specifically derived fragment of the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:4, respectively. Such PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-

MG.72 variant polypeptides include, for instance, PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-

MG.64 or PRO-C-MG.72 polypeptides wherein one or more amino acid residues are added, or deleted, at the N- and/or C-terminus, as well as within one or more internal domains, of the sequence of SEQ ID NO:2 or SEQ ID NO:4, respectively. Ordinarily, a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-

MG.72 variant polypeptide will have at least about 80% amino acid sequence identity, more preferably at least about 81% amino acid sequence identity, more preferably at least about 82% amino acid sequence identity,

more preferably at least about 83% amino acid sequence identity, more preferably at least about 84% amino acid sequence identity, more preferably at least about 85% amino acid sequence identity, more preferably at least about 86% amino acid sequence identity, more preferably at least about 87% amino acid sequence identity,

more preferably at least about 88% amino acid sequence identity, more preferably at least about 89% amino acid sequence identity, more preferably at least about 90% amino acid sequence identity, more preferably at least

about 91% amino acid sequence identity, more preferably at least about 92% amino acid sequence identity,

more preferably at least about 93% amino acid sequence identity, more preferably at least about 94% amino acid sequence identity, more preferably at least about 95% amino acid sequence identity, more preferably at least

about 96% amino acid sequence identity, more preferably at least about 97% amino acid sequence identity, more preferably at least about 98% amino acid sequence identity and yet more preferably at least about 99%

amino acid sequence identity with (a) residues about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to

about 633 of SEQ ID NO: 14, respectively, or (b) another specifically derived fragment of the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:4, respectively. PRO-C-MG.2, PRO-C-MG.12, PRO-C-

MG.45, PRO-C-MG.64 or PRO-C-MG.72 variant polypeptides do not encompass the native PRO-C-MG.2,

PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide sequence. Ordinarily, PRO-C-

MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 variant polypeptides are at least about 10 amino acids in length, often at least about 20 amino acids in length, more often at least about 30 amino

acids in length, more often at least about 40 amino acids in length, more often at least about 50 amino acids in length, more often at least about 60 amino acids in length, more often at least about 70 amino acids in length,

more often at least about 80 amino acids in length, more often at least about 90 amino acids in length, more often at least about 100 amino acids in length, more often at least about 150 amino acids in length, more often at

least about 200 amino acids in length, more often at least about 250 amino acids in length, more often at least

about 300 amino acids in length, or more.

"Percent (%) amino acid sequence identity" with respect to the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are obtained as described below by using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Table 1 has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or can be compiled from the source code provided in Table 1. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

Table 1

```

/*
 *
 * C-C increased from 12 to 15
 * Z is average of EQ
 * B is average of ND
 * match with stop is _M; stop-stop = 0; J (joker) match = 0
 */
#define _M -8 /* value of a match with a stop */

int _day[26][26] = {
/* A B C D E F G H I J K L M N O P Q R S T U V W X Y Z */
/* A */ { 2, 0, -2, 0, 0, -4, 1, -1, -1, 0, -1, -2, -1, 0, _M, 1, 0, -2, 1, 1, 0, 0, -6, 0, -3, 0},
/* B */ { 0, 3, -4, 3, 2, -5, 0, 1, -2, 0, 0, -3, -2, 2, _M, -1, 1, 0, 0, 0, 0, -2, -5, 0, -3, 1},
/* C */ {-2, -4, 15, -5, -5, -4, -3, -3, -2, 0, -5, -6, -5, -4, _M, -3, -5, -4, 0, -2, 0, -2, -8, 0, 0, -5},
/* D */ { 0, 3, -5, 4, 3, -6, 1, 1, -2, 0, 0, -4, -3, 2, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 2},
/* E */ { 0, 2, -5, 3, 4, -5, 0, 1, -2, 0, 0, -3, -2, 1, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 3},
/* F */ {-4, -5, -4, -6, -5, 9, -5, -2, 1, 0, -5, 2, 0, -4, _M, -5, -5, -4, -3, -3, 0, -1, 0, 0, 7, -5},
/* G */ { 1, 0, -3, 1, 0, -5, 5, -2, -3, 0, -2, -4, -3, 0, _M, -1, -1, -3, 1, 0, 0, -1, -7, 0, -5, 0},
/* H */ {-1, 1, -3, 1, 1, -2, -2, 6, -2, 0, 0, -2, -2, 2, _M, 0, 3, 2, -1, -1, 0, -2, -3, 0, 0, 2},
/* I */ {-1, -2, -2, -2, -2, 1, -3, -2, 5, 0, -2, 2, 2, -2, _M, -2, -2, -2, -1, 0, 0, 4, -5, 0, -1, -2},
/* J */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* K */ {-1, 0, -5, 0, 0, -5, -2, 0, -2, 0, 5, -3, 0, 1, _M, -1, 1, 3, 0, 0, 0, -2, -3, 0, -4, 0},
/* L */ {-2, -3, -6, -4, -3, 2, -4, -2, 2, 0, -3, 6, 4, -3, _M, -3, -2, -3, -3, -1, 0, 2, -2, 0, -1, -2},
/* M */ {-1, -2, -5, -3, -2, 0, -3, -2, 2, 0, 0, 4, 6, -2, _M, -2, -1, 0, -2, -1, 0, 2, -4, 0, -2, -1},
/* N */ { 0, 2, -4, 2, 1, -4, 0, 2, -2, 0, 1, -3, -2, 2, _M, -1, 1, 0, 1, 0, 0, -2, -4, 0, -2, 1},
/* O */ { _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M},
0, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M},
/* P */ { 1, -1, -3, -1, -1, -5, -1, 0, -2, 0, -1, -3, -2, -1, _M, 6, 0, 0, 1, 0, 0, -1, -6, 0, -5, 0},
/* Q */ { 0, 1, -5, 2, 2, -5, -1, 3, -2, 0, 1, -2, -1, 1, _M, 0, 4, 1, -1, -1, 0, -2, -5, 0, -4, 3},
/* R */ {-2, 0, -4, -1, -1, -4, -3, 2, -2, 0, 3, -3, 0, 0, _M, 0, 1, 6, 0, -1, 0, -2, 2, 0, -4, 0},
/* S */ { 1, 0, 0, 0, 0, -3, 1, -1, -1, 0, 0, -3, -2, 1, _M, 1, -1, 0, 2, 1, 0, -1, -2, 0, -3, 0},
/* T */ { 1, 0, -2, 0, 0, -3, 0, -1, 0, 0, 0, -1, -1, 0, _M, 0, -1, -1, 1, 3, 0, 0, -5, 0, -3, 0},
/* U */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* V */ { 0, -2, -2, -2, -2, -1, -1, -2, 4, 0, -2, 2, 2, -2, _M, -1, -2, -2, -1, 0, 0, 4, -6, 0, -2, -2},
/* W */ {-6, -5, -8, -7, -7, 0, -7, -3, -5, 0, -3, -2, -4, -4, _M, -6, -5, 2, -2, -5, 0, -6, 17, 0, 0, -6},
/* X */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* Y */ {-3, -3, 0, -4, -4, 7, -5, 0, -1, 0, -4, -1, -2, -2, _M, -5, -4, -4, -3, -3, 0, -2, 0, 0, 10, -4},
/* Z */ { 0, 1, -5, 2, 3, -5, 0, 2, -2, 0, 0, -2, -1, 1, _M, 0, 3, 0, 0, 0, 0, -2, -6, 0, -4, 4}
};

```

00500 "00500" 00500

```

/*
 */
#include <stdio.h>
#include <ctype.h>

#define MAXJMP      16      /* max jumps in a diag */
#define MAXGAP      24      /* don't continue to penalize gaps larger than this */
#define JMPS        1024    /* max jmps in an path */
#define MX          4       /* save if there's at least MX-1 bases since last jmp */

#define DMAT        3       /* value of matching bases */
#define DMIS        0       /* penalty for mismatched bases */
#define DINS0       8       /* penalty for a gap */
#define DINS1       1       /* penalty per base */
#define PINS0       8       /* penalty for a gap */
#define PINS1       4       /* penalty per residue */

struct jmp {
    short          n[MAXJMP];      /* size of jmp (neg for dely) */
    unsigned short x[MAXJMP];      /* base no. of jmp in seq x */
};
/* limits seq to 2^16 -1 */

struct diag {
    int            score;           /* score at last jmp */
    long           offset;          /* offset of prev block */
    short          ijmp;            /* current jmp index */
    struct jmp     jp;              /* list of jmps */
};

struct path {
    int spc;           /* number of leading spaces */
    short n[JMPS];     /* size of jmp (gap) */
    int x[JMPS];       /* loc of jmp (last elem before gap) */
};

char    *ofile;           /* output file name */
char    *namex[2];        /* seq names: getseqs() */
char    *prog;            /* prog name for err msgs */
char    *seqx[2];         /* seqs: getseqs() */
int      dmax;            /* best diag: nw() */
int      dmax0;           /* final diag */
int      dna;             /* set if dna: main() */
int      endgaps;         /* set if penalizing end gaps */
int      gapx, gapy;       /* total gaps in seqs */
int      len0, len1;       /* seq lens */
int      ngapx, ngapy;     /* total size of gaps */
int      smax;            /* max score: nw() */
int      *xbm;            /* bitmap for matching */
long     offset;          /* current offset in jmp file */
struct   diag    *dx;      /* holds diagonals */
struct   path    pp[2];    /* holds path for seqs */

char      *calloc(), *malloc(), *index(), *strcpy();
char      *getseq(), *g_calloc();

```



```

/* Needleman-Wunsch alignment program
 *
 * usage: progs file1 file2
 * where file1 and file2 are two dna or two protein sequences.
 * The sequences can be in upper- or lower-case and may contain ambiguity
 * Any lines beginning with ';', '>' or '<' are ignored
 * Max file length is 65535 (limited by unsigned short x in the jmp struct)
 * A sequence with 1/3 or more of its elements ACGTU is assumed to be DNA
 * Output is in the file "align.out"
 *
 * The program may create a tmp file in /tmp to hold info about traceback.
 * Original version developed under BSD 4.3 on a vax 8650
 */
#include "nw.h"
#include "day.h"

static _dbval[26] = {
    1,14,2,13,0,0,4,11,0,0,12,0,3,15,0,0,0,5,6,8,8,7,9,0,10,0
};

static _pbval[26] = {
    1, 2|(1<<('D'-'A'))|(1<<('N'-'A')), 4, 8, 16, 32, 64,
    128, 256, 0xFFFFFFFF, 1<<10, 1<<11, 1<<12, 1<<13, 1<<14,
    1<<15, 1<<16, 1<<17, 1<<18, 1<<19, 1<<20, 1<<21, 1<<22,
    1<<23, 1<<24, 1<<25|(1<<('E'-'A'))|(1<<('Q'-'A'))
};

main(ac, av)
    int ac;
    char *av[];
{
    prog = av[0];
    if (ac != 3) {
        fprintf(stderr, "usage: %s file1 file2\n", prog);
        fprintf(stderr, "where file1 and file2 are two dna or two protein sequences.\n");
        fprintf(stderr, "The sequences can be in upper- or lower-case\n");
        fprintf(stderr, "Any lines beginning with ';' or '<' are ignored\n");
        fprintf(stderr, "Output is in the file \"align.out\"\n");
        exit(1);
    }
    namex[0] = av[1];
    namex[1] = av[2];
    seqx[0] = getseq(namex[0], &len0);
    seqx[1] = getseq(namex[1], &len1);
    xbm = (dna)? _dbval : _pbval;

    endgaps = 0; /* 1 to penalize endgaps */
    ofile = "align.out"; /* output file */

    nw(); /* fill in the matrix, get the possible jumps */
    readjumps(); /* get the actual jumps */
    print(); /* print stats, alignment */

    cleanup(0); /* unlink any tmp files */
}

```

main

```

/* do the alignment, return best score: main()
 * dna: values in Fitch and Smith, PNAS, 80, 1382-1386, 1983
 * pro: PAM 250 values
 * When scores are equal, we prefer mismatches to any gap, prefer
 * a new gap to extending an ongoing gap, and prefer a gap in seqx
 * to a gap in seq y.
 */
nw()
{
    char          *px, *py;          /* seqs and ptrs */
    int           *ndely, *dely;      /* keep track of dely */
    int           ndelx, delx;        /* keep track of delx */
    int           *tmp;               /* for swapping row0, row1 */
    int           mis;                /* score for each type */
    int           ins0, ins1;         /* insertion penalties */
    register      id;                 /* diagonal index */
    register      ij;                 /* jmp index */
    register      *col0, *col1;       /* score for curr, last row */
    register      xx, yy;             /* index into seqs */

    dx = (struct diag *)g_calloc("to get diags", len0+len1+1, sizeof(struct diag));

    ndely = (int *)g_calloc("to get ndely", len1+1, sizeof(int));
    dely = (int *)g_calloc("to get dely", len1+1, sizeof(int));
    col0 = (int *)g_calloc("to get col0", len1+1, sizeof(int));
    col1 = (int *)g_calloc("to get col1", len1+1, sizeof(int));
    ins0 = (dna)? DINS0 : PINS0;
    ins1 = (dna)? DINS1 : PINS1;

    smax = -10000;
    if (endgaps) {
        for (col0[0] = dely[0] = -ins0, yy = 1; yy <= len1; yy++) {
            col0[yy] = dely[yy] = col0[yy-1] - ins1;
            ndely[yy] = yy;
        }
        col0[0] = 0;          /* Waterman Bull Math Biol 84 */
    }
    else
        for (yy = 1; yy <= len1; yy++)
            dely[yy] = -ins0;

    /* fill in match matrix
     */
    for (px = seqx[0], xx = 1; xx <= len0; px++, xx++) {
        /* initialize first entry in col
         */
        if (endgaps) {
            if (xx == 1)
                col1[0] = delx = -(ins0+ins1);
            else
                col1[0] = delx = col0[0] - ins1;
            ndelx = xx;
        }
        else {
            col1[0] = 0;
            delx = -ins0;
            ndelx = 0;
        }
    }
}

```

nw

...nw

```

for (py = seqx[1], yy = 1; yy <= len1; py++, yy++) {
    mis = col0[yy-1];
    if (dna)
        mis += (xbm[*px-'A']&xbm[*py-'A'])? DMAT : DMIS;
    else
        mis += _day[*px-'A'][*py-'A'];

    /* update penalty for del in x seq;
     * favor new del over ongong del
     * ignore MAXGAP if weighting endgaps
     */
    if (endgaps || ndely[yy] < MAXGAP) {
        if (col0[yy] - ins0 >= dely[yy]) {
            dely[yy] = col0[yy] - (ins0+ins1);
            ndely[yy] = 1;
        } else {
            dely[yy] -= ins1;
            ndely[yy]++;
        }
    } else {
        if (col0[yy] - (ins0+ins1) >= dely[yy]) {
            dely[yy] = col0[yy] - (ins0+ins1);
            ndely[yy] = 1;
        } else
            ndely[yy]++;
    }

    /* update penalty for del in y seq;
     * favor new del over ongong del
     */
    if (endgaps || ndelx < MAXGAP) {
        if (col1[yy-1] - ins0 >= delx) {
            delx = col1[yy-1] - (ins0+ins1);
            ndelx = 1;
        } else {
            delx -= ins1;
            ndelx++;
        }
    } else {
        if (col1[yy-1] - (ins0+ins1) >= delx) {
            delx = col1[yy-1] - (ins0+ins1);
            ndelx = 1;
        } else
            ndelx++;
    }

    /* pick the maximum score; we're favoring
     * mis over any del and delx over dely
     */
}

```

```

id = xx - yy + len1 - 1;
if (mis >= delx && mis >= dely[yy])
    coll[yy] = mis;
else if (delx >= dely[yy]) {
    coll[yy] = delx;
    ij = dx[id].ijmp;
    if (dx[id].jp.n[0] && (!dna || (ndelx >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writejumps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
    }
    dx[id].jp.n[ij] = ndelx;
    dx[id].jp.x[ij] = xx;
    dx[id].score = delx;
}
else {
    coll[yy] = dely[yy];
    ij = dx[id].ijmp;

if (dx[id].jp.n[0] && (!dna || (ndely[yy] >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
    dx[id].ijmp++;
    if (++ij >= MAXJMP) {
        writejumps(id);
        ij = dx[id].ijmp = 0;
        dx[id].offset = offset;
        offset += sizeof(struct jmp) + sizeof(offset);
    }
}
    dx[id].jp.n[ij] = -ndely[yy];
    dx[id].jp.x[ij] = xx;
    dx[id].score = dely[yy];
}
if (xx == len0 && yy < len1) {
    /* last col
    */
    if (endgaps)
        coll[yy] -= ins0 + ins1*(len1-yy);
    if (coll[yy] > smax) {
        smax = coll[yy];
        dmax = id;
    }
}
}
if (endgaps && xx < len0)
    coll[yy-1] -= ins0 + ins1*(len0-xx);
if (coll[yy-1] > smax) {
    smax = coll[yy-1];
    dmax = id;
}
tmp = col0; col0 = coll; coll = tmp;
}
(void) free((char *)ndely);
(void) free((char *)dely);
(void) free((char *)col0);
(void) free((char *)coll);
}

```

```

/*
 *
 * print() -- only routine visible outside this module
 *
 * static:
 * getmat() -- trace back best path, count matches: print()
 * pr_align() -- print alignment of described in array p[]: print()
 * dumpblock() -- dump a block of lines with numbers, stars: pr_align()
 * nums() -- put out a number line: dumpblock()
 * putline() -- put out a line (name, [num], seq, [num]): dumpblock()
 * stars() -- put a line of stars: dumpblock()
 * stripname() -- strip any path and prefix from a seqname
 */

#include "nw.h"

#define SPC      3
#define P_LINE  256    /* maximum output line */
#define P_SPC    3      /* space between name or num and seq */

extern __day[26][26];
int olen;              /* set output line length */
FILE *fx;              /* output file */

print()
{
    int lx, ly, firstgap, lastgap;    /* overlap */

    if ((fx = fopen(ofile, "w")) == 0) {
        fprintf(stderr, "%s: can't write %s\n", prog, ofile);
        cleanup(1);
    }
    fprintf(fx, "<first sequence: %s (length = %d)\n", namex[0], len0);
    fprintf(fx, "<second sequence: %s (length = %d)\n", namex[1], len1);
    olen = 60;
    lx = len0;
    ly = len1;
    firstgap = lastgap = 0;
    if (dmax < len1 - 1) { /* leading gap in x */
        pp[0].spc = firstgap = len1 - dmax - 1;
        ly -= pp[0].spc;
    }
    else if (dmax > len1 - 1) { /* leading gap in y */
        pp[1].spc = firstgap = dmax - (len1 - 1);
        lx -= pp[1].spc;
    }
    if (dmax0 < len0 - 1) { /* trailing gap in x */
        lastgap = len0 - dmax0 - 1;
        lx -= lastgap;
    }
    else if (dmax0 > len0 - 1) { /* trailing gap in y */
        lastgap = dmax0 - (len0 - 1);
        ly -= lastgap;
    }
    getmat(lx, ly, firstgap, lastgap);
    pr_align();
}

```

```

/*
 * trace back the best path, count matches
 */
static
getmat(lx, ly, firstgap, lastgap)
    int lx, ly;          /* "core" (minus endgaps) */
    int firstgap, lastgap; /* leading trailing overlap */
{
    int nm, i0, i1, siz0, siz1;
    char outx[32];
    double pct;
    register n0, n1;
    register char *p0, *p1;

    /* get total matches, score
    */
    i0 = i1 = siz0 = siz1 = 0;
    p0 = seqx[0] + pp[1].spc;
    p1 = seqx[1] + pp[0].spc;
    n0 = pp[1].spc + 1;
    n1 = pp[0].spc + 1;

    nm = 0;
    while ( *p0 && *p1 ) {
        if (siz0) {
            p1++;
            n1++;
            siz0--;
        }
        else if (siz1) {
            p0++;
            n0++;
            siz1--;
        }
        else {
            if (xbm[*p0-'A'] & xbm[*p1-'A'])
                nm++;
            if (n0++ == pp[0].x[i0])
                siz0 = pp[0].n[i0++];
            if (n1++ == pp[1].x[i1])
                siz1 = pp[1].n[i1++];
            p0++;
            p1++;
        }
    }

    /* pct homology:
    * if penalizing endgaps, base is the shorter seq
    * else, knock off overhangs and take shorter core
    */
    if (endgaps)
        lx = (len0 < len1)? len0 : len1;
    else
        lx = (lx < ly)? lx : ly;
    pct = 100.*((double)nm)/((double)lx);
    fprintf(fx, "\n");
    fprintf(fx, " < %d match%s in an overlap of %d: %.2f percent similarity\n",
        nm, (nm == 1)? "" : "es", lx, pct);
}

```

getmat

...getmat

```

fprintf(fx, "< gaps in first sequence: %d", gapx);
if (gapx) {
    (void) sprintf(outx, " (%d %s%s)",
        ngapx, (dna)? "base": "residue", (ngapx == 1)? "" : "s");
    fprintf(fx, "%s", outx);

    fprintf(fx, ", gaps in second sequence: %d", gapy);
    if (gapy) {
        (void) sprintf(outx, " (%d %s%s)",
            ngapy, (dna)? "base": "residue", (ngapy == 1)? "" : "s");
        fprintf(fx, "%s", outx);
    }
    if (dna)
        fprintf(fx,
            "\n< score: %d (match = %d, mismatch = %d, gap penalty = %d + %d per base)\n",
            smax, DMAT, DMIS, DINS0, DINS1);
    else
        fprintf(fx,
            "\n< score: %d (Dayhoff PAM 250 matrix, gap penalty = %d + %d per residue)\n",
            smax, PINS0, PINS1);
    if (endgaps)
        fprintf(fx,
            "< endgaps penalized. left endgap: %d %s%s, right endgap: %d %s%s\n",
            firstgap, (dna)? "base" : "residue", (firstgap == 1)? "" : "s",
            lastgap, (dna)? "base" : "residue", (lastgap == 1)? "" : "s");
    else
        fprintf(fx, "< endgaps not penalized\n");
}

static      nm;          /* matches in core -- for checking */
static      lmax;        /* lengths of stripped file names */
static      ij[2];       /* jmp index for a path */
static      nc[2];       /* number at start of current line */
static      ni[2];       /* current elem number -- for gapping */
static      siz[2];
static char *ps[2];      /* ptr to current element */
static char *po[2];      /* ptr to next output char slot */
static char out[2][P_LINE]; /* output line */
static char star[P_LINE]; /* set by stars() */

/*
 * print alignment of described in struct path pp[]
 */
static
pr_align()
{
    int      nn;          /* char count */
    int      more;
    register i;

    for (i = 0, lmax = 0; i < 2; i++) {
        nn = stripname(name[i]);
        if (nn > lmax)
            lmax = nn;

        nc[i] = 1;
        ni[i] = 1;
        siz[i] = ij[i] = 0;
        ps[i] = seqx[i];
        po[i] = out[i];
    }
}

```

pr_align

...pr_align

```

for (nn = nm = 0, more = 1; more; ) {
    for (i = more = 0; i < 2; i++) {
        /*
         * do we have more of this sequence?
         */
        if (!*ps[i])
            continue;

        more++;

        if (pp[i].spc) { /* leading space */
            *po[i]++ = ' ';
            pp[i].spc--;
        }
        else if (siz[i]) { /* in a gap */
            *po[i]++ = '-';
            siz[i]--;
        }
        else { /* we're putting a seq element
            */
            *po[i] = *ps[i];
            if (islower(*ps[i]))
                *ps[i] = toupper(*ps[i]);
            po[i]++;
            ps[i]++;

            /*
             * are we at next gap for this seq?
             */
            if (ni[i] == pp[i].x[ij[i]]) {
                /*
                 * we need to merge all gaps
                 * at this location
                 */
                siz[i] = pp[i].n[ij[i]] + 1;
                while (ni[i] == pp[i].x[ij[i]])
                    siz[i] += pp[i].n[ij[i]] + 1;
            }
            ni[i]++;
        }
    }
    if (++nn == olen || !more && nn) {
        dumpblock();
        for (i = 0; i < 2; i++)
            po[i] = out[i];
        nn = 0;
    }
}

/*
 * dump a block of lines, including numbers, stars: pr_align()
 */
static
dumpblock()
{
    register    i;

    for (i = 0; i < 2; i++)
        *po[i]-- = '\0';
}

```

dumpblock

...dumpblock

```

(void) putc('\n', fx);
for (i = 0; i < 2; i++) {
    if (*out[i] && (*out[i] != ' ' || *(po[i]) != ' ')) {
        if (i == 0)
            nums(i);
        if (i == 0 && *out[1])
            stars();
        putline(i);
        if (i == 0 && *out[1])
            fprintf(fx, star);
        if (i == 1)
            nums(i);
    }
}
}

```

```

/*
 * put out a number line: dumpblock()
 */

```

static

nums(ix)

nums

```

    int ix;          /* index in out[] holding seq line */
{
    char          nline[P_LINE];
    register      i, j;
    register char *pn, *px, *py;

    for (pn = nline, i = 0; i < lmax + P_SPC; i++, pn++)
        *pn = ' ';
    for (i = nc[ix], py = out[ix]; *py; py++, pn++) {
        if (*py == ' ' || *py == '-')
            *pn = ' ';
        else {
            if (i%10 == 0 || (i == 1 && nc[ix] != 1)) {
                j = (i < 0)? -i : i;
                for (px = pn; j /= 10, px--)
                    *px = j%10 + '0';
                if (i < 0)
                    *px = '-';
            }
            else
                *pn = ' ';
            i++;
        }
    }
    *pn = '\0';
    nc[ix] = i;
    for (pn = nline; *pn; pn++)
        (void) putc(*pn, fx);
    (void) putc('\n', fx);
}

```

```

/*
 * put out a line (name, [num], seq, [num]): dumpblock()
 */

```

static

putline(ix)

putline

```

    int ix;
{

```

...putline

```

int          i;
register char *px;

for (px = namex[ix], i = 0; *px && *px != ':'; px++, i++)
    (void) putc(*px, fx);
for (; i < lmax+P_SPC; i++)
    (void) putc(' ', fx);

/* these count from 1:
 * ni[] is current element (from 1)
 * nc[] is number at start of current line
 */
for (px = out[ix]; *px; px++)
    (void) putc(*px&0x7F, fx);
(void) putc('\n', fx);
}

```

```

/*
 * put a line of stars (seqs always in out[0], out[1]): dumpblock()
 */

```

static

stars()

stars

```

{
    int          i;
    register char *p0, *p1, cx, *px;

    if (!*out[0] || (*out[0] == ' ' && *(p0[0]) == ' ') ||
        !*out[1] || (*out[1] == ' ' && *(p0[1]) == ' '))
        return;
    px = star;
    for (i = lmax+P_SPC; i; i--)
        *px++ = ' ';

    for (p0 = out[0], p1 = out[1]; *p0 && *p1; p0++, p1++) {
        if (isalpha(*p0) && isalpha(*p1)) {
            if (xbm[*p0-'A']&xbm[*p1-'A']) {
                cx = '*';
                nm++;
            }
            else if (!dna && _day[*p0-'A'][*p1-'A'] > 0)
                cx = '.';
            else
                cx = ' ';
        }
        else
            cx = ' ';
        *px++ = cx;
    }
    *px++ = '\n';
    *px = '\0';
}

```

```

/*
 * strip path or prefix from pn, return len: pr_align()
 */
static
stripname(pn)
char *pn; /* file name (may be path) */
{
    register char *px, *py;

    py = 0;
    for (px = pn; *px; px++)
        if (*px == '/')
            py = px + 1;
    if (py)
        (void) strcpy(pn, py);
    return(strlen(pn));
}

```

stripname

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

```

/*
 * cleanup() -- cleanup any tmp file
 * getseq() -- read in seq, set dna, len, maxlen
 * g_malloc() -- calloc() with error checkin
 * readjumps() -- get the good jumps, from tmp file if necessary
 * writejumps() -- write a filled array of jumps to a tmp file: nw()
 */
#include "nw.h"
#include <sys/file.h>

char *jname = "/tmp/homgXXXXXX";          /* tmp file for jumps */
FILE *fj;

int cleanup();                             /* cleanup tmp file */
long lseek();

/*
 * remove any tmp file if we blow
 */
cleanup(i)                                cleanup
{
    int i;
    if (fj)
        (void) unlink(jname);
    exit(i);
}

/*
 * read, return ptr to seq, set dna, len, maxlen
 * skip lines starting with ';', '<', or '>'
 * seq in upper or lower case
 */
char *
getseq(file, len)                          getseq
{
    char *file; /* file name */
    int *len; /* seq len */

    {
        char line[1024], *pseq;
        register char *px, *py;
        int natgc, tlen;
        FILE *fp;

        if ((fp = fopen(file, "r")) == 0) {
            fprintf(stderr, "%s: can't read %s\n", prog, file);
            exit(1);
        }
        tlen = natgc = 0;
        while (fgets(line, 1024, fp)) {
            if (*line == ';' || *line == '<' || *line == '>')
                continue;
            for (px = line; *px != '\n'; px++)
                if (isupper(*px) || islower(*px))
                    tlen++;
        }
        if ((pseq = malloc((unsigned)(tlen+6))) == 0) {
            fprintf(stderr, "%s: malloc() failed to get %d bytes for %s\n", prog, tlen+6, file);
            exit(1);
        }
        pseq[0] = pseq[1] = pseq[2] = pseq[3] = '\0';
    }
}

```

...getseq

```

py = pseq + 4;
*len = tlen;
rewind(fp);

while (fgets(line, 1024, fp)) {
    if (*line == ';' || *line == '<' || *line == '>')
        continue;
    for (px = line; *px != '\n'; px++) {
        if (isupper(*px))
            *py++ = *px;
        else if (islower(*px))
            *py++ = toupper(*px);
        if (index("ATGCU", *(py-1)))
            natgc++;
    }
    *py++ = '\0';
    *py = '\0';
    (void) fclose(fp);
    dna = natgc > (tlen/3);
    return(pseq+4);
}

char *
g_calloc(msg, nx, sz)                                g_calloc
{
    char *msg; /* program, calling routine */
    int nx, sz; /* number and size of elements */
    {
        char *px, *calloc();

        if ((px = calloc((unsigned)nx, (unsigned)sz)) == 0) {
            if (*msg) {
                fprintf(stderr, "%s: g_calloc() failed %s (n=%d, sz=%d)\n", prog, msg, nx, sz);
                exit(1);
            }
        }
        return(px);
    }
}

/*
 * get final jmps from dx[] or tmp file, set pp[], reset dmax: main()
 */
readjmps()                                            readjmps
{
    int fd = -1;
    int siz, i0, i1;
    register i, j, xx;

    if (fj) {
        (void) fclose(fj);
        if ((fd = open(jname, O_RDONLY, 0)) < 0) {
            fprintf(stderr, "%s: can't open() %s\n", prog, jname);
            cleanup(1);
        }
    }
    for (i = i0 = i1 = 0, dmax0 = dmax, xx = len0; i++) {
        while (1) {
            for (j = dx[dmax].ijmp; j >= 0 && dx[dmax].jp.x[j] >= xx; j--)
                ;

```

Page 2 of nwsubr.c

...readjumps

```

        if (j < 0 && dx[dmax].offset && fj) {
            (void) lseek(fd, dx[dmax].offset, 0);
            (void) read(fd, (char *)&dx[dmax].jp, sizeof(struct jmp));
            (void) read(fd, (char *)&dx[dmax].offset, sizeof(dx[dmax].offset));
            dx[dmax].ijmp = MAXJMP-1;
        }
        else
            break;
    }
    if (i >= JMPS) {
        fprintf(stderr, "%s: too many gaps in alignment\n", prog);
        cleanup(1);
    }
    if (j >= 0) {
        siz = dx[dmax].jp.n[j];
        xx = dx[dmax].jp.x[j];
        dmax += siz;
        if (siz < 0) { /* gap in second seq */
            pp[1].n[i1] = -siz;
            xx += siz;

            /* id = xx - yy + len1 - 1
             */
            pp[1].x[i1] = xx - dmax + len1 - 1;
            gapy++;
            ngapy -= siz;
            /* ignore MAXGAP when doing endgaps */
            siz = (-siz < MAXGAP || endgaps)? -siz : MAXGAP;
            i1++;
        }
        else if (siz > 0) { /* gap in first seq */
            pp[0].n[i0] = siz;
            pp[0].x[i0] = xx;
            gapx++;
            ngapx += siz;
            /* ignore MAXGAP when doing endgaps */
            siz = (siz < MAXGAP || endgaps)? siz : MAXGAP;
            i0++;
        }
    }
    else
        break;
}

/* reverse the order of jumps
 */
for (j = 0, i0--; j < i0; j++, i0--) {
    i = pp[0].n[j]; pp[0].n[j] = pp[0].n[i0]; pp[0].n[i0] = i;
    i = pp[0].x[j]; pp[0].x[j] = pp[0].x[i0]; pp[0].x[i0] = i;
}
for (j = 0, i1--; j < i1; j++, i1--) {
    i = pp[1].n[j]; pp[1].n[j] = pp[1].n[i1]; pp[1].n[i1] = i;
    i = pp[1].x[j]; pp[1].x[j] = pp[1].x[i1]; pp[1].x[i1] = i;
}
if (fd >= 0)
    (void) close(fd);
if (fj) {
    (void) unlink(jname);
    fj = 0;
    offset = 0;
}
}

```

```

/*
 * write a filled jmp struct offset of the prev one (if any): nw()
 */
writejumps(ix)
    int ix;
{
    char      *mktemp();

    if (!fj) {
        if (mktemp(jname) < 0) {
            fprintf(stderr, "%s: can't mktemp() %s\n", prog, jname);
            cleanup(1);
        }
        if ((fj = fopen(jname, "w")) == 0) {
            fprintf(stderr, "%s: can't write %s\n", prog, jname);
            exit(1);
        }
    }
    (void) fwrite((char *)&dx[ix].jp, sizeof(struct jmp), 1, fj);
    (void) fwrite((char *)&dx[ix].offset, sizeof(dx[ix].offset), 1, fj);
}

```

writejumps

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Hypothetical exemplifications are shown in Table 2, Comparisons 1 to 4, for determining % amino acid sequence identity (Table 2, Comparisons 1 and 2) and % nucleic acid sequence identity (Table 2, Comparisons 3 and 4) using the ALIGN-2 sequence comparison computer program, wherein "PRO" represents the amino acid sequence of a hypothetical PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide of interest, "Comparison Protein" represents the amino acid sequence of a polypeptide against which the "PRO" polypeptide of interest is being compared, "PRO-DNA" represents a hypothetical PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-encoding nucleic acid sequence of interest, "Comparison DNA" represents the nucleotide sequence of a nucleic acid molecule against which the "PRO-DNA" nucleic acid molecule of interest is being compared, "X", "Y" and "Z" each represent different hypothetical amino acid residues and "N", "L" and "V" each represent different hypothetical nucleotides.

For purposes herein, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows: 100 times the fraction X/Y, where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. As examples of % amino acid sequence identity calculations, Table 2 below presents comparisons 1 and 2 demonstrate how to calculate the % amino acid sequence identity of the amino acid sequence designated "Comparison Protein" to the amino acid sequence designated "PRO".

TABLE 2

Comparison 1	PRO	XXXXXXXXXXXXXXXXX	(Length = 15 amino acids)
	Comparison Protein	XXXXXXXXXXXXXX	(Length = 12 amino acids)
	% amino acid sequence identity = (the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) = 5 divided by 15 = 33.3%		
Comparison 2	PRO	XXXXXXXXXX	(Length = 10 amino acids)
	Comparison Protein	XXXXXXXXXXXXZZYZ	(Length = 15 amino acids)
	% amino acid sequence identity = (the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) = 5 divided by 10 = 50%		

Comparison 3	PRO-DNA	NNNNNNNNNNNNNN	(Length = 14 nucleotides)
	Comparison DNA	NNNNNNLLLLLLLLLL	(Length = 16 nucleotides)
	% nucleic acid sequence identity = (the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) = 6 divided by 14 = 42.9%		
Comparison 4	PRO-DNA	NNNNNNNNNNNNNN	(Length = 12 nucleotides)
	Comparison DNA	NNNNLLLVV	(Length = 9 nucleotides)
	% nucleic acid sequence identity = (the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) = 4 divided by 12 = 33.3%		

Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as described above using the ALIGN-2 sequence comparison computer program. However, % amino acid sequence identity can also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., *Nucleic Acids Res.* 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program can be downloaded from <http://www.ncbi.nlm.nih.gov>. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask = yes, strand = all, expected occurrences = 10, minimum low complexity length = 15/5, multi-pass e-value = 0.01, constant for multi-pass = 25, dropoff for final gapped alignment = 25 and scoring matrix = BLOSUM62.

In situations where NCBI-BLAST2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows: 100 times the fraction X/Y, where X is the number of amino acid residues scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A.

"PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 variant polynucleotide" or "PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 variant nucleic acid sequence" means a nucleic acid molecule which encodes an active PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide as defined herein and which has at least about 80% nucleic acid sequence identity with either (a) a nucleic acid sequence which encodes residues about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively, or (b) a nucleic acid sequence which encodes another specifically derived fragment of the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:4, respectively. Ordinarily, a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45,

PRO-C-MG.64 or PRO-C-MG.72 variant polynucleotide will have at least about 80% nucleic acid sequence identity, more preferably at least about 81% nucleic acid sequence identity, more preferably at least about 82% nucleic acid sequence identity, more preferably at least about 83% nucleic acid sequence identity, more preferably at least about 84% nucleic acid sequence identity, more preferably at least about 85% nucleic acid sequence identity, more preferably at least about 86% nucleic acid sequence identity, more preferably at least about 87% nucleic acid sequence identity, more preferably at least about 88% nucleic acid sequence identity, more preferably at least about 89% nucleic acid sequence identity, more preferably at least about 90% nucleic acid sequence identity, more preferably at least about 91% nucleic acid sequence identity, more preferably at least about 92% nucleic acid sequence identity, more preferably at least about 93% nucleic acid sequence identity, more preferably at least about 94% nucleic acid sequence identity, more preferably at least about 95% nucleic acid sequence identity, more preferably at least about 96% nucleic acid sequence identity, more preferably at least about 97% nucleic acid sequence identity, more preferably at least about 98% nucleic acid sequence identity and yet more preferably at least about 99% nucleic acid sequence identity with either (a) a nucleic acid sequence which encodes residues about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively, or (b) a nucleic acid sequence which encodes another specifically derived fragment of the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:4, respectively. PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polynucleotide variants do not encompass the native PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 nucleotide sequence.

Ordinarily, PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 variant polynucleotides are at least about 30 nucleotides in length, often at least about 60 nucleotides in length, more often at least about 90 nucleotides in length, more often at least about 120 nucleotides in length, more often at least about 150 nucleotides in length, more often at least about 180 nucleotides in length, more often at least about 210 nucleotides in length, more often at least about 240 nucleotides in length, more often at least about 270 nucleotides in length, more often at least about 300 nucleotides in length, more often at least about 450 nucleotides in length, more often at least about 600 nucleotides in length, more often at least about 900 nucleotides in length, or more.

"Percent (%) nucleic acid sequence identity" with respect to the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide-encoding nucleic acid sequences identified herein is defined as the percentage of nucleotides in a candidate sequence that are identical with the nucleotides in a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide-encoding nucleic acid sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent nucleic acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared. For purposes herein,

however, % nucleic acid sequence identity values are obtained as described below by using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Table 1 has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or can be compiled from the source code provided in Table 1. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

For purposes herein, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows: 100 times the fraction W/Z, where W is the number of nucleotides scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C. As examples of % nucleic acid sequence identity calculations, Table 2, Comparisons 3 and 4, demonstrate how to calculate the % nucleic acid sequence identity of the nucleic acid sequence designated "Comparison DNA" to the nucleic acid sequence designated "PRO-DNA".

Unless specifically stated otherwise, all % nucleic acid sequence identity values used herein are obtained as described above using the ALIGN-2 sequence comparison computer program. However, % nucleic acid sequence identity can also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., Nucleic Acids Res. 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program can be downloaded from <http://www.ncbi.nlm.nih.gov>. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask = yes, strand = all, expected occurrences = 10, minimum low complexity length = 15/5, multi-pass e-value = 0.01, constant for multi-pass = 25, dropoff for final gapped alignment = 25 and scoring matrix = BLOSUM62.

In situations where NCBI-BLAST2 is employed for sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows: 100 times the fraction W/Z, where W is the number of nucleotides scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C.

In other embodiments, PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72

variant polynucleotides are nucleic acid molecules that encode an active PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide and which are capable of hybridizing, preferably under stringent hybridization and wash conditions, to nucleotide sequences encoding the full-length PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide shown in SEQ ID NO:2 or SEQ ID NO:4, respectively. PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 variant polypeptides can be those that are encoded by a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 variant polynucleotide.

The term "positives", in the context of the amino acid sequence identity comparisons performed as described above, includes amino acid residues in the sequences compared that are not only identical, but also those that have similar properties. Amino acid residues that score a positive value to an amino acid residue of interest are those that are either identical to the amino acid residue of interest or are a preferred substitution (as defined in Table 3) of the amino acid residue of interest.

For purposes herein, the % value of positives of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % positives to, with, or against a given amino acid sequence B) is calculated as follows: 100 times the fraction X/Y, where X is the number of amino acid residues scoring a positive value as defined above by the sequence alignment program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % positives of A to B will not equal the % positives of B to A.

"Isolated," when used to describe the various polypeptides disclosed herein, means polypeptide that has been identified and separated and/or recovered from a component of its natural environment. Preferably, the isolated polypeptide is free of association with all components with which it is naturally associated. Contaminant components of its natural environment are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and can include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes polypeptide *in situ* within recombinant cells, since at least one component of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

An "isolated" nucleic acid molecule encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-encoding nucleic acid. Preferably, the isolated nucleic is free of association with all components with which it is naturally associated. An isolated PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-encoding nucleic acid

molecule is other than in the form or setting in which it is found in nature. Isolated nucleic acid molecules therefore are distinguished from the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-encoding nucleic acid molecule as it exists in natural cells. However, an isolated nucleic acid molecule encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide includes PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-encoding nucleic acid molecules contained in cells that ordinarily express PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

The term "antibody" is used in the broadest sense and specifically covers, for example, single anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 monoclonal antibodies (including agonist, antagonist, and neutralizing antibodies), anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody compositions with polyepitopic specificity, single chain anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibodies, and fragments of anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibodies (see below). The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that can be present in minor amounts.

"Stringency" of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature which can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of

hybridization reactions, see Ausubel *et al.*, *Current Protocols in Molecular Biology*, Wiley Interscience Publishers, (1995).

"Stringent conditions" or "high stringency conditions", as defined herein, can be identified by those that:

- (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42°C; or (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC (sodium chloride/sodium citrate) and 50% formamide at 55°C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA at 55°C.

"Moderately stringent conditions" can be identified as described by Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, New York: Cold Spring Harbor Press, 1989, and include the use of washing solution and hybridization conditions (e.g., temperature, ionic strength and %SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37°C in a solution comprising: 20% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 mg/ml denatured sheared salmon sperm DNA, followed by washing the filters in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

The term "epitope tagged" when used herein refers to a chimeric polypeptide comprising a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody can be made, yet is short enough such that it does not interfere with activity of the polypeptide to which it is fused. The tag polypeptide preferably also is fairly unique so that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 and 50 amino acid residues (preferably, between about 10 and 20 amino acid residues).

As used herein, the term "immunoadhesin" designates antibody-like molecules which combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site of an antibody (i.e., is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin can be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD or IgM.

"Active" or "activity" for the purposes herein refers to form(s) of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 which retain a biological and/or an immunological activity of native

or naturally-occurring PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, wherein "biological" activity refers to a biological function (either inhibitory or stimulatory), which includes enzymatic activity, caused by a native or naturally-occurring PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 other than the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 and an "immunological" activity refers to the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72.

The term "antagonist" is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes a biological activity of a native PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide disclosed herein. In a similar manner, the term "agonist" is used in the broadest sense and includes any molecule that mimics a biological activity of a native PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide disclosed herein. Suitable agonist or antagonist molecules specifically include agonist or antagonist antibodies or antibody fragments, fragments or amino acid sequence variants of native PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides, peptides, antisense molecules, and small organic molecules. Methods for identifying agonists or antagonists of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide include contacting a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, mRNA or gene with a candidate agonist or antagonist molecule and measuring a detectable change in one or more biological activities normally associated with the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

"Treatment" refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) the targeted pathologic condition or disorder. More specifically, "treatment" is an intervention performed with the intention of preventing the development or altering the pathology of a cardiovascular, endothelial, neovascular or angiogenic disorder or condition. The concept of treatment is used in the broadest sense, and specifically includes the prevention (prophylaxis), moderation, reduction, and curing of the disorder or condition, at any stage. Accordingly, "treatment" refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) said disorder or condition. The disorder may result from any cause, including idiopathic, cardiopathic, or myotrophic causes, or ischemia or ischemic insults, such as myocardial infarction. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented.

"Chronic" administration refers to administration of the agent(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial therapeutic effect (activity) for an extended period of time.

"Intermittent" administration is treatment that is not consecutively done without interruption, but rather is cyclic in nature.

"Microarray" refers to an array of distinct polynucleotides or oligonucleotides arranged on a substrate such as paper, nylon or other type of membrane, filter, gel, polymer, chip, glass slide, or any other suitable support,

including solid supports. The polynucleotides or oligonucleotides (the backbone chemistry can be any available in the art) can be synthesized on a substrate or prepared before application to the substrate.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, cats, cattle, horses, sheep, pigs, goats, rabbits, etc. Preferably, the mammal is human.

Administration "in combination with" one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

"Carriers" as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers which are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution.

"Antibody fragments" comprise a portion of an intact antibody, preferably the antigen binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies (Zapata et al., Protein Eng. 8(10): 1057-1062 [1995]); single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, a designation reflecting the ability to crystallize readily. Pepsin treatment yields an F(ab')₂ fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

"Fv" is the minimum antibody fragment which contains a complete antigen-recognition and -binding site. This region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the VH-VL dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab fragments differ from Fab' fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa and lambda, based on the amino acid sequences of their constant domains.

Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these can be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and

IgA2.

"Single-chain Fv" or "sFv" antibody fragments comprise the VH and VL domains of antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the VH and VL domains which enables the sFv to form the desired structure for antigen binding. For a review of sFv, see Pluckthun in The Pharmacology of Monoclonal Antibodies, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (VH) connected to a light-chain variable domain (VL) in the same polypeptide chain (VH - VL). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger et al., Proc. Natl. Acad. Sci. USA, 90:6444-6448 (1993).

An "isolated" antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and can include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequencer, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody in situ within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

The word "label" when used herein refers to a detectable compound or composition which is conjugated directly or indirectly to the antibody so as to generate a "labeled" antibody. The label can be detectable by itself (e.g. radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, can catalyze chemical alteration of a substrate compound or composition which is detectable.

By "solid phase" is meant a non-aqueous matrix to which the antibody of the present invention can adhere. Examples of solid phases encompassed herein include those formed partially or entirely of glass (e.g., controlled pore glass), polysaccharides (e.g., agarose), polyacrylamides, polystyrene, polyvinyl alcohol and silicones. In certain embodiments, depending on the context, the solid phase can comprise the well of an assay plate; in others it is a purification column (e.g., an affinity chromatography column). This term also includes a discontinuous solid phase of discrete particles, such as those described in U.S. Patent No. 4,275,149.

A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug (such as a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or antibody thereto) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

A "small molecule" is defined herein to have a molecular weight below about 500 Daltons.

The phrases "vascular or angiogenic disorder", "vascular or angiogenic dysfunction" are used

interchangeably and refer in part to systemic disorders that affect vessels, such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins, and/or lymphatics. This would include indications that stimulate angiogenesis, cardiovascularization, and/or neovascularization, and those that inhibit angiogenesis, cardiovascularization, and/or neovascularization.

“Hypertrophy”, as used herein, is defined as an increase in mass of an organ or structure independent of natural growth that does not involve tumor formation. Hypertrophy of an organ or tissue is due either to an increase in the mass of the individual cells (true hypertrophy), or to an increase in the number of cells making up the tissue (hyperplasia), or both. Certain organs, such as the heart, lose the ability to divide shortly after birth. Accordingly, “cardiac hypertrophy” is defined as an increase in mass of the heart, which, in adults, is characterized by an increase in myocyte cell size and contractile protein content without concomitant cell division. The character of the stress responsible for inciting the hypertrophy, (*e.g.*, increased preload, increased afterload, loss of myocytes, as in myocardial infarction, or primary depression of contractility), appears to play a critical role in determining the nature of the response. The early stage of cardiac hypertrophy is usually characterized morphologically by increases in the size of myofibrils and mitochondria, as well as by enlargement of mitochondria and nuclei. At this stage, while muscle cells are larger than normal, cellular organization is largely preserved. At a more advanced stage of cardiac hypertrophy, there are preferential increases in the size or number of specific organelles, such as mitochondria, and new contractile elements are added in localized areas of the cells, in an irregular manner. Cells subjected to long-standing hypertrophy show more obvious disruptions in cellular organization, including markedly enlarged nuclei with highly lobulated membranes, which displace adjacent myofibrils and cause breakdown of normal Z-band registration. The phrase “cardiac hypertrophy” is used to include all stages of the progression of this condition, characterized by various degrees of structural damage of the heart muscle, regardless of the underlying cardiac disorder. Hence, the term also includes physiological conditions instrumental in the development of cardiac hypertrophy, such as elevated blood pressure, aortic stenosis, or myocardial infarction.

“Heart failure” refers to an abnormality of cardiac function where the heart does not pump blood at the rate needed for the requirements of metabolizing tissues. The heart failure can be caused by a number of factors, including ischemic, congenital, rheumatic, or idiopathic forms.

“Congestive heart failure” (CHF) is a progressive pathologic state where the heart is increasingly unable to supply adequate cardiac output (the volume of blood pumped by the heart over time) to deliver the oxygenated blood to peripheral tissues. As CHF progresses, structural and hemodynamic damages occur. While these damages have a variety of manifestations, one characteristic symptom is ventricular hypertrophy. CHF is a common end result of a number of various cardiac disorders.

“Myocardial infarction” generally results from atherosclerosis of the coronary arteries, often with superimposed coronary thrombosis. It may be divided into two major types: transmural infarcts, in which myocardial necrosis involves the full thickness of the ventricular wall, and subendocardial (nontransmural) infarcts, in which the necrosis involves the subendocardium, the intramural myocardium, or both, without extending all the way through the ventricular wall to the epicardium. Myocardial infarction is known to cause both a change in hemodynamic effects and an alteration in structure in the damaged and healthy zones of the

Supravalvular "aortic stenosis" is an inherited vascular disorder characterized by narrowing of the ascending aorta, but other arteries, including the pulmonary arteries, may also be affected. Untreated aortic stenosis may lead to increased intracardiac pressure resulting in myocardial hypertrophy and eventually heart failure and death. The pathogenesis of this disorder is not fully understood, but hypertrophy and possibly hyperplasia of medial smooth muscle are prominent features of this disorder. It has been reported that molecular variants of the elastin gene are involved in the development and pathogenesis of aortic stenosis. U.S. Patent No. 5,650,282 issued July 22, 1997.

“Valvular regurgitation” occurs as a result of heart diseases resulting in disorders of the cardiac valves. Various diseases, like rheumatic fever, can cause the shrinking or pulling apart of the valve orifice, while other diseases may result in endocarditis, an inflammation of the endocardium or lining membrane of the atrioventricular orifices and operation of the heart. Defects such as the narrowing of the valve stenosis or the defective closing of the valve result in an accumulation of blood in the heart cavity or regurgitation of blood past the valve. If uncorrected, prolonged valvular stenosis or insufficiency may result in cardiac hypertrophy and associated damage to the heart muscle, which may eventually necessitate valve replacement.

The treatment of all these, and other cardiovascular (endothelial-involved) and angiogenic disorders are encompassed by the present invention.

20 The terms "cancer", "cancerous", and "malignant" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth.

The term “neovascularization” refers to growth and development of blood vessels in tissue not normally containing them, or of blood vessels of a different kind than usual in tissue.

The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (e.g., ¹³¹I, ¹²⁵I, ⁹⁰Y, and ¹⁸⁶Re), chemotherapeutic agents, and toxins such as enzymatically active toxins of bacterial, fungal, plant, or animal origin, or fragments thereof.

A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include alkylating agents, folic acid antagonists, anti-metabolites of nucleic acid metabolism, antibiotics, pyrimidine analogs, 5-fluorouracil, cisplatin, purine nucleosides, amines, amino acids, triazol nucleosides, or corticosteroids. Specific examples include Adriamycin, Doxorubicin, 5-Fluorouracil, Cytosine arabinoside ("Ara-C"), Cyclophosphamide, Thiotepa, Busulfan, Cytosine, Taxol, Toxotere, Methotrexate, Cisplatin, Melphalan, Vinblastine, Bleomycin, Etoposide, Ifosfamide, Mitomycin C, Mitoxantrone, Vincristine, Vinorelbine, Carboplatin, Teniposide, Daunomycin, Carminomycin, Aminopterin, Dactinomycin, Mitomycins, Esperamicins (see U.S. Pat. No. 4,675,187), Melphalan, and other related nitrogen mustards. Also included in this definition are hormonal agents that act to regulate or inhibit hormone action on tumors, such as tamoxifen and onapristone.

A "growth-inhibitory agent" when used herein refers to a compound or composition that inhibits growth of

a cell, such as an Wnt-overexpressing cancer cell, either *in vitro* or *in vivo*, and includes and is used interchangeably herein with angiostatic agents. Thus, the growth-inhibitory agent is one which significantly reduces the percentage of malignant cells in S phase, for example. Examples of growth-inhibitory agents include agents that block cell cycle progression (at a place other than S phase), such as agents that induce G1 arrest and M-phase arrest. Classical M-phase blockers include the vincas (vincristine and vinblastine), taxol, and topo II inhibitors such as doxorubicin, daunorubicin, etoposide, and bleomycin. Those agents that arrest G1 also spill over into S-phase arrest, for example, DNA alkylating agents such as tamoxifen, prednisone, dacarbazine, mechlorethamine, cisplatin, methotrexate, 5-fluorouracil, and ara-C. Further information can be found in The Molecular Basis of Cancer, Mendelsohn and Israel, eds., Chapter 1, entitled "Cell cycle regulation, oncogenes, and antineoplastic drugs" by Murakami *et al.* (WB Saunders: Philadelphia, 1995), especially p. 13. Additional examples include tumor necrosis factor (TNF), an antibody capable of inhibiting or neutralizing the angiogenic activity of acidic or basic FGF or hepatocyte growth factor (HGF), an antibody capable of inhibiting or neutralizing the coagulant activities of tissue factor, protein C, or protein S (see, WO 91/01753, published 21 February 1991), or an antibody capable of binding to HER2 receptor (WO 89/06692), such as the 4D5 antibody (and functional equivalents thereof) (*e.g.*, WO 92/22653).

A "cardiovascular agent" refers generically to any drug that acts in treating cardiovascular disorders. Examples of cardiovascular agents are those that promote vascular homeostasis by modulating blood pressure, heart rate, heart contractility, and endothelial and smooth muscle biology, all of which factors have a role in cardiovascular disease. Specific examples of these include angiotensin-II receptor antagonists; endothelin receptor antagonists such as, for example, BOSENTAN™ and MOXONODIN™; interferon-gamma (IFN-γ); des-aspartate-angiotensin I; thrombolytic agents, *e.g.*, streptokinase, urokinase, t-PA, and a t-PA variant specifically designed to have longer half-life and very high fibrin specificity, TNK t-PA (a T103N, N117Q, KHR(296-299)AAAA t-PA variant, Keyt *et al.*, Proc. Natl. Acad. Sci. USA 91, 3670-3674 (1994)); inotropic or hypertensive agents such as digoxigenin and β-adrenergic receptor blocking agents, *e.g.*, propranolol, timolol, tertalolol, carteolol, nadolol, betaxolol, penbutolol, acetobutolol, atenolol, metoprolol, and carvedilol; angiotensin converting enzyme (ACE) inhibitors, *e.g.*, quinapril, captopril, enalapril, ramipril, benazepril, fosinopril, and lisinopril; diuretics, *e.g.*, chlorothiazide, hydrochlorothiazide, hydroflumethazide, methylchlorothiazide, benzthiazide, dichlorphenamide, acetazolamide, and indapamide; and calcium channel blockers, *e.g.*, diltiazem, nifedipine, verapamil, nicardipine.

"Angiogenic agents" and "endothelial agents" are active agents that promote angiogenesis and/or endothelial cell growth, or, if applicable, vasculogenesis. This would include factors that accelerate wound healing, such as growth hormone, insulin-like growth factor-I (IGF-I), VEGF, VIGF, PDGF, epidermal growth factor (EGF), CTGF and members of its family, FGF, and TGF-α and TGF-β.

"Angiostatic agents" are active agents that inhibit angiogenesis or vasculogenesis or otherwise inhibit or prevent growth of cancer cells. Examples include antibodies or other antagonists to angiogenic agents as defined above, such as antibodies to VEGF. They additionally include cytotherapeutic agents such as cytotoxic agents, chemotherapeutic agents, growth-inhibitory agents, apoptotic agents, and other agents to treat cancer, such as anti-HER-2, anti-CD20, and other bioactive and organic chemical agent.

"Endothelial cell" means the cells of endothelial tissue, which includes the membranes lining serous cavities, heart, blood and lymph vessels.

In a pharmacological sense, in the context of the present invention, a "therapeutically effective amount" of an active agent such as a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or agonist or antagonist thereto or an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody, refers to an amount effective in the treatment of a cardiovascular, endothelial or angiogenic disorder in a mammal and can be determined empirically. An effective amount will either prevent, lessen the worsening of, alleviate, or cure the treated condition., or stimulate, enhance, reduce or inhibit the cellular response, biological activity, or stated purpose.

As used herein, an "effective amount" of an active agent such as a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or agonist or antagonist thereto or an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody, refers to an amount effective for carrying out a stated purpose, wherein such amounts may be determined empirically for the desired effect. An effective amount can stimulate, enhance, reduce or inhibit the cellular response, biological activity, or other stated purpose.

II. Compositions and Methods of the Invention

A. Full-length PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72

Polypeptide

The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. In particular, cDNA encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide has been identified and isolated, as disclosed in further detail in the Examples below. For sake of simplicity, in the present specification the protein encoded by DNA-C-MG.2-1776, DNA-C-MG.12-1776, DNA-C-MG.45-1776, DNA-C-MG.64-1776 or DNA-C-MG.72-1776 as well as all further native homologues and variants included in the foregoing definition of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, will be referred to as "PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72", regardless of their origin or mode of preparation.

As disclosed in the Examples below, a cDNA clone designated herein as DNA-C-MG.2-1776, DNA-C-MG.12-1776, DNA-C-MG.45-1776, DNA-C-MG.64-1776 or DNA-C-MG.72-1776 has been deposited with the ATCC. The actual nucleotide sequence of the clone can readily be determined by the skilled artisan by sequencing of the deposited clone using routine methods in the art. The predicted amino acid sequence can be determined from the nucleotide sequence using routine skill. For the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide and encoding nucleic acid described herein, Applicants have identified what is believed to be the reading frame best identifiable with the sequence information available at the time.

SEQ ID NO:1 shows a cDNA containing a nucleotide sequence (nucleotides 1-2891) encoding native sequence PRO-C-MG.2, wherein the nucleotide sequence (SEQ ID NO:1) is a clone designated herein as "DNA-C-MG.2-1776." SEQ ID NO:2 shows the amino acid sequence (SEQ ID NO:2) of a native sequence

PRO-C-MG.2 polypeptide as derived from the coding sequence of SEQ ID NO:1.

SEQ ID NO:3 shows a cDNA containing a nucleotide sequence (nucleotides 1-2119) encoding native sequence PRO-C-MG.12, wherein the nucleotide sequence (SEQ ID NO:3) is a clone designated herein as "DNA-C-MG.12-1776." SEQ ID NO:4 shows the amino acid sequence (SEQ ID NO:4) of a native sequence

PRO-C-MG.12 polypeptide as derived from the coding sequence of SEQ ID NO:3.

B. PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 Variants

In addition to the full-length native sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides described herein, it is contemplated that PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 variants can be prepared. PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 variants can be prepared by introducing appropriate nucleotide changes into the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 DNA, and/or by synthesis of the desired PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide. Those skilled in the art will appreciate that amino acid changes can alter post-translational processes of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

Variations in the native full-length sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 or in various domains of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 described herein, can be made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations set forth, for instance, in U.S. Patent No. 5,364,934. Variations can be a substitution, deletion or insertion of one or more codons encoding the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 that results in a change in the amino acid sequence of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 as compared with the native sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. Optionally the variation is by substitution of at least one amino acid with any other amino acid in one or more of the domains of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. Guidance in determining which amino acid residue can be inserted, substituted or deleted without adversely affecting the desired activity can be found by comparing the sequence of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 with that of homologous known protein molecules and minimizing the number of amino acid sequence changes made in regions of high homology. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, i.e., conservative amino acid replacements. Insertions or deletions can optionally be in the range of about 1 to 5 amino acids. The variation allowed can be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the full-length or mature native sequence.

PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide fragments are provided herein. Such fragments can be truncated at the N-terminus or C-terminus, or can lack internal residues, for example, when compared with a full length native protein. Certain fragments lack amino acid

residues that are not essential for a desired biological activity of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 fragments can be prepared by any of a number of conventional techniques. Desired peptide fragments can be chemically synthesized. An alternative approach involves generating PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 fragments by enzymatic digestion, e.g., by treating the protein with an enzyme known to cleave proteins at sites defined by particular amino acid residues, or by digesting the DNA with suitable restriction enzymes and isolating the desired fragment. Yet another suitable technique involves isolating and amplifying a DNA fragment encoding a desired polypeptide fragment, by polymerase chain reaction (PCR). Oligonucleotides that define the desired termini of the DNA fragment are employed at the 5' and 3' primers in the PCR. Preferably, PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide fragments share at least one biological and/or immunological activity with the native PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide shown in SEQ ID NO:2 or SEQ ID NO:4, respectively.

In particular embodiments, conservative substitutions of interest are shown in Table 3 under the heading of preferred substitutions. If such substitutions result in a change in biological activity, then more substantial changes, denominated exemplary substitutions in Table 3, or as further described below in reference to amino acid classes, are introduced and the products screened.

Table 3

	<u>Original Residue</u>	<u>Exemplary Substitutions</u>	<u>Preferred Substitutions</u>
25	Ala (A)	val; leu; ile	val
	Arg (R)	lys; gln; asn	lys
	Asn (N)	gln; his; lys; arg	gln
	Asp (D)	glu	glu
	Cys (C)	ser	ser
30	Gln (Q)	asn	asn
	Glu (E)	asp	asp
	Gly (G)	pro; ala	ala
	His (H)	asn; gln; lys; arg	arg
	Ile (I)	leu; val; met; ala; phe; norleucine	leu
35	Leu (L)	norleucine; ile; val; met; ala; phe	ile
	Lys (K)	arg; gln; asn	arg
	Met (M)	leu; phe; ile	leu
40	Phe (F)	leu; val; ile; ala; tyr leu	
	Pro (P)	ala	ala
	Ser (S)	thr	thr
	Thr (T)	ser	ser
	Trp (W)	tyr; phe	tyr
45	Tyr (Y)	trp; phe; thr; ser	phe
	Val (V)	ile; leu; met; phe; ala; norleucine	leu

Substantial modifications in function or immunological identity of the PRO-C-MG.2, PRO-C-MG.12,

PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties: (1) hydrophobic: norleucine, met, ala, val, leu, ile; (2) neutral hydrophilic: cys, ser, thr; (3) acidic: asp, glu; (4) basic: asn, gln, his, lys, arg; (5) residues that influence chain orientation: gly, pro; and (6) aromatic: trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Such substituted residues also can be introduced into the conservative substitution sites or, more preferably, into the remaining (non-conserved) sites.

The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis [Carter et al., Nucl. Acids Res., 13:4331 (1986); Zoller et al., Nucl. Acids Res., 10:6487 (1987)], cassette mutagenesis [Wells et al., Gene, 34:315 (1985)], restriction selection mutagenesis [Wells et al., Philos. Trans. R. Soc. London SerA, 317:415 (1986)] or other known techniques can be performed on the cloned DNA to produce the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 variant DNA.

Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant [Cunningham and Wells, Science, 244: 1081-1085 (1989)]. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, The Proteins, (W.H. Freeman & Co., N.Y.); Chothia, J. Mol. Biol., 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isoteric amino acid can be used.

C. Modifications of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72

Covalent modifications of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C- terminal residues of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. Derivatization with bifunctional agents is useful, for instance, for crosslinking PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 to a water-insoluble support matrix or surface for use in the method for purifying anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibodies, and vice-versa. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[(p-azidophenyl)dithio]propioimide.

Other modifications include deamidation of glutamyl and asparagyl residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains [T.E. Creighton, Proteins: Structure and Molecular Properties, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

Another type of covalent modification of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means), and/or adding one or more glycosylation sites that are not present in the native sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

Addition of glycosylation sites to the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide can be accomplished by altering the amino acid sequence. The alteration can be made, for example, by the addition of, or substitution by, one or more serine or threonine residues to the native sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 (for O-linked glycosylation sites). The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 amino acid sequence can optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

Another means of increasing the number of carbohydrate moieties on the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide is by chemical or enzymatic coupling of glycosides to the polypeptide. Such methods are described in the art, e.g., in WO 87/05330 published 11 September 1987, and in Aplin and Wriston, CRC Crit. Rev. Biochem., pp. 259-306 (1981).

Removal of carbohydrate moieties present on the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide can be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, et al., Arch. Biochem. Biophys., 259:52 (1987) and by Edge et al., Anal. Biochem., 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., Meth. Enzymol., 138:350 (1987).

Another type of covalent modification of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 comprises linking the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689;

4,301,144; 4,670,417; 4,791,192 or 4,179,337.

The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 of the present invention can also be modified in a way to form a chimeric molecule comprising PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 fused to another, heterologous polypeptide or amino acid sequence.

In one embodiment, such a chimeric molecule comprises a fusion of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl- terminus of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. The presence of such epitope-tagged forms of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field et al., Mol. Cell. Biol., 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan et al., Molecular and Cellular Biology, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky et al., Protein Engineering, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp et al., BioTechnology, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin et al., Science, 255:192-194 (1992)]; an α -tubulin epitope peptide [Skinner et al., J. Biol. Chem., 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth et al., Proc. Natl. Acad. Sci. USA, 87:6393-6397 (1990)].

In an alternative embodiment, the chimeric molecule can comprise a fusion of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule (also referred to as an "immunoadhesin"), such a fusion could be to the Fc region of an IgG molecule. The Ig fusions preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide in place of at least one variable region within an Ig molecule. In a particularly preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions see also US Patent No. 5,428,130 issued June 27, 1995.

In another embodiment, the chimeric molecule includes a fusion of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 with a signal peptide to allow or enhance secretion of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 peptide or even to change its localization within the host cell. The signal sequence is generally placed at the amino- or carboxyl- terminus of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, more usually the N-terminus when secretion or membrane localization is desired. Such fusions are typically intermediate products, since the signal peptide is usually specifically cleaved by enzymes of the host cell. Provision of a signal peptide

enables the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 to be readily purified following its secretion to the culture medium. Various signal polypeptides, which allow secretion or targeting to compartments within the cell, are well known in the art and are available for use with numerous host cells, including yeast and mammalian cells.

5 D. Preparation of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72

The description below relates primarily to production of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 by culturing cells transformed or transfected with a vector containing PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 nucleic acid. It is, of course, contemplated that alternative methods, which are well known in the art, can be employed to prepare PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. For instance, the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 sequence, or portions thereof, can be produced by direct peptide synthesis using solid-phase techniques [see, e.g., Stewart et al., Solid-Phase Peptide Synthesis, W.H. Freeman Co., San Francisco, CA (1969); Merrifield, J. Am. Chem. Soc., **85**:2149-2154 (1963)]. *In vitro* protein synthesis can be performed using manual techniques or by automation. Automated synthesis can be accomplished, for instance, using an Applied Biosystems Peptide Synthesizer (Foster City, CA) using manufacturer's instructions. Various portions of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be chemically synthesized separately and combined using chemical or enzymatic methods to produce the full-length PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72.

20 1. Isolation of DNA Encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72

DNA encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be obtained from a cDNA library prepared from tissue believed to possess the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 mRNA and to express it at a detectable level. Accordingly, human PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 DNA can be conveniently obtained from a cDNA library prepared from human tissue, such as described in the Examples. The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-encoding gene can also be obtained from a genomic library or by known synthetic procedures (e.g., automated nucleic acid synthesis).

Libraries can be screened with probes (such as antibodies to the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 or oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe can be conducted using standard procedures, such as described in Sambrook et al., Molecular Cloning: A Laboratory Manual (New York: Cold Spring Harbor Laboratory Press, 1989). An alternative means to isolate the gene encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 is to use PCR methodology [Sambrook et al., supra; Dieffenbach et al., PCR Primer: A Laboratory Manual (Cold Spring Harbor Laboratory Press, 1995)].

The Examples below describe techniques for screening a cDNA library. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are

minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in the art, and include the use of radiolabels like ^{32}P -labeled ATP, biotinylation or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook et al., supra.

Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence databases. Sequence identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined using methods known in the art and as described herein.

Nucleic acid having protein coding sequence can be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in Sambrook et al., supra, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

2. Selection and Transformation of Host Cells

Host cells are transfected or transformed with expression or cloning vectors described herein for PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in Mammalian Cell Biotechnology: a Practical Approach, M. Butler, ed. (IRL Press, 1991) and Sambrook et al., supra.

Methods of eukaryotic cell transfection and prokaryotic cell transformation are known to the ordinarily skilled artisan, for example, CaCl_2 , CaPO_4 , liposome-mediated and electroporation. Depending on the host cell used, transformation is performed using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook et al., supra, or electroporation is generally used for prokaryotes. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., Gene, 23:315 (1983) and WO 89/05859 published 29 June 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, Virology, 52:456-457 (1978) can be employed. General aspects of mammalian cell host system transfections have been described in U.S. Patent No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen et al., J. Bact., 130:946 (1977) and Hsiao et al., Proc. Natl. Acad. Sci. (USA), 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene, polyornithine, can also be used. For various techniques for transforming mammalian cells, see Keown et al., Methods in Enzymology, 185:527-537 (1990) and Mansour et al., Nature, 336:348-352 (1988).

Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain

W3110 (ATCC 27,325) and K5 772 (ATCC 53,635). Other suitable prokaryotic host cells include Enterobacteriaceae such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published 12 April 1989),

5 *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. These examples are illustrative rather than limiting. Strain W3110 is one particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell secretes minimal amounts of proteolytic enzymes. For example, strain W3110 can be modified to effect a genetic mutation in the genes encoding proteins endogenous to the host, with examples of such hosts including *E. coli* W3110 strain 1A2, which has the

10 complete genotype *tonA*; *E. coli* W3110 strain 9E4, which has the complete genotype *tonA ptr3*; *E. coli* W3110 strain 27C7 (ATCC 55,244), which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT kan'*; *E. coli* W3110 strain 37D6, which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT rbs7 ilvG kan'*; *E. coli* W3110 strain 40B4, which is strain 37D6 with a non-kanamycin resistant *degP* deletion mutation; and an *E. coli* strain having mutant periplasmic protease disclosed in U.S. Patent No.

15 4,946,783 issued 7 August 1990. Alternatively, *in vitro* methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-encoding vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others

20 include *Schizosaccharomyces pombe* (Beach and Nurse, Nature, 290: 140 [1981]; EP 139,383 published 2 May 1985); *Kluyveromyces* hosts (U.S. Patent No. 4,943,529; Fleer et al., Bio/Technology, 9:968-975 (1991)) such as, e.g., *K. lactis* (MW98-8C, CBS683, CBS4574; Louvencourt et al., J. Bacteriol., 154(2):737-742 [1983]), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilum* (ATCC 36,906; Van den Berg et al., Bio/Technology, 8:135 (1990)), *K.*

25 *thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna et al., J. Basic Microbiol., 28:265-278 [1988]); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa* (Case et al., Proc. Natl. Acad. Sci. USA, 76:5259-5263 [1979]); *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published 31 October 1990); and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium* (WO 91/00357 published 10 January 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance et al., Biochem. Biophys. Res. Commun., 112:284-289 [1983]; Tilburn et al., Gene, 26:205-221 [1983]; Yelton et al., Proc. Natl. Acad. Sci. USA, 81: 1470-1474 [1984]) and *A. niger* (Kelly and Hynes, EMBO J., 4:475-479 [1985]).

30 Methylophilic yeasts are suitable herein and include, but are not limited to, yeast capable of growth on methanol selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*, *Torulopsis*, and *Rhodotorula*. A list of specific species that are exemplary of this class of yeasts can be found in

35 C. Anthony, The Biochemistry of Methylophilic Yeasts, 269 (1982).

Suitable host cells for the expression of glycosylated PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9, as well as plant cells. Examples of useful

mammalian host cell lines include Chinese hamster ovary (CHO) and COS cells. More specific examples include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., J. Gen Virol., 36:59 (1977)); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, Proc. Natl. Acad. Sci. USA, 77:4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod., 23:243-251 (1980)); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). The selection of the appropriate host cell is deemed to be within the skill in the art.

3. Selection and Use of a Replicable Vector

The nucleic acid (e.g., cDNA or genomic DNA) encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector can, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence can be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally include, but are not limited to, one or more of a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques which are known to the skilled artisan.

The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which can be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence can be a component of the vector, or it can be a part of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-encoding DNA that is inserted into the vector. The signal sequence can be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence can be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders, the latter described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published 4 April 1990), or the signal described in WO 90/13646 published 15 November 1990. In mammalian cell expression, mammalian signal sequences can be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders.

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 μ plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells.

Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical

nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-encoding nucleic acid, such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub et al., Proc. Natl. Acad. Sci. USA, 77:4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 [Stinchcomb et al., Nature, 282:39 (1979); Kingsman et al., Gene, 7:141 (1979); Tschemper et al., Gene, 10:157 (1980)]. The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 [Jones, Genetics, 85:12 (1977)].

Expression and cloning vectors usually contain a promoter operably linked to the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-encoding nucleic acid sequence to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the β -lactamase and lactose promoter systems [Chang et al., Nature, 275:615 (1978); Goeddel et al., Nature, 281:544 (1979)], alkaline phosphatase, a tryptophan (*trp*) promoter system [Goeddel, Nucleic Acids Res., 8:4057 (1980); EP 36,776], and hybrid promoters such as the *tac* promoter [deBoer et al., Proc. Natl. Acad. Sci. USA, 80:21-25 (1983)]. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72.

Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman et al., J. Biol. Chem., 255:2073 (1980)] or other glycolytic enzymes [Hess et al., J. Adv. Enzyme Reg., 7:149 (1968); Holland, Biochemistry, 17:4900 (1978)], such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

Transcription of a DNA encoding the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 by higher eukaryotes can be increased by inserting an enhancer sequence into the vector.

Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the

cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer can be spliced into the vector at a position 5' or 3' to the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 coding sequence, but is preferably located at a site 5' from the promoter.

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72.

15 Still other methods, vectors, and host cells suitable for adaptation to the synthesis of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 in recombinant vertebrate cell culture are described in Gething et al., Nature, 293:620-625 (1981); Mantei et al., Nature, 281:40-46 (1979); EP 117,060; and EP 117,058.

4. Detecting Gene Amplification/Expression

Gene amplification and/or expression can be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, Proc. Natl. Acad. Sci. USA, 77:5201-5205 (1980)], dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies can be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn can be labeled and the assay can be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression, alternatively, can be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids can be either monoclonal or polyclonal, and can be prepared in any mammal. Conveniently, the antibodies can be prepared against a native sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 DNA and encoding a specific antibody epitope.

5. Purification of Polypeptide

Forms of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the

membrane using a suitable detergent solution (e.g. Triton-X 100) or by enzymatic cleavage. Cells employed in expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents.

It can be desired to purify PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. Various methods of protein purification can be employed and such methods are known in the art and described for example in Deutscher, Methods in Enzymology, 182 (1990); Scopes, Protein Purification: Principles and Practice, Springer-Verlag, New York (1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 produced.

E. Uses for PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72

Nucleotide sequences (or their complement) encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 have various applications in the art of molecular biology, including uses as hybridization probes, in chromosome and gene mapping and in the generation of anti-sense RNA, DNA, and PNA (peptide nucleic acids). PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 nucleic acid will also be useful for the preparation of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides by the recombinant techniques described herein. Full-length or fragments of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide coding sequence find use as, for example, hybridization probes or for encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or fragment thereof that can optionally encode a polypeptide comprising a binding site for an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody.

The full-length native sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 gene (SEQ ID NO:1 or SEQ ID NO:3, respectively), or portions thereof, can be used as hybridization probes for a cDNA library to isolate the full-length PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 cDNA or to isolate still other cDNAs (for instance, those encoding naturally-occurring variants of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 or PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 from other species) which have a desired sequence identity to the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 sequence disclosed in SEQ ID NO:1 or SEQ ID NO:3, respectively. The hybridization probes can be derived from at least partially novel regions of the nucleotide sequence of SEQ ID NO:1 or SEQ ID NO:3, respectively, wherein those regions can be determined without undue experimentation or from genomic

sequences including promoters, enhancer elements and introns of native sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72.

Such nucleic acid fragments are usually at least about 20 nucleotides in length, preferably at least about 30 nucleotides in length, more preferably at least about 40 nucleotides in length, yet more preferably at least about 50 nucleotides in length, yet more preferably at least about 60 nucleotides in length, yet more preferably at least about 70 nucleotides in length, yet more preferably at least about 80 nucleotides in length, yet more preferably at least about 90 nucleotides in length, yet more preferably at least about 100 nucleotides in length, yet more preferably at least about 110 nucleotides in length, yet more preferably at least about 120 nucleotides in length, yet more preferably at least about 130 nucleotides in length, yet more preferably at least about 140 nucleotides in length, yet more preferably at least about 150 nucleotides in length, yet more preferably at least about 160 nucleotides in length, yet more preferably at least about 170 nucleotides in length, yet more preferably at least about 180 nucleotides in length, yet more preferably at least about 190 nucleotides in length, yet more preferably at least about 200 nucleotides in length, yet more preferably at least about 250 nucleotides in length, yet more preferably at least about 300 nucleotides in length, yet more preferably at least about 350 nucleotides in length, yet more preferably at least about 400 nucleotides in length, yet more preferably at least about 450 nucleotides in length, yet more preferably at least about 500 nucleotides in length, yet more preferably at least about 600 nucleotides in length, yet more preferably at least about 700 nucleotides in length, yet more preferably at least about 800 nucleotides in length, yet more preferably at least about 900 nucleotides in length and yet more preferably at least about 1000 nucleotides in length, wherein in this context the term "about" means the referenced nucleotide sequence length plus or minus 10% of that referenced length. In a preferred embodiment, the nucleotide sequence fragment is derived from any coding region of the nucleotide sequence shown in SEQ ID NO:1 or SEQ ID NO:3, respectively. In one embodiment the fragment size range is from 20 to 50 nucleotides, which is particularly useful for probe or antisense use. It is noted that novel fragments of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide-encoding nucleotide sequence can be determined in a routine manner by aligning the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide-encoding nucleotide sequence with other known nucleotide sequences using any of a number of well known sequence alignment programs and determining which PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide-encoding nucleotide sequence fragment(s) are novel. All of such PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide-encoding nucleotide sequences are contemplated herein and can be determined without undue experimentation. Also contemplated are the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide fragments encoded by these nucleotide molecule fragments, preferably those PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide fragments that comprise a binding site for an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody.

By way of example, a screening method will comprise isolating the coding region of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 gene using the known DNA sequence to synthesize a selected probe of about 40 bases. Hybridization probes can be labeled by a variety of labels,

including radionucleotides such as ^{32}P or ^{35}S , or enzymatic labels such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems. Labeled probes having a sequence complementary to that of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 gene of the present invention can be used to screen libraries of human cDNA, genomic DNA or mRNA to determine which members of such libraries the probe hybridizes to. Hybridization techniques are described in further detail in the Examples below.

Any EST sequences disclosed in the present application can similarly be employed as probes, using the methods disclosed herein.

Other useful fragments of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 nucleic acids include antigene (antisense or sense) oligonucleotides comprising a single-stranded nucleic acid sequence (either RNA or DNA) capable of binding to target PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 mRNA (sense) or PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 DNA (antisense) sequences. Antigene compounds comprise a fragment of the sequence of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 gene as discussed above and in more detail below. The fragment can include either 5' or 3' non-coding regions.

PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 oligonucleotides and probes can also be employed in PCR techniques to generate a pool of sequences for identification of closely related PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 coding sequences.

Nucleotide sequences encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can also be used to construct hybridization probes for mapping the gene which encodes that PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 and for the genetic analysis of individuals with genetic disorders. The nucleotide sequences provided herein can be mapped to a chromosome and specific regions of a chromosome using known techniques, such as *in situ* hybridization, linkage analysis against known chromosomal markers, and hybridization screening with libraries.

When the coding sequences for PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 encode a protein which binds to another protein, the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be used in assays to identify the other proteins or molecules involved in the binding interaction. By such methods, inhibitors of the binding interaction can be identified. Proteins involved in such binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction. Also, the receptor PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be used to isolate correlative ligand(s). Screening assays can be designed to find lead compounds that mimic the biological activity of a native PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 or a receptor for PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates. Small molecules contemplated include synthetic organic or inorganic compounds. The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays and cell based assays, which are well characterized in the art. Such high- and ultra-high throughput assays are can

also be used to test antisense molecules. One such assay includes the use of reporter molecules, such as beta-lactamase, in which a beta-lactamase expression cassette is integrated into the test cell genome in such a way that modulation of the biological response of interest, e.g. tube formation, is reflected as modulation of beta-lactamase activity, preferably measured by fluorescence (e.g., see WO 98/13353 and WO 98/52047).

5 Nucleic acids which encode PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 or its modified forms can also be used to generate either transgenic animals or "knock out" animals which, in turn, are useful in the development and screening of therapeutically useful reagents. A transgenic animal (e.g., a mouse or rat) is an animal having cells that contain a transgene, which transgene was introduced into the animal or an ancestor of the animal at a prenatal, e.g., an embryonic stage. A transgene is a DNA which
10 is integrated into the genome of a cell from which a transgenic animal develops. In one embodiment, cDNA encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be used to clone genomic DNA encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 in accordance with established techniques and the genomic sequences used to generate transgenic animals that contain cells which express DNA encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. Methods for generating transgenic animals, particularly animals such as mice or rats, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009. Typically, particular cells would be targeted for PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 transgene incorporation with tissue-specific enhancers. Transgenic animals that include a copy of a transgene encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 introduced into the germ line of the animal at an embryonic stage can be used to examine the effect of increased expression of DNA encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. Such animals can be used as tester animals for reagents thought to confer protection from, for example, pathological conditions associated with its overexpression. In accordance with this facet of the invention, an animal is treated with the reagent and a reduced incidence of the pathological condition,
25 compared to untreated animals bearing the transgene, would indicate a potential therapeutic intervention for the pathological condition.

Alternatively, non-human homologues of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be used to construct a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 "knock out" animal which has a defective or altered gene encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 as a result of homologous recombination between the endogenous gene encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 and altered genomic DNA encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 introduced into an embryonic stem cell of the animal. For example, cDNA encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be used to clone genomic DNA
30 encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 in accordance with established techniques. A portion of the genomic DNA encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be deleted or replaced with another gene, such as a gene encoding a selectable marker which can be used to monitor integration. Typically, several kilobases of

unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector [see e.g., Thomas and Capecchi, *Cell*, 51:503 (1987) for a description of homologous recombination vectors]. The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected [see e.g., Li et al., *Cell*, 69:915 (1992)]. The selected cells are then injected into a blastocyst of an animal (e.g., a mouse or rat) to form aggregation chimeras [see e.g., Bradley, in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed. (IRL, Oxford, 1987), pp. 113-152]. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a "knock out" animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knockout animals can be characterized for instance, for their ability to defend against certain pathological conditions and for their development of pathological conditions due to absence of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

Nucleic acid encoding the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides can also be used in gene therapy. In gene therapy applications, genes are introduced into cells in order to achieve *in vivo* synthesis of a therapeutically effective genetic product, for example for replacement of a defective gene. Alternatively, *in vivo* synthesis of an antisense form of the target gene can reduce unwanted target gene expression, such as in the case of tumors, viral infections, or conditions involving gene overexpression. "Gene therapy" includes both conventional gene therapy where a lasting effect is achieved by a acute treatment (e.g., a single treatment), and the administration of gene therapeutic agents, which involves the one time or repeated administration of a therapeutically effective DNA or mRNA. Antisense RNAs and DNAs can be used as therapeutic agents for blocking the expression of certain genes *in vivo*. It has already been shown that short antisense oligonucleotides can be imported into cells where they act as inhibitors, despite their low intracellular concentrations caused by their restricted uptake by the cell membrane. (Zamecnik *et al.*, Proc. Natl. Acad. Sci. USA 83:4143-4146 [1986]). The oligonucleotides can be modified to enhance their uptake, e.g. by substituting their negatively charged phosphodiester groups by uncharged groups such as in peptide nucleic acids (PNAs).

There are a variety of techniques available for introducing nucleic acids, including antigene oligonucleotides, into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells *in vitro*, or *in vivo* in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells *in vitro* include the use of liposomes, electroporation, microinjection, cell fusion, DEAE-dextran, the calcium phosphate precipitation method, etc. In one embodiment, *in vivo* gene transfer techniques include transfection with viral (typically retroviral, such as adenovirus, lentivirus, Herpes simplex I virus, or adeno-associated virus (AAV)) vectors, viral coat protein-liposome mediated transfection (Dzau et al., *Trends in Biotechnology* 11:205-210 [1993]), and lipid-based systems (for example, DOTMA, DOPE, and DC-Chol; see, e.g., Tonkinson *et al.*, *Cancer Investigation* 14(1): 54-65 (1996)). WO 99/22772 discloses particularly useful liposomes for use with antigene oligonucleotides. A viral vector such as a retroviral vector includes at least one transcriptional promoter/enhancer or locus-defining element(s), or other elements

that control gene expression by other means such as alternate splicing, nuclear RNA export, or post-translational modification of messenger. In addition, a viral vector such as a retroviral vector includes a nucleic acid molecule that, when transcribed in the presence of a gene encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, is operably linked thereto and acts as a translation initiation sequence. Such vector constructs also include a packaging signal, long terminal repeats (LTRs) or portions thereof, and positive and negative strand primer binding sites appropriate to the virus used (if these are not already present in the viral vector). In addition, such vector typically includes a signal sequence for secretion of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide from a host cell in which it is placed. Preferably the signal sequence for this purpose is a mammalian signal sequence. Should the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide contain an C-terminal or internal translocation peptide, it is preferable to delete it or inactivate it by mutation to avoid interference with the heterologous secretion signal peptide activity. Optionally, the vector construct may also include a signal that directs polyadenylation, as well as one or more restriction sites and a translation termination sequence. By way of example, such vectors will typically include a 5' LTR, a tRNA binding site, a packaging signal, an origin of second-strand DNA synthesis, and a 3' LTR or a portion thereof. Other vectors can be used that are non-viral, such as cationic lipids, polylysine, and dendrimers.

In some situations it is desirable to provide the nucleic acid source with an agent that targets the target cells, such as an antibody specific for a cell surface membrane protein or the target cell, a ligand for a receptor on the target cell, etc. Where liposomes are employed, proteins which bind to a cell surface membrane protein associated with endocytosis can be used for targeting and/or to facilitate uptake, e.g. capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu et al., *J. Biol. Chem.* 262, 4429-4432 (1987); and Wagner et al., *Proc. Natl. Acad. Sci. USA* 87, 3410-3414 (1990). For review of gene marking and gene therapy protocols see Anderson et al., *Science* 256, 808-813 (1992).

Chromosome Markers. The sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. Moreover, there is a current need for identifying particular sites on the chromosome. Few chromosome marking reagents based on actual sequence data (repeat polymorphisms) are presently available for marking chromosomal location. The mapping of DNAs to chromosomes according to the present invention is an important first step in correlating those sequences with genes associated with disease.

Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the cDNA. Computer analysis for the 3'- untranslated region is used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the primer will yield an amplified fragment.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular DNA to a particular chromosome. Using the present invention with the same oligonucleotide primers, sublocalization can be

achieved with panels of fragments from specific chromosomes or pools of large genomic clones in an analogous manner. Other mapping strategies that can similarly be used to map to its chromosome include *in situ* hybridization, prescreening with labeled flow-sorted chromosomes, and preselection by hybridization to construct chromosome-specific cDNA libraries.

Fluorescence *in situ* hybridization (FISH) of a cDNA clone to a metaphase chromosomal spread can be used to provide a precise chromosomal location in one step. This technique can be used with cDNA as short as 500 or 600 bases; however, clones larger than 2,000 bp have a higher likelihood of binding to a unique chromosomal location with sufficient signal intensity for simple detection. FISH requires use of the clones from which the gene encoding the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide was derived, and the longer the better. For example, 2,000 bp is good, 4,000 bp is better, and more than 4,000 is probably not necessary to get good results a reasonable percentage of the time. For a review of this technique, see, Verma *et al.*, Human Chromosomes: a Manual of Basic Techniques (Pergamon Press, New York, 1988).

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, *Mendelian Inheritance in Man* (available online through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region is then identified through linkage analysis (coinheritance of physically adjacent genes).

Next, it is necessary to determine the differences in the cDNA or genomic sequence between affected and unaffected individuals. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

With current resolution of physical mapping and genetic mapping techniques, a cDNA precisely localized to a chromosomal region associated with the disease could be one of between 50 and 500 potential causative genes. (This assumes 1 megabase mapping resolution and one gene per 20 kb).

The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides and nucleic acid molecules of the present invention can also be used for tissue typing, wherein the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides of the present invention can be differentially expressed in one tissue as compared to another. PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 nucleic acid molecules will find use for generating probes for PCR, Northern analysis, Southern analysis and Western analysis.

The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides described herein can also be employed as molecular weight markers for protein electrophoresis purposes.

F. PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 Antigene Compounds

PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 nucleic acids include antigene compounds, particularly oligonucleotides, for use in modulating the function of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, modulating the amount of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 produced by the cell, and ultimately modulating

the biological processes or responses in which PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 is critical. This can be accomplished by providing antigene compounds which specifically hybridize with one or more nucleic acids encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. As used herein, the terms "target nucleic acid" and "nucleic acid encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72" encompass DNA encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 (e.g., genomic DNA), RNA (including pre-mRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as "antisense" technology, however, is now more broadly referred to as "antigene" technology, which expressly includes both sense and antisense sequences and is used herein interchangeably with "antisense." Antigene compounds include peptide nucleic acids and ribozymes. The functions of DNA to be interfered with include replication and transcription. The functions of RNA to be interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. In the context of the present invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene. In the context of the present invention, inhibition is the preferred form of modulation of gene expression and mRNA is a preferred target.

In the present invention, the target is a nucleic acid molecule encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. Methods are available in the art to rapidly determine (within about a week) a site or sites within this gene for the antigene interaction to occur such that the desired effect, e.g., detection or modulation of expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, will result. A preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon." A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function *in vivo*. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). Eukaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used *in vivo* to initiate translation of an mRNA molecule transcribed from a gene encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, regardless of the sequence(s) of such codons. It is also known in the art that a translation termination codon (or "stop codon") of a gene may have

one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and

5 "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon.

The open reading frame (ORF) or "coding region," which refers to the region between the translation initiation codon and the translation termination codon, can also be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), which is the portion of an mRNA in the 5' direction from the

10 translation initiation codon and includes nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), which is the portion of an mRNA in the 3' direction from the translation termination codon and thus includes nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the

15 mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap. The 5' cap region is also a preferred target region.

While some eukaryotic mRNA transcripts are directly translated, many contain one or more regions known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated)

20 regions are known as "exons" and are spliced together to form a continuous mRNA sequence. When present, mRNA splice sites, i.e., intron-exon junctions, are also preferred target regions, and are particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also preferred targets. Introns are also effective target regions for antigene compounds targeted, for example, to

25 DNA or pre-mRNA.

Once one or more target sites have been identified, using techniques in the art, oligonucleotides are chosen which are sufficiently complementary to the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 gene target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect. The ability to derive an antisense or a sense oligonucleotide, based upon a cDNA sequence encoding a

30 given protein is described in, for example, Stein and Cohen (Cancer Res. 48:2659 (1988)) and van der Krol et al. (BioTechniques 6:958 (1988)). For example, targeting sites can be rapidly determined using combinatorial libraries, preferably in microarrays. Synthesis of peptide nucleic acid combinatorial libraries is disclosed in U.S. Patent 5,864,010. Antisense or sense oligonucleotides include PNAs or other molecules having modified backbones or modified nucleosides so long as they are designed upon and specific for a PRO-C-MG.2, PRO-C-

35 MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 nucleic acid sequence.

The sequence of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable, although 100% complementarity is preferred. An antigene compound is specifically hybridizable

when binding of the compound to the target PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed.

Methods for administration of antigene compounds to a variety of cells, including HUVEC, in order to modulate target gene function are known (e.g., Ackermann et al., J. Biol. Chem. 274(16):11245-52 (1999)).

The specificity and sensitivity of antigene is particularly suited for therapeutic uses. Antigene oligonucleotides have been employed as therapeutic moieties in the treatment of disease states in animals and man. Antigene oligonucleotides have been safely and effectively administered to humans and numerous clinical trials are presently underway. Antisense oligonucleotides have demonstrated acceptable safety and toxicity profiles in both animals and humans. Numerous antisense molecules are in Phase II and Phase III trials. An antisense compound has been approved and is marketed for treatment of CMV-induced retinitis. As a class, antisense molecules have been proven safe in animals and humans for systemic delivery. It has thus been established that antigene therapy can be a useful therapeutic modality that can be configured to be useful in treatment regimes for treatment of cells, tissues and animals, especially humans. Methods for testing toxicity and efficacy in animal models are thus well-established in the art.

In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally-occurring portions which function similarly. Such modified or substituted oligonucleotides are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases.

The antigene compounds in accordance with this invention preferably comprise from about 5 to about 60 nucleobases. Particularly preferred are antigene oligonucleotides comprising from about 8 to about 30 nucleobases (i.e. from about 8 to about 30 linked nucleosides), and most preferably from about 15 to about 25 nucleosides. Sequences of 17-18 bases are of special interest since this is the estimated length of unique sequences in the human genome. As is known in the art, a nucleoside is a nucleobase-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide

structure, the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

Accordingly, binding of antigene oligonucleotides, either antisense or sense oligonucleotides, to target nucleic acid sequences results in the formation of duplexes or triplexes that block transcription or translation of the target sequence by one of several means, including enhanced degradation of the duplexes, premature termination of transcription or translation, or by other means. The antisense oligonucleotides thus can be used to block expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 proteins. Antisense or sense oligonucleotides further comprise oligonucleotides having modified sugar-phosphodiester backbones (or other sugar linkages, such as those described in WO 91/06629) and wherein such sugar linkages are resistant to endogenous nucleases. Such oligonucleotides with resistant sugar linkages are stable *in vivo* (i.e., capable of resisting enzymatic degradation) but retain sequence specificity to be able to bind to target nucleotide sequences. Other examples of sense or antisense oligonucleotides include those oligonucleotides which are covalently linked to organic moieties, such as those described in WO 90/10048, and other moieties that increase affinity of the oligonucleotide for a target nucleic acid sequence, such as poly-(L-lysine). Further still, intercalating agents, such as ellipticine, and alkylating agents or metal complexes can be attached to sense or antisense oligonucleotides to modify binding specificities of the antisense or sense oligonucleotide for the target nucleotide sequence as discussed below.

Specific examples of preferred PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compounds include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included. Representative United States patents that teach the preparation of the phosphorus-containing linkages include, but are not limited to, U.S. Pat. Nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050, each of which is herein incorporated by reference.

Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and

alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH₂ component parts. Representative United States patents that teach the preparation of these oligonucleosides include, but are not limited to, U.S. Pat. Nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439, each of which is herein incorporated by reference.

In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent solubility, membrane-traversing, and hybridization properties, is referred to as a peptide nucleic acid (PNA; Nielsen *et al.*, *Science* 254:1497-1500 (1991)). In PNA compounds, the sugar-backbone is replaced with an amide containing backbone, e.g., an aminoethylglycine backbone. The nucleobases can be retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S. Pat. Nos. 5,539,082; 5,714,331; and 5,719,262, and PCT publication No. WO 97/33551, each of which is herein incorporated by reference. PNA compounds recognize and bind sequence-selectively and strand-selectively to double-stranded DNA (dsDNA), which is accomplished via strand displacement, in which the PNA binds via Watson-Crick binding to its complementary strand and extrudes the other strand in a virtually single-stranded conformation. PNA compounds also recognize and bind sequence-selectively to single-stranded DNA (ssDNA) and to RNA. This recognition by PNA of RNA, ssDNA or dsDNA can take place in sequences at least 5 bases long. A more preferred recognition sequence length is 5 to 60 base pairs long, and more preferably 8 to 30 base pairs long, and most preferably from about 15 to about 25 nucleosides. For therapeutic use of PNA compounds the targets of the PNA compounds would generally be double stranded DNA--in which case the PNA is effective in both the sense and antisense forms--and RNA. For diagnostic use, investigations methods and reagents where DNA is isolated outside of a cell, the DNA can be denatured to single stranded DNA and use of the PNA compound would be targeted to such single stranded DNA as well as RNA.

PNA compounds useful to effect binding to RNA, ssDNA and dsDNA and to form duplex and triplex complexes are polymeric strands formed from a polyamide, polythioamide, polysulfonamide or polysulfonamide backbone with a plurality of ligands located at spaced locations along the backbone, at least some of the ligands capable of hydrogen bonding with other ligands either on the compounds or nucleic acid targets. The amino acids which form the backbone may be identical or different, but those based on 2-aminoethyl-glycine are preferred. In some cases it may be of interest to attach ligands at either terminus to modulate the binding

characteristics of the PNAs. Representative ligands include DNA intercalators, which improve dsDNA binding or basic groups, such as lysine or polylysine, which strengthen the binding of the PNA due to electrostatic interaction. To decrease electrostatic repulsion charged groups such as carboxyl and sulfo groups could be used. Oligonucleotides and/or oligonucleoside can be covalently bound to either terminal positions to form chimeras containing PNA portions and oligonucleotide and/or oligonucleoside portions. Nucleosides and/or nucleotides (mono, di or tri-phosphates) also can be attached to the terminal positions. Moieties can also be located on non-terminal positions. In one embodiment, the PNA oligomers are conjugated to low molecular weight effector ligands such as ligands having nuclease activity or alkylating activity or reporter ligands (fluorescent, spin labels, radioactive, protein recognition ligands, for example, biotin or haptens). In another embodiment, the PNAs are conjugated to peptides or proteins, where the peptides have signaling activity and the proteins are, for example, enzymes, transcription factors or antibodies. Also, the PNAs can be attached to water-soluble or water-insoluble polymers. In yet another embodiment, the PNAs are conjugated to oligonucleotides or carbohydrates. When desired a PNA oligomer can be synthesized onto a moiety (e.g., a peptide chain, reporter, intercalator or other type of ligand-containing group) attached to a solid support.

In a further embodiment, PNA compounds also can be used as PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 gene-sequence specific gene activators and synthetic transcription factors, useful for selectively up-regulating PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. Transcription initiation by RNA polymerase involves the sequence specific recognition of the double-stranded DNA promoter either by the polymerase itself or by auxiliary transcription factors. Subsequently a transcription initiation open complex is formed in which about 12 base pairs of the DNA helix are melted, which exposes the bases of the template strand for base pairing with the RNA strand being synthesized. It has been demonstrated that an E. coli phage T7 RNA polymerase can utilize synthetic "RNA/DNA bubble duplex" complexes containing an RNA/DNA duplex and a single-stranded DNA D-loop for transcription elongation. In addition, homopyrimidine PNAs also form D-loop structures when binding to complementary double-stranded DNA by strand displacement, structures that behave like RNA/DNA open complex structures and are recognized by RNA polymerase.

Preferred embodiments of the invention are PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular -CH₂-NH-O-CH₂-, -CH₂-N(CH₃)-O-CH₂- [known as a methylene (methylimino) or MM1 backbone], -CH₂-O-N(CH₃)-CH₂-, -CH₂-N(CH₃)-N(CH₃)-CH₂- and -O-N(CH₃)-CH₂-CH₂- [wherein the native phosphodiester backbone is represented as -O-P-O-CH₂-] of the above referenced U.S. Pat. No. 5,489,677, and the amide backbones of U.S. Pat. 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures as described in U.S. Pat. 5,034,506.

Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C1 to C10 alkyl or C2 to C10 alkenyl and alkynyl. Particularly preferred are O[(CH₂)[n]O][m]CH₃, O(CH₂)[n]OCH₃, O(CH₂)[n]NH₂, O(CH₂)[n]CH₃, O(CH₂)[n]ONH₂, and

O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C1 to C10 lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O-CH₂CH₂OCH₃, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta*, 78:486-504 (1995); McKay et al., *J. Biol. Chem.* 274(3):1715-22 (1999)) i.e., an alkoxyalkoxy group. The incorporation of 2'-O-(2-methoxy)ethyl chemistry provides a number of significant improvements in oligonucleotide characteristics, including an increase in hybridization affinity toward a complementary RNA (1.5° C per modification) and an increase in resistance toward both 3'-exonuclease and intracellular nucleases. These improvements result in a substantial increase in oligonucleotide potency (e.g., >20-fold after 72 h). A further preferred modification includes 2'-dimethylaminoethoxy, i.e., a O(CH₂)₂ON(CH₃)₂ group, also known as 2'-DMAOE.

Other preferred modifications include 2'-methoxy (2'-O-CH₃), 2'-aminopropoxy (2'-OCH₂CH₂CH₂NH₂) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S. Pat. Nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920, each of which is herein incorporated by reference in its entirety.

Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further nucleobases include those disclosed in U.S. Pat. 3,687,808, in "The Concise Encyclopedia Of Polymer Science And Engineering," pages 858-859, Kroschwitz, ed. John Wiley & Sons, (1990), in Englisch et al., *Angewandte Chemie, International Edition*, 30:613 (1991), and by Sanghvi (Antisense Research and Applications, Chapter 15, pages 289-302, Crooke and Lebleu, ed., CRC Press, (1993)).

Nucleobases that are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. The 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2° C. (Sanghvi, *Id.* at pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications. Representative United States patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. Pat. No 3,687,808, as well as U.S. Pat. Nos. 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091; 5,614,617; 5,681,941; and 5,750,692, each of which is herein incorporated by reference.

Another modification of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., Proc. Natl. Acad. Sci. USA 86:6553-6556 (1989)), cholic acid (Manoharan et al., Bioorg. Med. Chem. Lett. 4:1053-1060 (1994)), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., Ann. N. Y. Acad. Sci. 660:306-309 (1992); Manoharan et al., Bioorg. Med. Chem. Lett. 3:2765-2770 (1993)), a thiocholesterol (Oberhauser et al., Nucl. Acids Res. 20:533-538 (1992)), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., EMBO J. 10:1111-1118 (1991); Kabanov et al., FEBS Lett. 259:327-330 (1990); Svinarchuk et al., Biochimie 75:49-54 (1993)), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., Tetrahedron Lett., 36:3651-3654 (1995); Shea et al., Nucl. Acids Res. 18:3777-3783 (1990)), a polyamine or a polyethylene glycol chain (Manoharan et al., Nucleosides & Nucleotides 14:969-973 (1995)), or adamantane acetic acid (Manoharan et al., Tetrahedron Lett. 36:3651-3654 (1995)), a palmityl moiety (Mishra et al., Biochim. Biophys. Acta 1264:229-237 (1995), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., J. Pharmacol. Exp. Ther., 277: 923-937 (1996)). Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S. Pat. Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941, each of which is herein incorporated by reference.

It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the modifications described can be incorporated into a single compound or even at a single nucleoside within an oligonucleotide. Accordingly, the present invention also includes PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compounds which are chimeric compounds. By

"chimeric PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene"

compounds or "antigene chimeras" is meant antigene compounds, particularly oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of an oligonucleotide compound. These oligonucleotides typically contain at least one region wherein the

oligonucleotide is modified so as to confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. An additional region of the oligonucleotide can serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids, such as an RNase. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of oligonucleotide inhibition of gene expression. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art. Chimeric antigene compounds of the invention can be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described herein. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3' "wing" segments of linked nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound.

Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides.

Oligonucleotides of the second type are also known in the art as "hemimers" or "wingmers." Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S. Pat. Nos. 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, each of which is herein incorporated by reference in its entirety. The term "prodrug" indicates a therapeutic agent that is prepared in an inactive form that is converted to an active form (i.e., drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or conditions. Included as PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compounds are their prodrug versions. For example, prodrug versions of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 oligonucleotides can be prepared as SATE [(S-acetyl-2-thioethyl) phosphate] derivatives according to the methods disclosed in WO 93/24510 or in WO 94/26764.

PNA compounds of the invention can be synthesized by any methodology, including those disclosed in WO 92/20702, WO/92/20703 and U.S. Patent 5,641,625.

The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene oligonucleotide compounds of the invention can be conveniently and routinely made through the well-known technique of solid phase synthesis. Any other means for such synthesis known in the art can additionally or alternatively be employed.

The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compounds of the invention can be admixed, encapsulated, conjugated or otherwise associated with other

molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption assisting formulations include, but are not limited to, U.S. Pat. Nos. 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291; 5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330; 4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170; 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of which is herein incorporated by reference.

The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the invention, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents. The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto. Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge et al., "Pharmaceutical Salts," J. of Pharma Sci. 66:1-19 (1977)). The base addition salts of said acidic compounds are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in the conventional manner. The free acid forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention. As used herein, a "pharmaceutical addition salt" includes a pharmaceutically acceptable salt of an acid form of one of the components of the compositions of the invention. These include organic or inorganic acid salts of the amines. Preferred acid salts are the hydrochlorides, acetates, salicylates, nitrates and phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid, glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic acid; and with amino acids, such as the 20 alpha-amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with

phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid.

- 5 Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible.

- For PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 oligonucleotides,
 10 preferred examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic
 15 acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine. The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene
 20 compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. For therapeutics, an animal, preferably a human, suspected of having a disease or disorder as discussed herein, which can be treated by modulating the expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, is treated by administering antigene compounds in accordance with the invention. The compounds of the invention can be utilized in pharmaceutical compositions by adding an
 25 effective amount of an antigene compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the antigene compounds and methods of the invention can also be useful prophylactically, e.g., to prevent or delay the desired response.

- The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene
 compounds, as research and diagnostic agents, hybridize to nucleic acids encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, enabling sandwich and other assays to easily be
 30 constructed. Hybridization of the antigene oligonucleotides of the invention with a nucleic acid encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be detected by means known in the art. Such means include conjugation of an enzyme to the oligonucleotide, radiolabelling of the
 oligonucleotide, fluorescence reporters, or any other suitable detection means. Kits using such detection means for detecting the level of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 in a
 35 sample can be prepared. PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compounds can be introduced into a cell containing the target nucleic acid sequence by any gene transfer method, including, for example, CaPO_4 -mediated DNA transfection, electroporation, or by using gene transfer vectors such as Epstein-Barr virus, and those discussed in detail herein. In brief, in a preferred

procedure, an antisense or sense oligonucleotide is inserted into a suitable retroviral vector. A cell containing the target nucleic acid sequence is contacted with the recombinant retroviral vector, either *in vivo* or *ex vivo*. Suitable retroviral vectors include, but are not limited to, those derived from the murine retrovirus M-MuLV, N2 (a retrovirus derived from M-MuLV), or the double copy vectors designated DCT5A, DCT5B and DCT5C (see WO 90/13641). Sense or antisense oligonucleotides also can be introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand binding molecule, as described in WO 91/04753. Suitable ligand binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially interfere with the ability of the ligand binding molecule to bind to its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell. Alternatively, a sense or an antisense oligonucleotide can be introduced into a cell containing the target nucleic acid sequence by formation of an oligonucleotide-lipid complex, as described in WO 90/10448. The sense or antisense oligonucleotide-lipid complex is preferably dissociated within the cell by an endogenous lipase. These and other methods are discussed in more detail herein.

Accordingly, the present invention also includes pharmaceutical compositions and formulations which include the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compounds of the invention. The pharmaceutical compositions of the present invention are administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal, oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration. PNAs, administered *i.p.*, have been shown to cross the blood-brain barrier and specifically reduce targeted gene expression (see e.g., Tyler et al., PNAS 96(12):7053-8 (1999)) *in vivo*.

Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful. Compositions and formulations for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, sachets or tablets. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable.

Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients. Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions may be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and

self-emulsifying semisolids.

The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

In one embodiment of the present invention the pharmaceutical compositions may be formulated and used as foams. Pharmaceutical foams include formulations such as, but not limited to, emulsions, microemulsions, creams, jellies and liposomes. While basically similar in nature these formulations vary in the components and the consistency of the final product. The preparation of such compositions and formulations is generally known to those skilled in the pharmaceutical and formulation arts and may be applied to the formulation of the compositions of the present invention.

Emulsions. The compositions of the present invention may be prepared and formulated as emulsions. Emulsions are typically heterogenous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1 μ m in diameter. (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199 (1988); Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245 (1988); Block in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335 (1988); Higuchi et al., in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pa., p. 301 (1985)). Emulsions are often biphasic systems comprising of two immiscible liquid phases intimately mixed and dispersed with each other. In general, emulsions may be either water in oil (w/o) or of the oil in water (o/w) variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase the resulting composition is called a water in oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase the resulting composition is called an oil in water (o/w) emulsion. Emulsions may contain additional components in addition to the dispersed phases and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed. Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil in water in oil (o/w/o) and water in oil in water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water

droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water stabilized in an oily continuous provides an o/w/o emulsion. Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into the external or continuous phase and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of stabilizing emulsions entail the use of emulsifiers that may be incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199 (1988)).

Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285 (1988); Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199 (1988)). Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting surfactants in the preparation of formulations. Surfactants may be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and amphoteric (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285 (1988)).

Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations. These include polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as bentonite, attapulgite, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants (Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335 (1988); Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199 (1988)).

Hydrophilic colloids or hydrocolloids include naturally occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth), cellulose derivatives (for example, carboxymethyl cellulose and carboxypropyl cellulose), and synthetic polymers (for example, carbomers, cellulose ethers, and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming strong interfacial films around the

dispersed-phase droplets and by increasing the viscosity of the external phase.

Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides that may readily support the growth of microbes, these formulations often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added to emulsion formulations to prevent deterioration of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

The application of emulsion formulations via dermatological, oral and parenteral routes and methods for their manufacture have been reviewed in the literature (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199 (1988)). Emulsion formulations for oral delivery have been very widely used because of reasons of ease of formulation, efficacy from an absorption and bioavailability standpoint. (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245 (1988); Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199 (1988)). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

Microemulsions. In one embodiment of the present invention, the compositions of oligonucleotides and nucleic acids are formulated as microemulsions. A microemulsion may be defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245 (1988)). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, in: *Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., VCH Publishers, New York, pages 185-215 (1989)). Microemulsions commonly are prepared via a combination of three to five components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and geometric packing of the polar heads and hydrocarbon tails of the surfactant molecules (Schott, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pa., p. 271 (1985)).

The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245 (1988); Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335 (1988)). Compared to conventional emulsions,

microemulsions offer the advantage of solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed spontaneously.

Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers, polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprate (MCA750), decaglycerol monooleate (MO750), decaglycerol sequioleate (SO750), decaglycerol decaoleate (DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art. The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain (C8-C12) mono, di, and triglycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides, saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone oil.

Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides (Constantinides et al., *Pharmaceutical Research*, 11:1385-1390 (1994); Ritschel, *Meth. Find. Exp. Clin. Pharmacol.* 13:205 (1993)). Microemulsions afford advantages of improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity (Constantinides et al., *Pharmaceutical Research*, 11:1385 (1994); Ho et al., *J. Pharm. Sci.*, 85:138-143 (1996)). Often microemulsions may form spontaneously when their components are brought together at ambient temperature. This may be particularly advantageous when formulating thermolabile drugs, peptides or oligonucleotides. Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of oligonucleotides and nucleic acids from the gastrointestinal tract, as well as improve the local cellular uptake of oligonucleotides and nucleic acids within the gastrointestinal tract, vagina, buccal cavity and other areas of administration.

Microemulsions of the present invention may also contain additional components and additives such as sorbitan monostearate and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the oligonucleotides and nucleic acids of the present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as belonging to one of five broad categories-surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, p. 92 (1991)), as discussed

Liposomes. There are many organized surfactant structures besides microemulsions that have been

studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the duration of action they offer from the standpoint of drug delivery. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

5 Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as efficiently with the cell wall, are taken up by macrophages in vivo.

10 In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each with a diameter less than 50 nm, under the influence of a suitable transdermal gradient. Therefore, it is desirable to use a liposome which is highly deformable and able to pass through such fine pores.

15 Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245 (1988)). Important considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

20 Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes. As the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

25 Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

30 Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the skin. The majority of applications resulted in the targeting of the upper epidermis.

35 Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang et al., *Biochem. Biophys. Res. Commun.*, 147:980-985 (1987)).

 Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes

have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou et al., *Journal of Controlled Release*, 19:269-274 (1992)).

One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

Several studies have assessed the topical delivery of liposomal drug formulations to the skin. Application of liposomes containing interferon to guinea pig skin resulted in a reduction of skin herpes sores while delivery of interferon via other means (e.g. as a solution or as an emulsion) were ineffective (Weiner et al., *Journal of Drug Targeting*, 2:405-410 (1992)). Further, an additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon using an aqueous system, and concluded that the liposomal formulation was superior to aqueous administration (du Plessis et al., *Antiviral Research* 18:259-265 (1992)).

Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin, in particular systems comprising non-ionic surfactant and cholesterol. Non-ionic liposomal formulations comprising Novasome TM I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome TM II (glyceryl distearate/cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the dermis of mouse skin. Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of cyclosporin-A into different layers of the skin.

Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside G[M1], or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen et al., *FEBS Letters*, 223:42 (1987); Wu et al., *Cancer Research* 53:3765 (1993)).

Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos et al. (*Ann. N.Y. Acad. Sci.* 507:64 (1987)) reported the ability of monosialoganglioside G[M1], galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon et al. (*PNAS* 85:6949 (1988)). U.S. Pat. No. 4,837,028 and WO 88/04924 disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside G[M1] or a galactocerebroside sulfate ester. U.S. Pat.No.

5,543,152 discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-sn-dimyristoylphosphatidylcholine are disclosed in WO 97/13499. Synthetic versions of these molecules are preferred.

Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto et al. (*Bull. Chem. Soc. Jpn.* 53:2778 (1980)) described liposomes comprising a nonionic detergent, 2C1215G, that contains a PEG moiety. Illum et al. (*FEBS Lett.* 167:79 (1984)) noted that hydrophilic coating of polystyrene particles with polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Pat. Nos. 4,426,330 and 4,534,899). Klibanov et al. (*FEBS Lett.* 268:235 (1990)) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume et al. (*Biochimica et Biophysica Acta* 1029:91 (1990)) extended such observations to other PEG-derivatized phospholipids, e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle et al. (U.S. Pat. Nos. 5,013,556 and 5,356,633) and Martin et al. (U.S. Pat. No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Pat. No. 5,225,212 (both to Martin et al.) and in WO 94/20073 (Zalipsky et al.) Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi et al.). U.S. Pat. Nos. 5,540,935 (Miyazaki et al.) and 5,556,948 (Tagawa et al.) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces. A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry et al. discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Pat. No. 5,264,221 to Tagawa et al. discloses protein-bonded liposomes and asserts that the contents of such liposomes may include an antisense RNA. U.S. Pat. 5,665,710 to Rahman et al. describes certain methods of encapsulating oligodeoxynucleotides in liposomes. WO 97/04787 to Love et al. discloses liposomes comprising antisense oligonucleotides targeted to the raf gene.

Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, e.g. they are self-optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

Surfactants find wide application in formulations such as emulsions (including microemulsions) and

liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, N.Y. p. 285 (1988)).

If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values. In general their HLB values range from 2 to about 18 depending on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included in this class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of this class.

If the surfactant molecule has the ability to carry either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

The use of surfactants in drug products, formulations and in emulsions has been reviewed (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, N.Y., p. 285 (1988)).

Penetration Enhancers. In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides, to the skin of animals. Most drugs are present in solution in both ionized and nonionized forms. However, usually only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs. Penetration enhancers may be classified as belonging to one of five broad categories, i.e., surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., *Crit. Rev. Ther. Drug Carrier Systems* p.92 (1991)). Each of the above mentioned classes of penetration enhancers are described below in greater detail.

Surfactants: In connection with the present invention, surfactants (or "surface-active agents") are chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid, with the result that absorption of

oligonucleotides through the mucosa is enhanced. In addition to bile salts and fatty acids, these penetration enhancers include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether) (Lee et al., Crit. Rev. Ther. Drug Carrier Systems, p.92 (1991)); and perfluorochemical emulsions, such as FC-43. Takahashi et al., J. Pharm. Pharmacol. 40:252 (1988)).

5 Fatty acids: Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprate, monoolein (1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arachidonic acid, glycerol 1-monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, C[1-10]-alkyl esters thereof (e.g., methyl, isopropyl and t-butyl), and mono- and di-glycerides thereof (i.e.,
10 oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., Crit. Rev. Ther. Drug Carrier Systems p.92 (1991); Muranishi, Crit. Rev. Ther. Drug Carrier Systems 7:1-33 (1990); El Hariri et al., J. Pharm. Pharmacol. 44: 651-654 (1992)).

Bile salts: The physiological role of bile includes the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's The Pharmacological Basis of Therapeutics, 9th Ed., Hardman et al. Eds., McGraw-Hill, New York pp. 934-935 (1996)). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. The bile salts of the invention include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), glucolic acid (sodium
20 glucolate), glycholic acid (sodium glycocholate), glycodeoxycholic acid (sodium glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate), ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydro-fusidate (STDHF), sodium glycodihydrofusidate and polyoxyethylene-9-lauryl ether (POE) (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, page 92 (1991); Swinyard, Chapter 39 In: Remington's Pharmaceutical
25 Sciences, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, Pa., pages 782-783 (1990); Muranishi, Critical Reviews in Therapeutic Drug Carrier Systems, 7: 1-33 (1990); Yamamoto et al., J. Pharm. Exp. Ther. 263:25 (1992); Yamashita et al., J. Pharm. Sci. 79:579-583 (1990)).

Chelating Agents: Chelating agents, as used in connection with the present invention, can be defined as compounds that remove metallic ions from solution by forming complexes therewith, with the result that
30 absorption of oligonucleotides through the mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal ion for catalysis and are thus inhibited by chelating agents (Jarrett, J. Chromatogr. 618:315-339 (1993)). Chelating agents of the invention include but are not limited to disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate,
35 5-methoxysalicylate and homovanilate), N-acyl derivatives of collagen, laureth-9 and N-amino acyl derivatives of beta-diketones (enamines)(Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, page 92 (1991); Muranishi, Critical Reviews in Therapeutic Drug Carrier Systems, 7:1-33 (1990); Buur et al., J. Control Rel. 14:43-51 (1990)).

Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant activity as chelating agents or as surfactants but that nonetheless enhance absorption of oligonucleotides through the alimentary mucosa (Muranishi, Critical Reviews in Therapeutic Drug Carrier Systems, 7: 1-33 (1990)). This class of penetration enhancers include, for example, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, page 92 (1991)); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., J. Pharm. Pharmacol. 39:621-626 (1987)).

Agents that enhance uptake of oligonucleotides at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi et al, U.S. Pat. No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo et al., PCT Application WO 97/30731), are also known to enhance the cellular uptake of oligonucleotides.

Other agents may be utilized to enhance the penetration of the administered nucleic acids, including glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

Carriers. Certain compositions of the present invention also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a nucleic acid, or analog thereof, which is inert (i.e., does not possess biological activity per se) but is recognized as a nucleic acid by in vivo processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For example, the recovery of a partially phosphorothioate oligonucleotide in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'-isothiocyano-stilbene-2,2'-disulfonic acid (Miyao et al., Antisense Res. Dev. 5:115-121 (1995); Takakura et al., Antisense & Nucl. Acid Drug Dev. 6:177-183 (1996)).

Excipients. In contrast to a carrier compound, a "pharmaceutical carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); lubricants (e.g., magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, etc.); disintegrants (e.g., starch, sodium starch glycolate, etc.); and

wetting agents (e.g., sodium lauryl sulphate, etc.).

Pharmaceutically acceptable organic or inorganic excipient suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous solutions, non-aqueous solutions in common solvents such as alcohols, or solutions of the nucleic acids in liquid or solid oil bases. The solutions may also contain buffers, diluents and other suitable additives.

Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral administration which do not deleteriously react with nucleic acids can be used.

Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Other Components. The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compositions of the present invention can additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact with the nucleic acid(s) of the formulation.

Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

Administration. The formulation of therapeutic compositions and their subsequent administration is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can generally be estimated based on EC50s found to be effective in in vitro and in vivo animal models. In general, dosage is from 0.01 ug to 100 g per kg of body weight, and can be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years. Preferred dosage is from 0.005 to 35 mg/kg body weight, even more preferred is 0.05 to 20 mg/kg body weight,

and yet more preferred is 0.01 to 10 mg/kg body weight. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 ug to 100 g per kg of body weight, once or more daily, to once every 20 years.

5 G. Screening Assays for Drug Candidates. This invention encompasses methods of screening compounds to identify those that mimic the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide (agonists) or prevent the effect of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide (antagonists). Screening assays for antagonist drug candidates are designed to identify compounds that bind or complex with the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides encoded by the genes identified herein, or with a gene and
10 mRNAs encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, or otherwise interfere with the interaction of the encoded polypeptides with other cellular proteins. These screening assays will include assays amenable to high- or ultra-high-throughput screening of chemical libraries, making them particularly suitable for identifying antigene (antisense or sense) and small molecule drug
15 candidates.

Polypeptide-targeted assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays, target nucleic acid binding assays, and cell-based assays, which are well characterized in the art. A drug candidate is contacted with a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide encoded by a nucleic acid identified herein
20 under conditions and for a time sufficient to allow these two components to interact.

In binding assays, the interaction is binding and the complex formed can be isolated or detected in the reaction mixture. In a particular embodiment, the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide encoded by the gene identified herein or the drug candidate is immobilized on a solid phase, e.g., on a microtiter plate, by covalent or non-covalent attachments. Non-covalent attachment generally is accomplished by coating the solid surface with a solution of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide and drying. Alternatively, an immobilized antibody, e.g., a monoclonal antibody, specific for the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide to be immobilized can be used to anchor it to a solid surface. The assay is performed by adding the non-immobilized component, which can be labeled by a
25 detectable label, to the immobilized component, e.g., the coated surface containing the anchored component. When the reaction is complete, the non-reacted components are removed, e.g., by washing, and complexes anchored on the solid surface are detected. When the originally non-immobilized component carries a detectable label, the detection of label immobilized on the surface indicates that complexing occurred. Where the originally non-immobilized component does not carry a label, complexing can be detected, for example, by
30 using a labeled antibody specifically binding the immobilized complex.
35

If the candidate compound interacts with but does not bind to a particular PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide encoded by a gene identified herein, its interaction with that polypeptide can be assayed by methods well known for detecting protein-protein

interactions. Such assays include traditional approaches, such as, e.g., cross-linking, co-immunoprecipitation, and co-purification through gradients or chromatographic columns. In addition, protein-protein interactions can be monitored by using a yeast-based genetic system described by Fields and co-workers (Fields and Song, Nature (London), 340:245-246 (1989); Chien et al., Proc. Natl. Acad. Sci. USA, 88:9578-9582 (1991)) as

5 disclosed by Chevray and Nathans, Proc. Natl. Acad. Sci. USA, 89: 5789-5793 (1991). Many transcriptional activators, such as yeast GAL4, consist of two physically discrete modular domains, one acting as the DNA-binding domain, the other one functioning as the transcription-activation domain. The yeast expression system described in the foregoing publications (generally referred to as the "two-hybrid system") takes advantage of this property, and employs two hybrid proteins, one in which the target protein is fused to the DNA-binding
10 domain of GAL4, and another, in which candidate activating proteins are fused to the activation domain. The expression of a GAL1-*lacZ* reporter gene under control of a GAL4-activated promoter depends on reconstitution of GAL4 activity via protein-protein interaction. Colonies containing interacting polypeptides are detected with a chromogenic substrate for β -galactosidase. A complete kit (MATCHMAKER™) for identifying protein-protein interactions between two specific proteins using the two-hybrid technique is
15 commercially available from Clontech. This system can also be extended to map protein domains involved in specific protein interactions as well as to pinpoint amino acid residues that are crucial for these interactions.

Compounds that interfere with the interaction of a gene encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide identified herein and other intra- or extracellular components can be tested as follows: usually a reaction mixture is prepared containing the product of the gene
20 and the intra- or extracellular component under conditions and for a time allowing for the interaction and binding of the two products. To test the ability of a candidate compound to inhibit binding, the reaction is run in the absence and in the presence of the test compound. In addition, a placebo can be added to a third reaction mixture, to serve as positive control. The binding (complex formation) between the test compound and the intra- or extracellular component present in the mixture is monitored as described herein. The formation of a
25 complex in the control reaction(s) but not in the reaction mixture containing the test compound indicates that the test compound interferes with the interaction of the test compound and its reaction partner. A particularly useful assay system is a microarray assay, such as chip upon which a nucleic acid fragment-sequence library--based on the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 gene sequence--is synthesized.

30 Oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein can be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously (to produce a transcript image), to identify genetic variants, mutations and polymorphisms, to identify effective nucleic acid binding molecules such as antisense molecules, regulatory proteins, ribosomes or polymerases. This information may be used to determine gene function, to understand
35 the genetic basis of disease, to diagnose disease, to identify therapeutic molecules (e.g., antisense), and to develop, and monitor the activities of therapeutic agents.

In one embodiment, the microarray can be prepared and used according to the methods known in the art, such as those described in WO95/11995 (Chee *et al.*), Lockhart, D. J., *et al.* (*Nat. Biotech.* 14: 1675-1680

(1996)), and Schena, M., *et al.* (*Proc. Natl. Acad. Sci.* 93: 10614-10619 (1996)) or in WO 99/24463.

The microarray is preferably composed of a large number of unique, single-stranded nucleic acid sequences, usually either synthetic antisense oligonucleotides or fragments of cDNAs, fixed to a solid support. The oligonucleotides are preferably about 5 to 60 nucleotides in length, more preferably about 8 to 30, even more preferably about 15 to 30 nucleotides in length, even more preferably 15 to 25, and most preferably about 20 to 25 nucleotides in length. For a certain type of microarray, it may be preferable to use oligonucleotides that are only 7 to 10 nucleotides in length. The microarray can contain oligonucleotides which cover the known 5' (or 3') sequence or untranslated regions, sequential oligonucleotides which cover the full-length sequence or unique oligonucleotides selected from particular areas along the length of the sequence including untranslated regions. Polynucleotides used in the microarray can be oligonucleotides that are specific to a gene or genes of interest, preferably a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 gene, in which at least a fragment of the sequence is known or that are specific to one or more unidentified cDNAs that are common to a particular cell or tissue type or to a normal, developmental, or disease state. In certain situations, it is appropriate to use pairs of oligonucleotides on a microarray. The pairs will be identical, except for one nucleotide preferably located in the center of the sequence. The second oligonucleotide in the pair (mismatched by one) serves as a control. The number of oligonucleotide pairs may range from 2 to 1,000,000. Microarrays can also contain fragments in DNA duplex form, which are particularly useful in identifying molecules that bind to PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 genomic DNA.

For producing oligonucleotides to a known sequence for a microarray, the gene of interest is examined using a computer algorithm which starts at the 5' or more preferably at the 3' end of the nucleotide sequence. The algorithm identifies oligomers of defined length that are unique to the gene, have a GC content within a range suitable for hybridization, and lack predicted secondary structure that may interfere with hybridization.

In one aspect, the oligonucleotides are synthesized at designated areas on the surface of a substrate, for example by using a light-directed chemical coupling procedure and an inkjet application apparatus, such as that described in WO95/251116 (Baldeschweiler *et al.*). The substrate may be paper, nylon or any other type of membrane, filter, chip, glass slide, or any other suitable solid support. In another aspect, a "gridded" array analogous to a dot or slot blot (HYBRIDOT apparatus, GIBCO/BRL) may be used to arrange and link cDNA fragments or oligonucleotides to the surface of a substrate using a vacuum system, thermal, UV, mechanical or chemical bonding procedures. In a most preferred embodiment, each of the different predefined regions is physically separated from each other of the different regions. In yet another aspect, an array may be produced by hand or by using available devices, materials, and machines (including BRINKMANN multichannel pipettors or robotic instruments). Such an array may contain 8, 24, 96, 384, 1536, or 6144 oligonucleotides, or any other multiple from 2 to 1,000,000 that lends itself to the efficient use of commercially available instrumentation. In one preferred embodiment the array includes at least 1,000 different oligonucleotides attached to surface of the solid support, and more preferably at least 10,000 different oligonucleotides. Oligonucleotides are preferably attached to the first surface of the solid support through a linker group. The oligonucleotide in the different predefined regions are at least 20% pure, more preferably are at least 50% pure,

even more preferably at least 80% pure, and most preferably at least 90% pure.

In one embodiment, the array contains a planar, non-porous solid support having at least a first surface, and a plurality of different oligonucleotides attached to the first surface of the solid support at a density exceeding 400 different oligonucleotides per square cm, wherein each of the different oligonucleotides is attached to the surface of the solid support in a different predefined region, has a different determinable sequence, and is at least 6 nucleotides in length, with preferred lengths as discussed above, wherein at least one of the different oligonucleotides is a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 sequence. In this embodiment each different oligonucleotides is from about 6 to about 20 nucleotides in length, more preferably at least 10 nucleotides in length, and most preferably at least 20 nucleotides in length.

In a most preferred embodiment, each of the different predefined regions is physically separated from each other of the different regions. Oligonucleotides are preferably attached to the first surface of the solid support through a linker group. The oligonucleotide in the different predefined regions are at least 20% pure, more preferably are at least 50% pure, even more preferably at least 80% pure, and most preferably at least 90% pure.

Sample analysis using the microarrays can be conducted by extracting polynucleotides from a biological sample. The biological samples are obtained from any bodily fluid (blood, urine, saliva, phlegm, gastric juices, *etc.*), cultured cells, biopsies, or other tissue preparations. The polynucleotides extracted from the sample can be used to produce, as probes, nucleic acid sequences that are complementary to the nucleic acids on the microarray. If the microarray consists of cDNAs, antisense RNAs (aRNA) are appropriate probes. Therefore, in one aspect, mRNA is used to produce cDNA that, in turn and in the presence of fluorescent nucleotides, is used to produce fragment or oligonucleotide aRNA probes. These fluorescently-labeled probes are incubated with the microarray so that the probe sequences hybridize to the cDNA oligonucleotides of the microarray. In another aspect, nucleic acid sequences used as probes can include polynucleotides, fragments, and complementary or antisense sequences produced using restriction enzymes, PCR technologies, and OLIGOLABELING™ or TRANSPROBE™ kits (Pharmacia) well known in the area of hybridization technology. In an alternative microarray embodiment, oligonucleotides (preferably antisense molecules) are employed on the support and the target cDNA is the soluble binding component of the assay.

Incubation conditions are adjusted so that hybridization occurs with precise complementary matches or with various degrees of less complementarity. After removal of nonhybridized probes, a scanner is used to determine the levels and patterns of fluorescence. The scanned images are examined to determine degree of complementarity and the relative abundance of each oligonucleotide sequence on the microarray. A detection system may be used to measure the absence, presence, and amount of hybridization for all of the distinct sequences simultaneously. This data may be used for large-scale correlation studies or functional analysis of the sequences, mutations, variants, or polymorphisms among samples (Heller, R. A., et al., Proc. Natl. Acad. Sci. 94: 2150-55 (1997)).

For gene mapping, a gene or a cloned DNA fragment is hybridized to an ordered array of DNA fragments, and the identity of the DNA elements applied to the array is unambiguously established by the pixel or pattern of pixels of the array that are detected. In constructing physical maps of the genome, arrays of immobilized cloned DNA fragments are hybridized with other cloned DNA fragments to establish whether the

cloned fragments in the probe mixture overlap and are therefore contiguous to the immobilized clones on the array. For example, Meier-Ewert et al., (J. Biotech. 35(2-3):191-203 (1994)) disclose such an application.

The arrays of immobilized DNA fragments may also be used for genetic diagnostics. For example, array containing multiple forms of a mutated gene or genes can be probed with a labeled mixture of a patient's DNA which will preferentially interact with only one of the immobilized versions of the gene. The detection of this interaction can provide a medical diagnosis. Arrays of immobilized DNA fragments can also be used in DNA probe diagnostics. For unambiguous genotyping or identifying a DNA- or RNA-containing sample as that of a human, the identity of the test sample can be established unambiguously by hybridizing the sample to an array containing DNA from different organisms, including human, wherein one or more PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 genes sequences are included in the array. Other molecules of genetic interest, such as cDNAs and RNAs can be immobilized on the array or alternately used as the labeled probe mixture that is applied to the array.

In one embodiment, a potential antagonist includes a polypeptide or small molecule that binds to the fusions of immunoglobulin with PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, and, in particular are antibodies including, without limitation, poly- and monoclonal antibodies and antibody fragments, single-chain antibodies, anti-idiotypic antibodies, and chimeric or humanized versions of such antibodies or fragments, as well as human antibodies and antibody fragments. Alternatively, a potential antagonist can be a closely related protein or peptide, for example, a mutated form of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide that recognizes a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 binding protein or substrate but imparts no effect, thereby competitively inhibiting the action of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

Another potential PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide antagonist is an antigene (antisense or sense) construct, as described herein, prepared using antisense technology, where, for example, the antisense molecule acts to reduce directly the translation of mRNA by hybridizing to targeted PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 mRNA or the sense or antisense molecule reduces transcription of the mRNA by hybridizing to PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 genomic DNA (typically through triple-helix formation), both means preventing or reducing protein translation of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. For example, the 5' coding portion of the polynucleotide sequence, which encodes the mature PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides herein, is used to design an antisense RNA or DNA or PNA oligonucleotide of from about 5 to 60 base pairs in length. The antisense oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of the mRNA molecule into the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide (antisense--Okano, Neurochem., 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression (CRC Press: Boca Raton, FL, 1988). A PNA sense or antisense oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix--see Lee et al., Nucl. Acids Res., 6:3073 (1979); Cooney et al., Science, 241: 456 (1988); Dervan et al., Science,

251:1360 (1991)), thereby preventing transcription and the production of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide. The oligonucleotides described above can also be delivered to cells such that the antigene molecule can be expressed *in vivo* to inhibit production of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide. When antisense DNA is used, oligodeoxyribonucleotides derived from the translation-initiation site, e.g., between about -10 and +10 positions of the target gene nucleotide sequence, are preferred.

Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA.

Ribozymes act by sequence-specific hybridization to the complementary target RNA, followed by endonucleolytic cleavage. Specific ribozyme cleavage sites within a potential RNA target can be identified by known techniques. For further details see, e.g., Rossi, Current Biology, 4:469-471 (1994), and PCT publication No. WO 97/33551 (published September 18, 1997).

As discussed herein, nucleic acid molecules in triple-helix formation used to inhibit transcription can be single-stranded and composed of deoxynucleotides. Such molecules can have backbone bonds not naturally found in DNA or RNA. A preferred form are PNAs. Such molecules that form a triplex with PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 gene can also act as agonists to up-modulate PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 transcription when appropriately targeted as discussed herein.

Potential antagonists include small molecules that bind to the active site, the protein binding site, or other relevant binding site (e.g., co-factor binding site, substrate binding site) of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, thereby blocking the normal biological activity of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide. Examples of small molecules include, but are not limited to, small peptides or peptide-like molecules, preferably soluble peptides, and synthetic non-peptidyl organic or inorganic compounds.

These small molecules can be identified by any one or more of the screening assays discussed herein and/or by any other screening techniques well known for those skilled in the art.

For example, the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 agonist can be screened for the ability to stimulate or reduce the proliferation of or tube formation of endothelial cells as described herein. In brief, in the proliferation assay, human umbilical vein endothelial cells are obtained and cultured in 96-well flat-bottomed culture plates (Costar, Cambridge, MA) and supplemented with a reaction mixture appropriate for facilitating proliferation of the cells. The compound to be screened is added and, after incubation at 37°C, cultures are pulsed with 3-H-thymidine and harvested onto glass fiber filters (phD; Cambridge Technology, Watertown, MA). Mean 3-H-thymidine incorporation (cpm) of triplicate cultures is determined using a liquid scintillation counter (Beckman Instruments, Irvine, CA). Significant 3-(H)thymidine incorporation indicates stimulation of endothelial cell proliferation. To assay for antagonists, the assays described herein can be performed. For example, in the above assay, a compound to be screened is added and its ability to inhibit 3-(H)thymidine incorporation is determined.

The compositions useful in the treatment of disorders and conditions provided herein include, without limitation, antibodies, small organic and inorganic molecules, peptides, phosphopeptides, antisense and

ribozyme molecules, triple-helix molecules, etc., that inhibit the expression and/or activity of the target gene product.

The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides and nucleic acid molecules of the present invention are particularly useful for detecting, monitoring, analyzing, or identifying, as described herein, the occurrence or progression of angiogenesis or vasculogenesis, as can occur, for example, in blood vessel repair and formation after trauma, such as after surgery, or during disorders or conditions such as cancer, tumor growth, or neovascularization. Angiogenesis, in which endothelial cells differentiate into endothelial cell tube-like structures that are precursor structures to vessel formation, is an important component of a variety of diseases and disorders including trauma, tumor growth and metastasis, rheumatoid arthritis, psoriasis, atherosclerosis, diabetic retinopathy, retrolental fibroplasia, neovascular glaucoma, age-related macular degeneration, hemangiomas, immune rejection of transplanted corneal tissue and other tissues, and chronic inflammation. By reducing vessel formation, the invention reduces the vasculature supporting a tumor, inhibiting tumor size or growth and reducing the tumor burden of the mammal. Conversely, by enhancing vessel formation, the invention increases or restores the vasculature supporting damaged tissue. Accordingly, the present invention provides means to detect, monitor, analyze, identify, or treat the occurrence or progression of angiogenesis or vasculogenesis in these and other related conditions, and to identify drugs, e.g., antisense, small molecule, antibody, useful to treat these and other related conditions.

Various assays can be used to test the polypeptide herein for angiogenic activity. Such assays include those provided in the Examples below.

Assays for tissue generation activity include, without limitation, those described in WO 95/16035 (bone, cartilage, tendon); WO 95/05846 (nerve, neuronal), and WO 91/07491 (skin, endothelium).

Assays for wound-healing activity include, for example, those described in Winter, Epidermal Wound Healing, Maibach, HI and Rovee, DT, eds. (Year Book Medical Publishers, Inc., Chicago), pp. 71-112, as modified by the article of Eaglstein and Mertz, J. Invest. Dermatol., 71: 382-384 (1978).

Cell-Based Assays

Cell-based assays and animal models for angiogenic disorders, such as tumors, can be used to verify the findings of an angiogenic or angiostatic assay herein, and further to understand the relationship between the genes identified herein and the development and pathogenesis of undesirable angiogenic cell growth. The role of gene products identified herein in the development and pathology of desirable or undesirable angiogenic cell growth, e.g., endothelial cells, tumor cells, can be tested by using cells or cells lines that have been identified as being stimulated or inhibited by the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or its agonists or antagonists, herein. Such cells include, for example, those set forth in the Examples below.

In a different approach, cells of a cell type known to be involved in a particular angiogenic activity or disorder are transfected with the cDNAs herein, and the ability of these cDNAs to induce excessive growth or inhibit growth is analyzed. If the angiogenic disorder is cancer, suitable tumor cells include, for example, stable tumor cells lines such as the B104-1-1 cell line (stable NIH-3T3 cell line transfected with the *neu* protooncogene) and *ras*-transfected NIH-3T3 cells, which can be transfected with the desired gene and

monitored for tumorigenic growth. Such transfected cell lines can then be used to test the ability of poly- or monoclonal antibodies or antibody compositions to inhibit tumorigenic cell growth by exerting cytostatic or cytotoxic activity on the growth of the transformed cells, or by mediating antibody-dependent cellular cytotoxicity (ADCC). Cells transfected with the coding sequences of the genes identified herein can further be used to identify drug candidates for the treatment of angiogenic disorders such as cancer.

In another assay, human umbilical cord endothelial cells (HUVECS) undergoing tube formation in three-dimensional gels in the presence of growth factors, mimic the angiogenic environment of endothelial cells in vivo, providing a well-accepted system for angiogenesis and vasculogenesis, both in normal and neoplastic conditions (see for example Davis, et al. *Exp. Cell Res.* 1996 224:39-51 (1996) and the Examples herein). For example, in one tube formation assay, endothelial cells are suspended in a three-dimensional collagen lattice of type I collagen and undergo rapid morphogenesis. Within 4 hours numerous vacuoles are observed in the majority of endothelial cells. At 24 hours the formation of tube-like structures can be observed. And at 48 hours an interconnected network of tube-like structures is observed. In this and other tube formation assays, inhibitors of protein synthesis (cycloheximide) and mRNA synthesis (actinomycin D) completely block tube-formation. The three dimensional gel is pre-requisite for the differentiation and fusion of endothelial cells into tubes; HUVECS grown on the surface of gelatin or on plastic do not undergo tube-formation. HUVECS can be grown under various conditions, inductive or non-inductive to tube formation, either on gelatin or collagen film (non-inductive) or in collagen gels (inductive), with or without the addition of growth factors to simulate normal angiogenic- or tumor-derived factors. HUVEC cells can be transfected with the cDNAs (or their antisense) herein, and the ability of these nucleic acids to induce excessive growth or tube formation or inhibit growth or tube formation is analyzed. HUVEC cells expressing coding sequences of the genes identified herein can further be used to identify drug candidates. PCR can be used detect the expression of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 mRNA in the endothelial cells cultured in 3D gels, as well as in any other cell or organism. In addition, primary cultures derived from tumors in transgenic animals (as described above) can be used in the cell-based assays herein, although stable cell lines are preferred. Techniques to derive continuous cell lines from transgenic animals are well known in the art. See, e.g., Small *et al.*, Mol. Cell. Biol., 5: 642-648 (1985).

For cancer, a variety of well-known animal models can be used to further understand the role of the genes identified herein in the development and pathogenesis of tumors, and to test the efficacy of candidate therapeutic agents, including antibodies and other antagonists of the native PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides, such as small-molecule antagonists. The *in vivo* nature of such models makes them particularly predictive of responses in human patients. Animal models of tumors and cancers (e.g., breast cancer, colon cancer, prostate cancer, lung cancer, etc.) include both non-recombinant and recombinant (transgenic) animals. Non-recombinant animal models include, for example, rodent, e.g., murine models. Such models can be generated by introducing tumor cells into syngeneic mice using standard techniques, e.g., subcutaneous injection, tail vein injection, spleen implantation, intraperitoneal implantation, implantation under the renal capsule, or orthotopic implantation, e.g., colon cancer cells implanted in colonic tissue. See, e.g., PCT publication No. WO 97/33551, published September 18, 1997. Probably the

most often used animal species in oncological studies are immunodeficient mice and, in particular, nude mice. The observation that the nude mouse with thymic hypo/aplasia could successfully act as a host for human tumor xenografts has lead to its widespread use for this purpose. The autosomal recessive *nu* gene has been introduced into a very large number of distinct congenic strains of nude mouse, including, for example, ASW, A/He, AKR, BALB/c, B10.LP, C17, C3H, C57BL, C57, CBA, DBA, DDD, I/st, NC, NFR, NFS, NFS/N, NZB, NZC, NZW, P, RIII, and SJL. In addition, a wide variety of other animals with inherited immunological defects other than the nude mouse have been bred and used as recipients of tumor xenografts. For further details see, e.g., The Nude Mouse in Oncology Research, E. Boven and B. Winograd, eds. (CRC Press, Inc., 1991).

The cells introduced into such animals can be derived from known tumor/cancer cell lines, such as any of the above-listed tumor cell lines, and, for example, the B104-1-1 cell line (stable NIH-3T3 cell line transfected with the *neu* protooncogene); *ras*-transfected NIH-3T3 cells; Caco-2 (ATCC HTB-37); or a moderately well-differentiated grade II human colon adenocarcinoma cell line, HT-29 (ATCC HTB-38); or from tumors and cancers. Samples of tumor or cancer cells can be obtained from patients undergoing surgery, using standard conditions involving freezing and storing in liquid nitrogen. Karmali *et al.*, Br. J. Cancer, **48**: 689-696 (1983).

Tumor cells can be introduced into animals such as nude mice by a variety of procedures. The subcutaneous (s.c.) space in mice is very suitable for tumor implantation. Tumors can be transplanted s.c. as solid blocks, as needle biopsies by use of a trochar, or as cell suspensions. For solid-block or trochar implantation, tumor tissue fragments of suitable size are introduced into the s.c. space. Cell suspensions are freshly prepared from primary tumors or stable tumor cell lines, and injected subcutaneously. Tumor cells can also be injected as subdermal implants. In this location, the inoculum is deposited between the lower part of the dermal connective tissue and the s.c. tissue.

Animal models of breast cancer can be generated, for example, by implanting rat neuroblastoma cells (from which the *neu* oncogene was initially isolated), or *neu*-transformed NIH-3T3 cells into nude mice, essentially as described by Drebin *et al.* Proc. Nat. Acad. Sci. USA, **83**: 9129-9133 (1986).

Similarly, animal models of colon cancer can be generated by passaging colon cancer cells in animals, e.g., nude mice, leading to the appearance of tumors in these animals. An orthotopic transplant model of human colon cancer in nude mice has been described, for example, by Wang *et al.*, Cancer Research, **54**: 4726-4728 (1994) and Too *et al.*, Cancer Research, **55**: 681-684 (1995). This model is based on the so-called "METAMOUSE"TM sold by AntiCancer, Inc., (San Diego, California).

Tumors that arise in animals can be removed and cultured *in vitro*. Cells from the *in vitro* cultures can then be passaged to animals. Such tumors can serve as targets for further testing or drug screening. Alternatively, the tumors resulting from the passage can be isolated and RNA from pre-passage cells and cells isolated after one or more rounds of passage analyzed for differential expression of genes of interest. Such passaging techniques can be performed with any known tumor or cancer cell lines.

For example, Meth A, CMS4, CMS5, CMS21, and WEHI-164 are chemically induced fibrosarcomas of BALB/c female mice (DeLeo *et al.*, J. Exp. Med., **146**: 720 (1977)), which provide a highly controllable model system for studying the anti-tumor activities of various agents. Palladino *et al.*, J. Immunol., **138**: 4023-

4032 (1987). Briefly, tumor cells are propagated *in vitro* in cell culture. Prior to injection into the animals, the cell lines are washed and suspended in buffer, at a cell density of about 10×10^6 to 10×10^7 cells/ml. The animals are then infected subcutaneously with 10 to 100 μ l of the cell suspension, allowing one to three weeks for a tumor to appear.

5 In addition, the Lewis lung (3LL) carcinoma of mice, which is one of the most thoroughly studied experimental tumors, can be used as an investigational tumor model. Efficacy in this tumor model has been correlated with beneficial effects in the treatment of human patients diagnosed with small-cell carcinoma of the lung (SCCL). This tumor can be introduced in normal mice upon injection of tumor fragments from an affected mouse or of cells maintained in culture. Zupi *et al.*, Br. J. Cancer, **41**: suppl. 4, 30 (1980). Evidence indicates
10 that tumors can be started from injection of even a single cell and that a very high proportion of infected tumor cells survive. For further information about this tumor model see, Zacharski, Haemostasis, **16**: 300-320 (1986).

One way of evaluating the efficacy of a test compound in an animal model with an implanted tumor is to measure the size of the tumor before and after treatment. Traditionally, the size of implanted tumors has been measured with a slide caliper in two or three dimensions. The measure limited to two dimensions does not
15 accurately reflect the size of the tumor; therefore, it is usually converted into the corresponding volume by using a mathematical formula. However, the measurement of tumor size is very inaccurate. The therapeutic effects of a drug candidate can be better described as treatment-induced growth delay and specific growth delay. Another important variable in the description of tumor growth is the tumor volume doubling time. Computer programs for the calculation and description of tumor growth are also available, such as the program reported by Rygaard and Spang-Thomsen, Proc. 6th Int. Workshop on Immune-Deficient Animals, Wu and Sheng eds. (Basel,
20 1989), p. 301. It is noted, however, that necrosis and inflammatory responses following treatment may actually result in an increase in tumor size, at least initially. Therefore, these changes need to be carefully monitored, by a combination of a morphometric method and flow cytometric analysis.

Further, recombinant (transgenic) animal models can be engineered by introducing the coding portion
25 of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 genes identified herein into the genome of animals of interest, using standard techniques for producing transgenic animals. Animals that can serve as a target for transgenic manipulation include, without limitation, mice, rats, rabbits, guinea pigs, sheep, goats, pigs, and non-human primates, *e.g.*, baboons, chimpanzees and monkeys. Techniques known in the art to introduce a transgene into such animals include pronucleic microinjection (U.S. Patent No.
30 4,873,191); retrovirus-mediated gene transfer into germ lines (*e.g.*, Van der Putten *et al.*, Proc. Natl. Acad. Sci. USA, **82**: 6148-615 (1985)); gene targeting in embryonic stem cells (Thompson *et al.*, Cell, **56**: 313-321 (1989)); electroporation of embryos (Lo, Mol. Cell. Biol., **3**: 1803-1814 (1983)); and sperm-mediated gene transfer. Lavitrano *et al.*, Cell, **57**: 717-73 (1989). For a review, see for example, U.S. Patent No. 4,736,866.

For the purpose of the present invention, transgenic animals include those that carry the transgene only
35 in part of their cells ("mosaic animals"). The transgene can be integrated either as a single transgene, or in concatamers, *e.g.*, head-to-head or head-to-tail tandems. Selective introduction of a transgene into a particular cell type is also possible by following, for example, the technique of Lasko *et al.*, Proc. Natl. Acad. Sci. USA, **89**: 6232-636 (1992). The expression of the transgene in transgenic animals can be monitored by standard

techniques. For example, Southern blot analysis or PCR amplification can be used to verify the integration of the transgene. The level of mRNA expression can then be analyzed using techniques such as *in situ* hybridization, Northern blot analysis, PCR, or immunocytochemistry. The animals are further examined for signs of tumor or cancer development.

5 Alternatively, "knock-out" animals can be constructed that have a defective or altered gene encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide identified herein, as a result of homologous recombination between the endogenous gene encoding the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide and altered genomic DNA encoding the same polypeptide introduced into an embryonic cell of the animal. For example, cDNA encoding a particular
10 PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide can be used to clone genomic DNA encoding that polypeptide in accordance with established techniques. A portion of the genomic DNA encoding a particular PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide can be deleted or replaced with another gene, such as a gene encoding a selectable marker that can be used to monitor integration. Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector. See, *e.g.*, Thomas and Capecchi, Cell, 51: 503 (1987) for a description of homologous recombination vectors. The vector is introduced into an embryonic stem cell line (*e.g.*, by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected. See, *e.g.*, Li *et al.*, Cell, 69: 915 (1992). The selected cells are then injected into a blastocyst of an animal (*e.g.*, a mouse or rat) to form aggregation chimeras. See, *e.g.*, Bradley, in Teratocarcinomas and
20 Embryonic Stem Cells: A Practical Approach, E. J. Robertson, ed. (IRL: Oxford, 1987), pp. 113-152. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a "knock-out" animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knock-out animals can also be generated, as is well known in the art, by administering an antisense molecule of the invention. Animals comprising such antisense molecules are specifically contemplated as an embodiment of the invention. Knockout animals can be characterized, for instance, by their ability to defend against certain pathological conditions and by their development of pathological conditions due to absence (knock-out) of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

30 The efficacy of antibodies specifically binding the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides identified herein, and other drug candidates, can be tested also in the treatment of spontaneous animal tumors. A suitable target for such studies is the feline oral squamous cell carcinoma (SCC). Feline oral SCC is a highly invasive, malignant tumor that is the most common oral malignancy of cats, accounting for over 60% of the oral tumors reported in this species. It rarely metastasizes to distant sites, although this low incidence of metastasis may merely be a reflection of the short survival times for cats with this tumor. These tumors are usually not amenable to surgery, primarily because of the anatomy of the feline oral cavity. At present, there is no effective treatment for this tumor. Prior to entry into the study, each cat undergoes complete clinical examination and biopsy, and is scanned by computed tomography (CT). Cats

diagnosed with sublingual oral squamous cell tumors are excluded from the study. The tongue can become paralyzed as a result of such tumor, and even if the treatment kills the tumor, the animals may not be able to feed themselves. Each cat is treated repeatedly, over a longer period of time. Photographs of the tumors will be taken daily during the treatment period, and at each subsequent recheck. After treatment, each cat undergoes another CT scan. CT scans and thoracic radiograms are evaluated every 8 weeks thereafter. The data are evaluated for differences in survival, response, and toxicity as compared to control groups. Positive response may require evidence of tumor regression, preferably with improvement of quality of life and/or increased life span.

In addition, other spontaneous animal tumors, such as fibrosarcoma, adenocarcinoma, lymphoma, chondroma, or leiomyosarcoma of dogs, cats, and baboons can also be tested. Of these, mammary adenocarcinoma in dogs and cats is a preferred model as its appearance and behavior are very similar to those in humans. However, the use of this model is limited by the rare occurrence of this type of tumor in animals. Other *in vitro* and *in vivo* angiogenic tests known in the art are also suitable herein.

H. Anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 Antibodies

The present invention further provides anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibodies. Exemplary antibodies include polyclonal, monoclonal, humanized, bispecific, and heteroconjugate antibodies.

1. Polyclonal Antibodies

The anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibodies can comprise polyclonal antibodies. Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent can include the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or a fusion protein thereof. It can be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants which can be employed include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). The immunization protocol can be selected by one skilled in the art without undue experimentation.

2. Monoclonal Antibodies

The anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibodies can, alternatively, be monoclonal antibodies. Monoclonal antibodies can be prepared using hybridoma methods, such as those described by Kohler and Milstein, Nature, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing

agent. Alternatively, the lymphocytes can be immunized *in vitro*.

The immunizing agent will typically include the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or a fusion protein thereof. Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell [Goding, Monoclonal Antibodies: Principles and Practice, Academic Press, (1986) pp. 59-103]. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells can be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Manassas, Virginia. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies [Kozbor, J. Immunol., 133:3001 (1984); Brodeur et al., Monoclonal Antibody Production Techniques and Applications, Marcel Dekker, Inc., New York, (1987) pp. 51-63].

The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980).

After the desired hybridoma cells are identified, the clones can be subcloned by limiting dilution procedures and grown by standard methods [Goding, supra]. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells can be grown *in vivo* as ascites in a mammal.

The monoclonal antibodies secreted by the subclones can be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

The monoclonal antibodies can also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding

specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA can be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also can be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences [U.S. Patent No. 4,816,567; Morrison et al., supra] or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

The antibodies can be monovalent antibodies. Methods for preparing monovalent antibodies are well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy chain crosslinking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

In vitro methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to produce fragments thereof, particularly, Fab fragments, can be accomplished using routine techniques known in the art.

3. Human and Humanized Antibodies

The anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibodies of the invention can further comprise humanized antibodies or human antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies can also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin [Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-329 (1988); and Presta, Curr. Op. Struct. Biol., 2:593-596 (1992)].

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an

"import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers [Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-327 (1988); Verhoeyen et al., *Science*, 239:1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

Human antibodies can also be produced using various techniques known in the art, including phage display libraries [Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks et al., *J. Mol. Biol.*, 222:581 (1991)]. The techniques of Cole et al. and Boerner et al. are also available for the preparation of human monoclonal antibodies (Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985) and Boerner et al., *J. Immunol.*, 147(1):86-95 (1991)]. Similarly, human antibodies can be made by introducing of human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following scientific publications: Marks *et al.*, *Bio/Technology* 10, 779-783 (1992); Lonberg *et al.*, *Nature* 368 856-859 (1994); Morrison, *Nature* 368, 812-13 (1994); Fishwild *et al.*, *Nature Biotechnology* 14, 845-51 (1996); Neuberger, *Nature Biotechnology* 14, 826 (1996); Lonberg and Huszar, *Intern. Rev. Immunol.* 13 65-93 (1995).

4. Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, the other one is for any other antigen, and preferably for a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities [Milstein and Cuello, *Nature*, 305:537-539 (1983)]. Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker et al., *EMBO J.*, 10:3655-3659 (1991).

Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least

one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh et al., Methods in Enzymology, 121:210 (1986).

5 According to another approach described in WO 96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 region of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory “cavities”
10 of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')₂ bispecific antibodies). Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan *et al.*, Science 229:81 (1985) describe a procedure wherein intact antibodies are
15 proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of
20 the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Fab' fragments can be directly recovered from *E. coli* and chemically coupled to form bispecific antibodies. Shalaby *et al.*, J. Exp. Med. 175:217-225 (1992) describe the production of a fully humanized
25 bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling *in vitro* to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

Various technique for making and isolating bispecific antibody fragments directly from recombinant cell
30 culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny *et al.*, J. Immunol. 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The “diabody” technology
35 described by Hollinger *et al.*, Proc. Natl. Acad. Sci. USA 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker which is too short to allow pairing between the

two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby forming two antigen-binding sites.

Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See, Gruber *et al.*, J. Immunol. 152:5368 (1994).

5 Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt *et al.*, J. Immunol. 147:60 (1991).

Exemplary bispecific antibodies can bind to two different epitopes on a given PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide herein. Alternatively, an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide arm can be combined
10 with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g. CD2, CD3, CD28, or B7), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16) so as to focus cellular defense mechanisms to the cell expressing the particular PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide. Bispecific antibodies can also be used to localize cytotoxic agents to cells which express a particular PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45,
15 PRO-C-MG.64 or PRO-C-MG.72 polypeptide. These antibodies possess a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-binding arm and an arm which binds a cytotoxic agent or a radionuclide chelator, such as EOTUBE, DPTA, DOTA, or TETA. Another bispecific antibody of interest binds the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide and further binds tissue factor (TF).

20 5. Heteroconjugate Antibodies

Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360; WO 92/200373; EP 03089). It is contemplated that the antibodies can be prepared *in vitro* using
25 known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins can be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Patent No. 4,676,980.

6. Effector Function Engineering

30 It can be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, *e.g.*, the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) can be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated can have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron *et al.*, J. Exp Med., 176: 1191-1195 (1992) and Shopes, J. Immunol., 148: 2918-2922 (1992). Homodimeric antibodies
35 with enhanced anti-tumor activity can also be prepared using heterobifunctional cross-linkers as described in Wolff *et al.* Cancer Research, 53: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and can thereby have enhanced complement lysis and ADCC capabilities. See Stevenson *et al.*,

Anti-Cancer Drug Design, 3: 219-230 (1989).

7. Immunoconjugates

The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (*e.g.*, an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (*i.e.*, a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcumin, croton, saponaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include ^{212}Bi , ^{131}I , ^{131}In , ^{90}Y , and ^{186}Re .

Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta *et al.*, Science, **238**: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminopentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026.

In another embodiment, the antibody can be conjugated to a "receptor" (such streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (*e.g.*, avidin) that is conjugated to a cytotoxic agent (*e.g.*, a radionucleotide).

8. Immunoliposomes

The antibodies disclosed herein can also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein *et al.*, Proc. Natl. Acad. Sci. USA, **82**: 3688 (1985); Hwang *et al.*, Proc. Natl. Acad. Sci. USA, **77**: 4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin *et al.*, J. Biol. Chem., **257**: 286-288 (1982) via a disulfide-interchange reaction. A chemotherapeutic agent (such as Doxorubicin) is optionally contained within the liposome. See Gabizon *et al.*,

J. National Cancer Inst., **81**(19): 1484 (1989).

9. Pharmaceutical Compositions of Antibodies

Antibodies specifically binding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide identified herein, as well as other molecules identified by the screening assays disclosed herein, can be administered for the treatment of various disorders in the form of pharmaceutical compositions.

If the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide is intracellular and whole antibodies are used as inhibitors, internalizing antibodies are preferred. However, lipofections or liposomes can also be used to deliver the antibody, or an antibody fragment, into cells. Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein is preferred. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. See, *e.g.*, Marasco *et al.*, Proc. Natl. Acad. Sci. USA, **90**: 7889-7893 (1993). The formulation herein can also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition can comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

The active ingredients can also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, *supra*.

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

Sustained-release preparations can be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, *e.g.*, films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOTTM (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they can denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange,

stabilization can be achieved by modifying sulphydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

I. Uses for anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72

Antibodies

5 The anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibodies of the invention have various utilities. For example, anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibodies can be used in diagnostic assays for PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, *e.g.*, detecting its expression in specific cells, tissues, or serum. Various diagnostic assay techniques known in the art can be used, such as competitive binding assays, direct or indirect sandwich assays and immunoprecipitation assays conducted in either heterogeneous or homogeneous phases (Zola, Monoclonal Antibodies: A Manual of Techniques, CRC Press, Inc. (1987) pp. 147-158). The antibodies used in the diagnostic assays can be labeled with a detectable moiety. The detectable moiety should be capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety can be a radioisotope, such as ^3H , ^{14}C , ^{32}P , ^{35}S , or ^{125}I , a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin, or an enzyme, such as alkaline phosphatase, beta-galactosidase or horseradish peroxidase. Any method known in the art for conjugating the antibody to the detectable moiety can be employed, including those methods described by Hunter et al., Nature, 144:945 (1962); David et al., Biochemistry, 13:1014 (1974); Pain et al., J. Immunol. Meth., 40:219 (1981); and Nygren, J. Histochem. and Cytochem., 30:407 (1982).

20 Anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibodies also are useful for the affinity purification of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 from recombinant cell culture or natural sources. In this process, the antibodies against PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 are immobilized on a suitable support, such as a Sephadex resin or filter paper, using methods well known in the art. The immobilized antibody then is contacted with a sample containing the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 to be purified, and thereafter the support is washed with a suitable solvent that will remove substantially all the material in the sample except the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, which is bound to the immobilized antibody. Finally, the support is washed with another suitable solvent that will release the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 from the antibody.

J. Use of Gene as Diagnostic

This invention is also related to the use of the gene encoding the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide as a diagnostic. Detection of a mutated form of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide will allow a diagnosis of an angiogenic disease or a susceptibility to an angiogenic disease, such as a tumor, since mutations in the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide may cause tumors.

5

10

25

30

3.

3.

support and the labeled PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide and an appropriately processed sample derived from the subject are passed over the solid support, wherein the amount of label detected attached to the solid support is correlated to a quantity of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide in the sample.

- 5 Since expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 mRNA is correlated with angiogenesis as disclosed herein, in another embodiment a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 specific nucleic acid of the invention can be used in an RNA detection or quantification method, such as in situ hybridization or PCR amplification, to diagnose or detect vascular disease or neovascularization associated with tumor formation.

10 L. Types of Angiogenic Disorders to be Treated

- The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides, or agonists or antagonists thereto, that have activity in the cardiovascular, angiogenic, and endothelial assays described herein, are likely to have therapeutic uses in a variety of angiogenic disorders, including systemic disorders that affect vessels, such as diabetes mellitus. Their therapeutic utility could include diseases of the arteries, capillaries, veins, and/or lymphatics. The compounds of the invention thus have use in treatment of diseases or disorders characterized by undesirable excessive neovascularization. Vascular or angiogenic dysfunction further includes diseases of the vessels themselves, such as of the arteries, capillaries, veins, and/or lymphatics. This would include indications that stimulate angiogenesis, cardiovascularization, and/or neovascularization, and those that inhibit angiogenesis, cardiovascularization, and/or neovascularization. Such disorders include, for example, arterial disease, such as atherosclerosis, hypertension, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, and arterial restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, cancer such as vascular tumors, e.g., hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, Kaposi's sarcoma, lymphangioma, and lymphangiosarcoma, tumor angiogenesis, trauma such as wounds, burns, and other injured tissue, implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, psoriasis, retinopathy, retrolental fibroplasia, neovascular glaucoma, age-related macular degeneration, thyroid hyperplasias, Grave's disease, tissue transplantation, chronic inflammation, lung inflammation, obesity, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. This would also include angina, myocardial infarctions such as acute myocardial infarctions and heart failure such as congestive heart failure.

- The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides or agonists or antagonists thereto may also be employed to stimulate wound healing or tissue regeneration and associated therapies concerned with re-growth of tissue, such as connective tissue, skin, bone, cartilage, muscle, lung, or kidney, to promote angiogenesis, and to proliferate the growth of vascular smooth muscle and endothelial cell production, and improving allograft and xenograft success. The increase in angiogenesis is mediated by the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or antagonist would be beneficial to ischemic tissues and to collateral coronary development in the heart

subsequent to coronary stenosis. Antagonists are used to inhibit the action of such polypeptides, for example, to limit the production of excess connective tissue during wound healing or pulmonary fibrosis if the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide promotes such production. This would include treatment of acute myocardial infarction and heart failure, other trauma of the vasculature, and muscle wasting disease.

Moreover, the present invention concerns the treatment of cardiac hypertrophy, regardless of the underlying cause, by administering a therapeutically effective dose of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or agonist or antagonist thereto. If the objective is the treatment of human patients, the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide preferably is recombinant human PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide (rhPRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 or rhPRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide). The treatment for cardiac hypertrophy can be performed at any of its various stages, which may result from a variety of diverse pathologic conditions, including myocardial infarction, hypertension, hypertrophic cardiomyopathy, and valvular regurgitation. The treatment extends to all stages of the progression of cardiac hypertrophy, with or without structural damage of the heart muscle, regardless of the underlying cardiac disorder.

The decision of whether to use the molecule itself or an agonist thereof for any particular indication, as opposed to an antagonist to the molecule, would depend mainly on whether the molecule herein promotes cardiovascularization, genesis of endothelial cells, or angiogenesis or inhibits these conditions. For example, if the molecule promotes angiogenesis, an antagonist thereof would be useful for treatment of disorders where it is desired to limit or prevent angiogenesis. Examples of such disorders include vascular tumors such as haemangioma, tumor angiogenesis, neovascularization in the retina, choroid, or cornea, associated with diabetic retinopathy or premature infant retinopathy or macular degeneration and proliferative vitreoretinopathy, rheumatoid arthritis, Crohn's disease, atherosclerosis, ovarian hyperstimulation, psoriasis, endometriosis associated with neovascularization, restenosis subsequent to balloon angioplasty, scar tissue overproduction, for example, that seen in a keloid that forms after surgery, fibrosis after myocardial infarction, or fibrotic lesions associated with pulmonary fibrosis.

Excessive endometrial angiogenesis has been proposed as an important mechanism in the pathogenesis of endometriosis. The endometrium of women with endometriosis has an increased capacity to proliferate, implant and grow in the peritoneal cavity. The endometrium of patients with endometriosis shows enhanced endothelial cell proliferation. Cell adhesion molecule integrin $\alpha v \beta 3$ is expressed in more blood vessels in the endometrium of women with endometriosis when compared with normal women. Taken together, these results provide evidence for increased endometrial angiogenesis in women with endometriosis when compared with normal subjects (Healy et al. *Hum. Reprod. Update* 4(5):736-40 (1998)). Endometriosis is one of the family of angiogenic diseases, as discussed herein. Inhibition of angiogenesis as taught herein will provide benefit in treating such a disease.

If, however, the molecule inhibits angiogenesis, it would be expected to be used directly for treatment of

the above conditions.

On the other hand, if the molecule stimulates angiogenesis it would be used itself (or an agonist thereof) for indications where angiogenesis is desired such as peripheral vascular disease, hypertension, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, arterial restenosis, thrombophlebitis, lymphangitis, lymphedema, wound healing and tissue repair, ischemia reperfusion injury, angina, myocardial infarctions such as acute myocardial infarctions, chronic heart conditions, heart failure such as congestive heart failure, and osteoporosis.

If, however, the molecule inhibits angiogenesis, an antagonist thereof would be used for treatment of those conditions where angiogenesis is desired.

Specific types of diseases are described herein, where the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or antagonists thereof may serve as useful for vascular-related drug targeting or as therapeutic targets for the treatment or prevention of the disorders. Atherosclerosis is a disease characterized by accumulation of plaques of intimal thickening in arteries, due to accumulation of lipids, proliferation of smooth muscle cells, and formation of fibrous tissue within the arterial wall. The disease can affect large, medium, and small arteries in any organ. Changes in endothelial and vascular smooth muscle cell function are known to play an important role in modulating the accumulation and regression of these plaques.

Hypertension is characterized by raised vascular pressure in the systemic arterial, pulmonary arterial, or portal venous systems. Elevated pressure may result from or result in impaired endothelial function and/or vascular disease.

Inflammatory vasculitides include giant cell arteritis, Takayasu's arteritis, polyarteritis nodosa (including the microangiopathic form), Kawasaki's disease, microscopic polyangiitis, Wegener's granulomatosis, and a variety of infectious-related vascular disorders (including Henoch-Schonlein purpura). Altered endothelial cell function has been shown to be important in these diseases.

Reynaud's disease and Reynaud's phenomenon are characterized by intermittent abnormal impairment of the circulation through the extremities on exposure to cold. Altered endothelial cell function has been shown to be important in this disease.

Aneurysms are saccular or fusiform dilatations of the arterial or venous tree that are associated with altered endothelial cell and/or vascular smooth muscle cells.

Arterial restenosis (restenosis of the arterial wall) may occur following angioplasty as a result of alteration in the function and proliferation of endothelial and vascular smooth muscle cells.

Thrombophlebitis and lymphangitis are inflammatory disorders of veins and lymphatics, respectively, that may result from, and/or in, altered endothelial cell function. Similarly, lymphedema is a condition involving impaired lymphatic vessels resulting from endothelial cell function.

Another use for the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides herein or antagonists thereto is in the prevention or treatment of cancer, and preferably vascular tumors. Examples of cancer include but are not limited to, carcinoma including adenocarcinoma, lymphoma, blastoma, melanoma, sarcoma, and leukemia. More particular examples of such cancers include squamous cell cancer, small-cell lung cancer, non-small cell lung cancer, gastrointestinal cancer, Hodgkin's and non-

Hodgkin's lymphoma, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer such as hepatic carcinoma and hepatoma, bladder cancer, breast cancer, colon cancer, colorectal cancer, endometrial carcinoma, salivary gland carcinoma, kidney cancer such as renal cell carcinoma and Wilms' tumors, basal cell carcinoma, melanoma, prostate cancer, vulval cancer, thyroid cancer, testicular cancer, esophageal cancer, and various types of head and neck cancer. The preferred cancers for treatment herein are breast, colon, lung, melanoma, ovarian, and others involving vascular tumors as noted above of tumor angiogenesis, which involves vascularization of a tumor to enable it to grow and/or metastasize. This process is dependent on the growth of new blood vessels. Further examples of preferred neoplasms and related conditions that involve tumor angiogenesis include breast carcinomas, gastric carcinomas, esophageal carcinomas, colorectal carcinomas, liver carcinomas, thecomas, arrhenoblastomas, cervical carcinomas, endometrial carcinoma, endometrial hyperplasia, endometriosis, fibrosarcomas, choriocarcinoma, nasopharyngeal carcinoma, laryngeal carcinomas, hepatoblastoma, Kaposi's sarcoma, melanoma, skin carcinomas, hemangioma, cavernous hemangioma, hemangioblastoma, pancreas carcinomas, retinoblastoma, astrocytoma, glioblastoma, Schwannoma, oligodendroglioma, medulloblastoma, neuroblastomas, rhabdomyosarcoma, osteogenic sarcoma, leiomyosarcomas, urinary tract carcinomas, thyroid carcinomas, Wilm's tumor, renal cell carcinoma, prostate carcinoma, abnormal vascular proliferation associated with phakomatoses, edema (such as that associated with brain tumors), and Meigs' syndrome. The family of benign vascular tumors, also included herein for treatment, are also characterized by abnormal proliferation and growth of cellular elements of the vascular system. For example, lymphangiomas are benign tumors of the lymphatic system that are congenital, often cystic, malformations of the lymphatics that usually occur in newborns. Cystic tumors tend to grow into the adjacent tissue. Cystic tumors usually occur in the cervical and axillary region. They can also occur in the soft tissue of the extremities. The main symptoms are dilated, sometimes reticular, structured lymphatics and lymphocysts surrounded by connective tissue. Lymphangiomas are assumed to be caused by improperly connected embryonic lymphatics or their deficiency. The result is impaired local lymph drainage (Griener *et al.*, *Lymphology* 4:140-144 (1971)).

In one embodiment of the method of the invention, a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antagonist is administered to a mammal, e.g. a human patient, in need thereof to reduce the tumor burden in the mammal. The compound, by inhibiting angiogenesis, is useful for the treatment of diseases or disorders characterized by undesirable excessive neovascularization, including by way of example tumors, and especially solid malignant tumors as mentioned herein, and non-neoplastic disorders including angina, myocardial infarctions such as acute myocardial infarctions, and heart failure such as congestive heart failure, psoriasis, diabetic and other proliferative retinopathies including retinopathy of prematurity, retrolental fibroplasia, neovascular glaucoma, thyroid hyperplasias (including Grave's disease), corneal and other tissue transplantation, chronic inflammation, lung inflammation, nephrotic syndrome, preeclampsia, ascites, pericardial effusion (such as that associated with pericarditis), pleural effusion, rheumatoid arthritis, atherosclerosis, hemangiomas, obesity, and age-related macular degeneration.

Age-related macular degeneration (AMD) is a leading cause of severe visual loss in the elderly population. The exudative form of AMD is characterized by choroidal neovascularization and retinal pigment epithelial cell

detachment. Because choroidal neovascularization is associated with a dramatic worsening in prognosis, the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides or antagonist thereto is expected to be useful in reducing the severity of AMD.

Healing of trauma such as wound healing and tissue repair is also a targeted use for the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides herein or their antagonists. Formation and regression of new blood vessels is essential for tissue healing and repair. This category includes bone, cartilage, tendon, ligament, and/or nerve tissue growth or regeneration, as well as wound healing and tissue repair and replacement, and in the treatment of burns, incisions, and ulcers. A PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or antagonist thereof that induces cartilage and/or bone growth in circumstances where bone is not normally formed has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or antagonist thereof may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. *De novo* bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma-induced, or oncologic, resection-induced craniofacial defects, and also is useful in cosmetic plastic surgery. PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides or antagonists thereto may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

It is expected that a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or antagonist thereto may also exhibit activity for generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, or endothelium), muscle (smooth, skeletal, or cardiac), and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring to allow normal tissue to regenerate.

A PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide herein or antagonist thereto may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage. Also, the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or antagonist thereto may be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells, or for inhibiting the growth of tissues described above.

A PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or antagonist thereto may also be used in the treatment of periodontal diseases and in other tooth-repair processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells, or induce differentiation of progenitors of bone-forming cells. A PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide herein or an antagonist thereto may also be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue destruction (collagenase activity, osteoclast activity, etc.) mediated

by inflammatory processes, since blood vessels play an important role in the regulation of bone turnover and growth.

Another category of tissue regeneration activity that may be attributable to the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide herein or antagonist thereto is

tendon/ligament formation. A protein that induces tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed has application in the healing of tendon or ligament tears, deformities, and other tendon or ligament defects in humans and other animals. Such a preparation may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue. *De novo* tendon/ligament-like tissue formation induced by a composition of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide herein or antagonist thereto contributes to the repair of congenital, trauma-induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions herein may provide an environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions herein may also be useful in the treatment of tendinitis, carpal tunnel syndrome, and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

Ischemia-reperfusion injury is another indication. Endothelial cell dysfunction may be important in both the initiation of, and in regulation of the sequelae of events that occur following ischemia-reperfusion injury.

Rheumatoid arthritis is a further indication. Blood vessel growth and targeting of inflammatory cells through the vasculature is an important component in the pathogenesis of rheumatoid and sero-negative forms of arthritis.

In view of the above, the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides or agonists or antagonists thereof described herein, which are shown to alter or impact endothelial cell function, proliferation, and/or form, are likely to play an important role in the etiology and pathogenesis of many or all of the disorders noted above, and as such can serve as therapeutic targets to augment or inhibit these processes or for vascular-related drug targeting in these disorders.

M. Administration Protocols, Schedules, Doses, and Formulations

The molecules herein and agonists and antagonists thereto, including antigene compounds, are pharmaceutically useful as a prophylactic and therapeutic agent for various disorders and diseases as set forth above. Antigene compounds, such as antisense oligonucleotides, are more preferably formulated and administered as discussed above.

Therapeutic compositions of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides or agonists or antagonists are prepared for storage by mixing the desired molecule having the appropriate degree of purity with optional pharmaceutically acceptable carriers, excipients, or stabilizers (Remington's Pharmaceutical Sciences, 16th edition, Osol, A. ed. (1980)), in the form of lyophilized

formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrins; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; lipids such as cationic lipids; salt-forming counter-ions such as sodium; metal complexes (*e.g.*, Zn-protein complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG).

Additional examples of such carriers include ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts, or electrolytes such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, and polyethylene glycol. Carriers for topical or gel-based forms of antagonist include polysaccharides such as sodium carboxymethylcellulose or methylcellulose, polyvinylpyrrolidone, polyacrylates, polyoxyethylene-polyoxypropylene-block polymers, polyethylene glycol, and wood wax alcohols. For all administrations, conventional depot forms are suitably used. Such forms include, for example, microcapsules, nano-capsules, liposomes, plasters, inhalation forms, nose sprays, sublingual tablets, and sustained-release preparations. The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides or agonists or antagonists will typically be formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml.

PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or antagonist to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to or following lyophilization and reconstitution. Therapeutic compositions herein generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle. PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide ordinarily will be stored in lyophilized form or in solution if administered systemically. If in lyophilized form, PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or antagonist thereto is typically formulated in combination with other ingredients for reconstitution with an appropriate diluent at the time for use. An example of a liquid formulation of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or antagonist is a sterile, clear, colorless unpreserved solution filled in a single-dose vial for subcutaneous injection. Preserved pharmaceutical compositions suitable for repeated use may contain, for example, depending mainly on the indication and type of polypeptide: a) PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or agonist or antagonist thereto; b) a buffer

capable of maintaining the pH in a range of maximum stability of the polypeptide or other molecule in solution, preferably about 4-8; c) a detergent/surfactant primarily to stabilize the polypeptide or molecule against agitation-induced aggregation; d) an isotonifier; e) a preservative selected from the group of phenol, benzyl alcohol and a benzethonium halide, *e.g.*, chloride; and f) water.

5 If the detergent employed is non-ionic, it may, for example, be polysorbates (*e.g.*, POLYSORBATE™ (TWEEN™) 20, 80, etc.) or poloxamers (*e.g.*, POLOXAMER™ 188). The use of non-ionic surfactants permits the formulation to be exposed to shear surface stresses without causing denaturation of the polypeptide. Further, such surfactant-containing formulations may be employed in aerosol devices such as those used in a pulmonary dosing, and needleless jet injector guns (see, *e.g.*, EP 257,956).

10 An isotonifier may be present to ensure isotonicity of a liquid composition of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or antagonist thereto, and includes polyhydric sugar alcohols, preferably trihydric or higher sugar alcohols, such as glycerin, erythritol, arabitol, xylitol, sorbitol, and mannitol. These sugar alcohols can be used alone or in combination. Alternatively, sodium chloride or other appropriate inorganic salts may be used to render the solutions isotonic.

15 The buffer may, for example, be an acetate, citrate, succinate, or phosphate buffer depending on the pH desired. The pH of one type of liquid formulation of this invention is buffered in the range of about 4 to 8, preferably about physiological pH.

The preservatives phenol, benzyl alcohol and benzethonium halides, *e.g.*, chloride, are known antimicrobial agents that may be employed.

20 Examples of pharmacologically acceptable salts of molecules that form salts and are useful hereunder include alkali metal salts (*e.g.*, sodium salt, potassium salt), alkaline earth metal salts (*e.g.*, calcium salt, magnesium salt), ammonium salts, organic base salts (*e.g.*, pyridine salt, triethylamine salt), inorganic acid salts (*e.g.*, hydrochloride, sulfate, nitrate), and salts of organic acid (*e.g.*, acetate, oxalate, p-toluenesulfonate).

25 The route of administration is in accord with known methods, *e.g.* injection or infusion by intravenous, intraperitoneal, intracerebral, intracerebrospinal, intraocular, intraarticular, intrasynovial, intrathecal, intraarterial or intralesional routes, oral, topical administration, or by sustained release systems. The compounds of the invention are also suitably administered by intratumoral, peritumoral, intralesional, or perilesional routes, to exert local as well as systemic therapeutic effects. The intraperitoneal route is expected to be particularly useful, for example, in the treatment of ovarian tumors. The formulations can also be administered as repeated
30 intravenous (*i.v.*), subcutaneous (*s.c.*), or intramuscular (*i.m.*) injections, or as aerosol formulations suitable for intranasal or intrapulmonary delivery (for intrapulmonary delivery see, *e.g.*, EP 257,956). If a peptide or small molecule is employed as an antagonist or agonist, it is preferably administered orally in the form of a liquid or solid.

35 PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide can also be administered in the form of sustained-released preparations. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the protein, which matrices are in the form of shaped articles, *e.g.*, films, or microcapsules. Microencapsulation of recombinant proteins for sustained release has been successfully performed with human growth hormone (rhGH), interferon- (rhIFN-),

interleukin-2, and MN rgp120. Johnson et al., Nat. Med., 2:795-799 (1996); Yasuda, Biomed. Ther., 27:1221-1223 (1993); Hora et al., Bio/Technology, 8:755-758 (1990); Cleland, "Design and Production of Single Immunization Vaccines Using Polylactide Polyglycolide Microsphere Systems," in Vaccine Design: The Subunit and Adjuvant Approach, Powell and Newman, eds, (Plenum Press: New York, 1995), pp. 439-462; WO 97/03692, WO 96/40072, WO 96/07399; and U.S. Pat. No. 5,654,010. Examples of sustained-release matrices include polyesters, hydrogels (e.g., poly(2-hydroxyethyl-methacrylate) as described by Langer *et al.*, J. Biomed. Mater. Res., 15: 167-277 (1981) and Langer, Chem. Tech., 12: 98-105 (1982) or poly(vinylalcohol)), polylactides (U.S. Patent No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman *et al.*, Biopolymers, 22: 547-556 (1983)), non-degradable ethylene-vinyl acetate (Langer *et al.*, *supra*), degradable lactic acid-glycolic acid copolymers such as the Lupron Depot™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid (EP 133,988). The sustained-release formulations developed using poly-lactic-co-glycolic acid (PLGA) polymer are preferred due to its biocompatibility and wide range of biodegradable properties. The degradation products of PLGA, lactic and glycolic acids, can be cleared quickly within the human body. Moreover, the degradability of this polymer can be adjusted from months to years depending on its molecular weight and composition. Lewis, "Controlled release of bioactive agents from lactide/glycolide polymer," in: M. Chasin and R. Langer (Eds.), Biodegradable Polymers as Drug Delivery Systems (Marcel Dekker: New York, 1990), pp. 1-41.

While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated proteins remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for protein stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

Sustained-release PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide compositions also include liposomally entrapped PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides. Liposomes containing the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide are prepared by methods known *per se*: DE 3,218,121; Epstein *et al.*, Proc. Natl. Acad. Sci. USA, 82: 3688-3692 (1985); Hwang *et al.*, Proc. Natl. Acad. Sci. USA, 77: 4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese patent application 83-118008; U.S. Patent Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. % cholesterol, the selected proportion being adjusted for the optimal therapy.

The therapeutically effective dose of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, agonist, or antagonist thereto will, of course, vary depending on such factors as the pathological condition to be treated (including prevention), the method of administration, the type of compound

being used for treatment, any co-therapy involved, the patient's age, weight, general medical condition, medical history, etc., and its determination is well within the skill of a practicing physician. Accordingly, it will be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the maximal therapeutic effect. The progress of this therapy is easily monitored by appropriate clinical diagnostic methods. Animal experiments provide reliable guidance for the determination of effective doses for human therapy. Interspecies scaling of effective doses can be performed following the principles laid down by Mordenti, J. and Chappell, W. "The use of interspecies scaling in toxicokinetics" In Toxicokinetics and New Drug Development, Yacobi et al., Eds., Pergamon Press, New York 1989, pp. 42-96.

When *in vivo* administration of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, agonist, or antagonist thereof is employed, normal dosage amounts can vary from about 10 ng/kg to up to 100 mg/kg of mammal body weight or more per day, preferably about 1 µg/kg/day to 10 mg/kg/day, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature; see, for example, U.S. Pat. Nos. 4,657,760; 5,206,344; or 5,225,212. It is anticipated that different formulations will be effective for different treatment compounds and different disorders, that administration targeting one organ or tissue, for example, can necessitate delivery in a manner different from that to another organ or tissue.

Another formulation comprises incorporating a compound of the invention into formed articles. Such articles can be used in modulating endothelial cell growth, angiogenesis, and tumor invasion and metastasis. The therapeutic method includes administering the composition topically, systemically, or locally as an implant or device. When administered, the therapeutic composition for use is in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably be encapsulated or injected in a viscous form for delivery to the targeted site. Topical administration may be suitable for wound healing, tissue repair, and skin and oral lesions and cancers. A kit, contains at least an article of manufacture comprising a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or an agonist or an antagonist thereof useful for the diagnosis or treatment of the disorders described above, and a label. The article of manufacture can be a container, including, for example, bottles, vials, syringes, test tubes, implants, and pumps. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition that is effective for diagnosing or treating the condition and may have a sterile access port (for example, the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The active agent in the composition is a compound of the invention. The label on, or associated with, the container indicates that the composition is used for diagnosing or treating the condition of choice. The kit can further comprise a second container comprising a pharmaceutically-acceptable buffer, such as phosphate-buffered saline, Ringer's solution, and dextrose solution. It can further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for use. The kit can also comprise a second or third container with another active agent as described herein.

N. Combination Therapies

The effectiveness of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-

MG.72 polypeptide or an agonist or antagonist thereof in preventing or treating the disorder in question may be improved by administering the active agent serially or in combination with another agent that is effective for those purposes, either in the same composition or as separate compositions. Positive regulators of angiogenesis include aFGF, bFGF, TGF- α , TGF- β , HGF, TNF- α , angiogenin, IL-8, etc. (see for example, Folkman *et al.*, *J. Biol. Chem.*, 267: 10931-10934 (1992) and Klagsbrun *et al.*, *Annu. Rev. Physiol.*, 53: 217-239 (1991)) and VEGF (Ferrara *et al.*, *Endocr. Rev.*, 18: 4-25 (1997)). Negative regulators include thrombospondin (Good *et al.*, *Proc. Natl. Acad. Sci. USA.*, 87: 6624-6628 (1990)), the 16-kilodalton N-terminal fragment of prolactin (Clapp *et al.*, *Endocrinology*, 133: 1292-1299 (1993)), angiostatin (O'Reilly *et al. Cell*, 79: 315-328 (1994)), and endostatin (O'Reilly *et al.*, *Cell*, 88: 277-285 (1996)). The effectiveness of the agonist in preventing or treating disease may be improved by administering the agonist serially or in combination with yet another agent that is effective for those purposes, such as immunoadhesins, ribozymes, antisense agents, tumor necrosis factor (TNF), an antibody capable of inhibiting or neutralizing the angiogenic activity of acidic or basic fibroblast growth factor (FGF), vascular endothelial growth factor (VEGF), or hepatocyte growth factor (HGF), an antibody capable of inhibiting or neutralizing the coagulant activities of tissue factor, protein C, or protein S (see Esmon, *et al.*, PCT Patent Publication No. WO 91/01753, published 21 February 1991), an antibody capable of binding to HER2 receptor (see Hudziak, *et al.*, PCT Patent Publication No. WO 89/06692, published 27 July 1989), or one or more conventional therapeutic agents such as, for example, alkylating agents, folic acid antagonists, anti-metabolites of nucleic acid metabolism, antibiotics, pyrimidine analogs, 5-fluorouracil, cisplatin, purine nucleosides, amines, amino acids, triazol nucleosides, corticosteroids and proteins such as angiostatin, endostatin, thrombospondin, and platelet factor 4. For example, vascularization of tumors can be blocked with combination therapy, in which one or more antagonists are administered to tumor-bearing patients at therapeutically effective doses as determined for example by observing necrosis of the tumor or its metastatic foci, if any. This therapy is continued until such time as no further beneficial effect is observed or clinical examination shows no trace of the tumor or any metastatic foci. The antagonist is administered, alone or in combination, with an auxiliary agent such as α -, β -, or γ -interferon, anti-HER2 antibody, heregulin, anti-heregulin antibody, D-factor, interleukin-1 (IL-1), interleukin-2 (IL-2), granulocyte-macrophage colony stimulating factor (GM-CSF), or agents that promote microvascular coagulation in tumors, such as anti-protein C antibody, anti-protein S antibody, or C4b binding protein (Esmon, *et al.*, PCT Patent Publication No. WO 91/01753). Such other agents may be present in the composition being administered or may be administered separately. Also, the antagonist is suitably administered serially or in combination with heat or radiological treatments, whether involving irradiation or administration of radioactive substances. Examples of chemotherapeutic agents include, but are not limited to, anticancer drugs such as daunorubicin, dactinomycin, doxorubicin, bleomycin, mitomycin, nitrogen mustard, chlorambucil, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine (CA), 5-fluorouracil (5-FU), floxuridine (5-FuDR), methotrexate (MTX), colchicine, vincristine, vinblastine, etoposide, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, The Merck Manual of Diagnosis and Therapy, 15th Ed., Berkow *et al.*, eds., 1987, Rahway, N.J., pages 1206-1228).

Since the auxiliary agents will vary in their effectiveness it is desirable to compare their impact on the

tumor by matrix screening in conventional fashion. For example, the administration of an antagonist of the invention and auxiliary agent is repeated until the desired clinical effect is achieved. In instances where solid tumors are found in the limbs or in other locations susceptible to isolation from the general circulation, the therapeutic agents described herein are administered to the isolated tumor or organ. In a preferred embodiment, a FGF, platelet-derived growth factor (PDGF), or VEGF antagonist, such as an anti-FGF, anti-VEGF, or an anti-PDGF neutralizing antibody, is administered to the patient in conjunction with an antagonist compound of the invention. Treatment with an antagonist can be suspended during periods of wound healing or desirable vascularization, or alternatively an agonist of the invention can be used to promote such benefit.

For other indications, PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides or their agonists can be combined with other agents beneficial to the treatment of the defect, wound, or tissue in question. These agents include various growth factors such as EGF, PDGF, TGF- α or TGF- β , IGF, FGF, and CTGF, as discussed herein.

In addition, PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides or their antagonists used to treat cancer may be combined with cytotoxic, chemotherapeutic, or growth-inhibitory agents as identified above. Also, for cancer treatment, the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or antagonist thereof is suitably administered serially or in combination with radiological treatments, whether involving irradiation or administration of radioactive substances.

Certain preferred embodiments of the invention provide pharmaceutical compositions containing (a) one or more antigene compounds and (b) one or more other agents which function by a non-antigene mechanism, such as chemotherapeutic, angiogenic, or angiostatic agents as discussed. Two or more combined compounds may be used together or sequentially.

In another related embodiment, compositions of the invention may contain one or more antigene compounds, particularly oligonucleotides, targeted to a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 nucleic acid and one or more additional antigene compounds targeted to a second nucleic acid target. Numerous examples of antigene compounds are known in the art. Two or more combined compounds may be used together or sequentially. For example, Im et al., Cancer Research 59(4):895-900 (1999) reported inhibition of tumor growth using antisense VEGF. The recombinant adenoviral vector Ad5CMV carried the coding sequence of wild-type VEGF165 cDNA in an antisense orientation. Infection of U-87 MG malignant glioma cells with the vector resulted in reduction of the level of the endogenous VEGF mRNA and drastically decreased the production of the targeted secretory form of the VEGF protein. Treatment of s.c. human glioma tumors established in nude mice with intralesional injection of antisense VEGF vector inhibited tumor growth. Sharma et al. (J. Clin. Invest. 102(8):1599-608 (1998)) reported that by blocking perlecan expression by using either constitutive CMV-driven or doxycycline-inducible antisense constructs, growth of colon carcinoma cells was markedly attenuated. In both tumor xenografts induced by human colon carcinoma cells and tumor allografts induced by highly invasive mouse melanoma cells, perlecan suppression caused substantial inhibition of tumor growth and neovascularization. It is noted that the mouse system, in combination with an appropriate human tumor or tumor cell line xenograft or mouse allograft, also provides a

rapid means to screen for maximally effective PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene compounds *in vivo*.

The effective amounts of the therapeutic agents administered in combination with the compounds of the invention will be at the physician's or veterinarian's discretion. Dosage administration and adjustment is done to achieve maximal management of the conditions to be treated. The dose will additionally depend on such factors as the type of the therapeutic agent to be used and the specific patient being treated. Typically, the amount employed, at least as a starting point, will be the same dose as that used, if the given therapeutic agent is administered without a compound of the invention.

According to the present invention, angiogenesis, vascular and neovascular conditions are particularly well-suited to antigene therapy (see e.g., Thomas et al. Radiographics 18(6):1373-94 (1998)). Accordingly, local gene transfer into the vascular wall offers a promising alternative to treat angiogenic-related diseases and disorders described herein. Blood vessels, and the vascular wall, are among the easiest targets for gene therapy because of ready accessibility and of novel percutaneous, catheter-based treatment methods. Vascular and interventional radiology techniques also are ideally suited for minimally invasive, readily monitored gene delivery, of either viral or nonviral vectors or of syntehtetic oligo compounds. Recombinant genes can be delivered *ex vivo* and *in vivo*; the latter approaches can involve well-known open surgical, percutaneous injection, or endovascular catheter-based methods. Perforated, hydrogel-coated, and double balloon catheters can also be readily used. Catheter systems for gene transfer enable delivery of the vector to the precise anatomic location with transfection limited to the cells of interest and will minimize shedding of the vector to distal sites, systemic effects of the therapeutic agent, and morbidity from the delivery method. On the other hand, gene transfer to the artery wall can also be accomplished from adventitia, and in some situations intramuscular gene delivery. Promising therapeutic effects have been obtained in animal models of restenosis with the transfer of genes for vascular endothelial growth factor, fibroblast growth factor, thymidine kinase, p53, bcl-x, nitric oxide synthase and retinoblastoma. Also, growth arrest homeobox gene and antisense oligonucleotides against transcription factors or cell cycle regulatory proteins have produced beneficial therapeutic effects. Antiangiogenic tumor therapies using antigen technology can provide broad-spectrum action, low toxicity, and, in the case of direct endothelial targeting, an absence of drug resistance. Gene therapy offers a potential way to achieve sustained therapeutic release of potent antiangiogenic substances. An alternative for longer term administration (or as combination therapy) are recombinant vectors carrying antisense genes. For example, adeno-associated virus (rAAV) vectors carrying genes coding for angiostatin, endostatin, and an antisense mRNA species against vascular endothelial growth factor (VEGF), efficiently transduced three human tumor cell lines tested. Transduction with an rAAV-encoding antisense VEGF mRNA inhibited the production of endogenous tumor cell VEGF (Nguyen et al., Cancer Research, 58(24):5673-7 (1998)).

The following examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

All patent and literature references cited in the present specification are hereby incorporated by reference in their entirety.

EXAMPLES

Commercially available reagents referred to in the examples were used according to manufacturer's instructions unless otherwise indicated. The source of those cells identified in the following examples, and throughout the specification, by ATCC accession numbers is the American Type Culture Collection, Manassas, VA.

EXAMPLE 1

Isolation of cDNA Clones Encoding a Human PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72

Formation of Three-Dimensional Collagen Gels: Tube formation by endothelial cells is a critical process in the development of a blood vessel during angiogenesis and vasculogenesis. Human umbilical cord endothelial cells (HUVEC) undergoing tube formation in collagen gels in the presence of growth factors, mimic the angiogenic environment of endothelial cells in vivo, providing a well-accepted system for angiogenesis and vasculogenesis, both in normal and neoplastic conditions. The three dimensional gel is prerequisite for the differentiation and fusion of endothelial cells into tubes, as HUVECs grown on the surface of gelatin or on plastic do not undergo tube-formation.

In brief, HUVECs were grown under various conditions, inductive or non-inductive to tube formation, either on collagen film (non-inductive) or in collagen gels (inductive), with or without the addition of growth factors to simulate induction by normal angiogenic factors or by tumor-derived factors. Differential cDNA screening was used to identify genes critical to this process. The particular method used to quantitate endothelial cell gene expression was Quantitative Expression Analysis (QEA; U.S. Patent 5,871,697). HUVEC total RNA was prepared, followed by mRNA purification and double stranded cDNA synthesis. The cDNA was digested with restriction enzyme pairs to produce cDNA fragments, which were then ligated with linkers. Primer pairs bearing the specific sequences of the linkers were used to amplify the restricted products in a PCR reaction. Quantification and identification of amplified products revealed modulated genes, thus identifying genes critical to angiogenesis.

Tube formation was achieved as follows. Collagen gels were formed by mixing together an ice-cold gelation solution of a 100:27.7:50:10:750:62.5 ratio (by volume) of a ten-fold-concentrated M199 stock, water, 0.53 M NaHCO₃, 200 mM L-glutamine, type I collagen, and 0.1 M NaOH. This was mixed with HUVEC cells (in 1X basal medium at a concentration of 3×10^6 cells/ml) at a ratio of 4 volumes gelation solution to 1 volume of cells. The gels were allowed to form by incubation in a CO₂-free incubator at 37°C for 30 min to 1 hour. The gels were then overlaid with 1X basal medium consisting of M199 supplemented with 1% FBS, 1X ITS, 2mM L-glutamine, 50 mg/ml ascorbic acid, 26.5 mM NaHCO₃, 100 U/ml penicillin and 100 U/ml streptomycin. In the tube-forming experiments, the culture media was supplemented with 80 nM PMA, 40 ng/ml bFGF and 40 ng/ml VEGF. In a parallel set of experiments, endothelial cells were cultured on the surface of type I collagen, or on pig skin gelatin (DIFCO, USA) in 1X basal medium consisting of M199 supplemented with 1% FBS, 1X ITS, 2mM L-glutamine, 50 mg/ml ascorbic acid, 26.5 mM NaHCO₃, 100 U/ml penicillin and 100 U/ml streptomycin, without or with 80 nM PMA, 40 ng/ml bFGF and 40 ng/ml VEGF.

For the differential cDNA screening experiment (by QEA, also referred to as GeneCalling™, Curagen Corp., USA), mRNA was isolated from cells incubated in the above conditions for 4 hr, 24 hr and 48 hr.

mRNA was isolated and cDNA synthesized as follows. Media was aspirated from the surface of the collagen gels and the gels were scraped into a 50 ml polypropylene tube containing 3 volumes of Tri-Reagent-LS (Molecular Research Center, Cincinnati, OH). The tubes were incubated 10 min at 23°C with intermittent gentle agitation. The tubes were stored at -80°C until all experimental samples had been collected. The tubes were then thawed at room temperature and the mRNA extracted following manufacturer's specifications. The RNA pellets were resuspended in DEPC-treated water and RNA content quantified spectroscopically at 260 nm. RNA samples were stored -20°C. Samples used for GeneCalling™ analysis were shipped on dry ice to Curagen Corp. (New Haven, CT). Samples from time points of 4, 24 and 48 hrs were used for the GeneCalling™ analysis, and in a separate experiments, samples cells grown in collagen gels and on the surface of type I collagen in 1X basal medium supplemented with 80 nM PMA, 40 ng/ml bFGF and 40 ng/ml VEGF from time points of 30 min, and 2, 4, 8, 16, 24, 38 and 46.5 hrs were prepared for Taqman PCR confirmation. For the quantitative expression analysis, contaminating DNA was removed by treatment of the isolated RNA with DNase I (Promega, Madison, WI). Poly-A+ RNA was prepared by fractionation of total RNA using an mRNA purification kit that utilized biotinylated oligo-dT-Streptavidin magnetic bead method (MPG, Lincoln Park, NJ) followed by cDNA synthesis by reverse transcription of oligo-dT primed mRNA (Superscript II, Life Technologies) and second strand synthesis. Terminal phosphate removal was achieved by treatment with Artic Shrimp Alkaline Phosphatase (Amersham Life Sciences, Piscataway, NJ) followed by purification of cDNA by phenol-chloroform extraction. Yield of cDNA was quantitated by fluorometry using PicoGreen dye (Molecular Probes, Eugene OR). Double stranded DNA was digested using pairs of restriction enzymes with 6 base-pair recognition sites. More than 48 enzyme pairs were used and were chosen such that a representative coverage of most of the possible sequences in a given DNA sample was achieved. PCR amplification using specific linkers was carried out as described in US Patent 5,871,697. The final DNA products were denatured by heating to 96°C and electrophoresed on ultra-thin polyacrylamide gels under denaturing conditions in 6M urea. PCR products were visualized by the presence of FAM label on the product using a multi-color laser excitation Niagara (Curagen Corp., New Haven CT) imaging system.

GeneCalling analysis used a fully integrated Web-based interactive bioinformatics data gathering and analysis suite called "GeneScape." The data obtained from Niagara gels were GeneCalled against public and proprietary databases using present statistical and mathematical criteria (US Patent 5,871,697) and a gene list was generated from the cDNA fragment data that is a list of likely genes that the cDNA fragment can belong to based on the size of the fragment and the position of the restriction enzyme pair that produced it in the known sequence. If a gene candidate could not be obtained, the cDNA fragment was designated as belonging to a putative novel gene.

A GeneCall was defined as the probability of a cDNA fragment belonging to a known gene. GeneCalls were confirmed in a poisoning reaction where the known sequence of the likely gene of interest is used to design poisoning primers as previously described (US Patent 5,871,697; Shimkets *et al. Nature Biotechnology* 17(8):798-803 (1999)). Ablation of the cDNA fragment of interest confirmed that the cDNA fragment

belonged to the gene for which the primer was designed.

If no GeneCall was obtained for a cDNA fragment, the putative novel cDNA fragment was eluted from, and subcloned into *E. coli* using standard TA-cloning vector (Invitrogen, Palo Alto, CA). The putative novel cDNA fragment was then sequenced and the resulting sequence used to design primer primers for confirmation as described above.

To confirm the expression data from GeneCalling by an independent technique, Quantitative RT-PCR (Taqman), in which gene specific PCR oligonucleotide primer pairs and oligonucleotide probes labeled with a reporter fluorescent dye at the 5' end and quencher fluorescent dye at the 3' end were designed using the Oligo 4.0 software (National Bioscience, Plymouth MN). Total RNA (50 ng) was added to a 50 µl RT-PCR reaction mixture according to the manufacturer's protocol (Roche Molecular Systems Inc., Branchburg, NJ). The thermal cycling conditions included 1 cycle at 48°C for 30 min, 1 cycle at 95°C for 10 min, 40 cycles at 95°C for 15s, annealing at 60°C for 1 min, and a final hold at 25°C for 2 min. Standard curves for the expression of each gene were generated by serial dilution of a standard preparation of total RNA isolated from quiescent HUVEC grown in monolayer culture. Data were expressed as the fold induction normalized to the same gene from quiescent HUVEC RNA.

The GeneCalling process was used in the selection (gating) of differentially expressed genes in tube formation. The experimental design was based on the observation that endothelial cells grown on the surface of type I collagen in 1X basal medium supplemented with 80 nM PMA, 40 ng/ml bFGF and 40 ng/ml VEGF do not form tubes, but rather remain as a monolayer. This result also occurs if the cells are grown on gelatin, a form of denatured collagen. However, if the cells are suspended in a three dimensional collagen gel, and grown in 1X basal media supplemented with 80 nM PMA, 40 ng/ml bFGF and 40 ng/ml VEGF, the cells undergo a synchronous differentiation into an interconnected tube like network. The tubular structures contain lumen-like structures. At 4 hours, large intracellular vacuoles are forming, but the cells are still round. At 24 hrs, the cells have become elongated and many cells are touching other cells. By 48 hrs, the cells have become interconnected and share common lumens. To select for genes that play a role in this differentiation, an array of GeneCalling differences was set up such that cDNA fragments that changed more than 2 fold between 24 hours and 4 hours, between 48 hours and 4 hours, or between 48 hours and 24 hrs, in the 3-D gel environment, but which were unchanged or changed less than 2 fold in the 2D (surface of type I collagen or gelatin) environment at the same pair-wise time comparisons were preferentially selected and identified. In addition, those cDNA fragments which demonstrate large (greater than 8 fold) changes in gene expression were also identified.

Full length cDNAs corresponding to the differentially expressed genes identified by their GeneCalled fragments were prepared and sequenced as follows. An oligo dT primed cDNA library was prepared from mRNA isolated from human HUVEC cells as above, using reagents and protocols from Invitrogen, San Diego, CA (Fast Track 2). This RNA was used to generate an oligo dT primed cDNA library in the vector pRK5D using reagents and protocols from Life Technologies, Gaithersburg, MD (Super Script Plasmid System). In this procedure, the double stranded cDNA was sized to greater than 1000 bp and the SalI/NotI linked cDNA was cloned into XhoI/NotI cleaved vector. pRK5D is a cloning vector that has an sp6 transcription initiation site followed by an SfiI restriction enzyme site preceding the XhoI/NotI cDNA cloning sites.

Oligonucleotides probes based upon the above described GeneCall fragment sequence were then synthesized to identify by PCR a cDNA library that contained the sequence of interest, and to use as probes to isolate a clone of the full-length coding sequence for the differentially expressed gene of interest. Forward and reverse PCR primers generally range from 20 to 30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In order to screen several libraries for a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley and Sons (1997), with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs.

PRO-C-MG.2. The initial GeneCall assembled fragment sequence was SEQ ID No. 5. To obtain the full-length clone, oligonucleotide probes based on this sequence were as follows:

forward PCR primer 5' GGACGACACGGTGCCGCTGACAGC 3' (SEQ ID NO:6)

reverse PCR primer 5' GTTTTCCAGAGAAATTCCTCTTTGCACTCGA 3' (SEQ ID NO:8)

hybridization probe 5' GCCATCGAGGCGAGCCAGAGCCTGCAGTCCCACACGGAATATATTATTCGA 3' (SEQ ID NO:7)

A full length clone was identified that contained a single open reading frame with an apparent translational initiation site at nucleotide positions 66-68 and a stop signal at nucleotide positions 1794-1796 (SEQ ID NO:1). The predicted polypeptide is 577 amino acids long, has a calculated molecular weight of approximately 64935.06 daltons and an estimated pI of approximately 9.94. Analysis of the full-length PRO-C-MG.2 sequence shown in SEQ ID NO:2 evidences the presence of a variety of important polypeptide domains. The locations given for those important polypeptide domains are approximate as described: cAMP- and cGMP-dependent protein kinase phosphorylation site at amino acid positions 54-58, 441-445, and 464-468; casein kinase II phosphorylation site at amino acid positions 32-36, 57-61, 110-114, 179-183, 190-194, 216-220, 233-237, 402-406, 452-456, 470-474; tyrosine kinase phosphorylation site at amino acid positions 116-125 and 117-125; N-myristoylation site at amino acid positions 489-495, 545-551, and 549-555; leucine zipper pattern at amino acid positions 289-311; a PX kinase domain at amino acid position 16-122; and, a pkinase domain from amino acid positions 230-284. This gene had a greater than 4-fold increase in gene expression as determined by the GeneCalling approach. clone 12 is +2.5X by GeneCalling

Clone DNA-C-MG.2-1776 has been deposited with ATCC on September 28, 1999, and is assigned ATCC deposit no. PTA-799.

PRO-C-MG.12. The initial GeneCall assembled fragment sequence was SEQ ID No. 9. To obtain the full-length clone, oligonucleotide probes based on this sequence were as follows:

forward PCR primer 5' GACCTATTGGGACACCTTCTGGAG 3' (SEQ ID NO:10)

reverse PCR primer 5' CTTGGTCAGACGAGAGGAGCTGATC 3' (SEQ ID NO:12)

hybridization probe 5' CCCGCCTAGTGCCAAGCAACCCTCCAAGATGCTAGTTATCAAA 3' (SEQ ID NO:11)

A full length clone was identified that contained a single open reading frame with an apparent translational initiation site at nucleotide positions 465-467 and a stop signal at nucleotide positions 1884-1886 (SEQ ID

NO:3). The predicted polypeptide is 474 amino acids long, has a calculated molecular weight of approximately 52573.30 daltons and an estimated pI of approximately 6.66. Analysis of the full-length PRO-C-MG.12 sequence shown in SEQ ID NO:4 evidences the presence of a variety of important polypeptide domains. The locations given for those important polypeptide domains are approximate as described: cAMP- and cGMP-dependent protein kinase phosphorylation site at amino acid positions 199-203 and 316-320; casein kinase II phosphorylation site at amino acid positions 61-65, 81-85, 202-206, 266-270, 292-296, 328-332, 353-357, 411-415, 458-462, 463-467, 467-471, and 468-472; N-myristoylation site at amino acid positions 112-118, 122-128, 177-183, 218-224, 224-230, 262-268, 287-293, and 364-370; and a potential autocatalytic peptide splicing site at position 99-104 (LPRGhD; see Gu et al., J. Biol. Chem. 268(10):7372-81 (1993)). This gene had a greater than 2.5-fold increase in gene expression as determined by the GeneCalling approach.

Clone DNA-C-MG.12-1776 has been deposited with ATCC on September 28, 1999, and is assigned ATCC deposit no. PTA-798.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in SEQ ID NO:4, evidenced sequence identity between the PRO-C-MG.12 amino acid sequence and the following Dayhoff sequences: Accession P_Y00281. P_Y00281 was reported as a 337 amino acid long human allegedly secreted protein encoded by gene 24 in WO9906423. This 337 amino acid sequence matches PRO-C-MG.12 from amino acid position 138 to 474. WO9906423 alleges that P_Y00281 gene maps to chromosome 1, and was expressed primarily in brain, and to a lesser extent in ovaries and activated T-cells, and consequently was alleged to find use primarily in brain neurodegenerative disorders, immune deficiencies and reproduction.

PRO-C-MG.45. The initial GeneCall assembled fragment sequence was SEQ ID No. 24. This was similar to EST accession # AA461480. To obtain a longer sequence, SEQ ID NO: 24 was used to search public EST and other sequence databases (e.g., GenBank). The search was performed using the computer program BLAST or BLAST-2 (Altschul *et al.*, Methods in Enzymology, 266:460-480 (1996)). Comparisons with a BLAST score of preferably 70 (or in some cases, 90) or greater that did not encode known proteins were clustered and assembled into a consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, Washington). Using this homology screen, consensus DNA sequences were assembled relative to other identified EST sequences using phrap. In addition, the consensus DNA sequences obtained were often (but not always) extended using repeated cycles of BLAST and phrap to extend the consensus sequence as far as possible using the sources of EST sequences discussed above.

From either the GeneCalled sequence or the consensus sequence or both, oligonucleotides are then synthesized and used to identify by PCR a cDNA library that contains the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for a PRO polypeptide. Forward and reverse PCR primers generally range from 20 to 30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In some cases, additional oligonucleotides are synthesized when the consensus sequence is greater than about 1-1.5 kbp. In order to screen several libraries for a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel *et al.*, Current Protocols in Molecular Biology, *supra*, with the PCR primer pair. A positive library

was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs. To obtain the full-length clone, preferred oligonucleotide primers are preferably as follows:

forward PCR primer 5' atgcagttct tttcaactt cc (SEQ ID NO:26)

reverse PCR primer 5' ctagagacca atctaagtaa 3' (SEQ ID NO:27)

- 5 The probe can be a unique region preferably from about 19 to about 100 base pairs.

The cDNA libraries used to isolate the cDNA clones are constructed by standard methods using commercially available reagents, such as those from Invitrogen, San Diego, CA. The cDNA is primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

- 15 **PRO-C-MG.64.** The initial GeneCall assembled fragment sequence was SEQ ID No. 5. This was similar to EST accession # AA913939. To obtain a longer sequence, SEQ ID NO: 24 was used to search public EST and other sequence databases (*e.g.*, GenBank). The search was performed using the computer program BLAST or BLAST-2 (Altschul *et al.*, Methods in Enzymology, 266:460-480 (1996)). Comparisons with a BLAST score of preferably 70 (or in some cases, 90) or greater that did not encode known proteins were clustered and assembled into a consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, Washington). Using this homology screen, consensus DNA sequences were assembled relative to other identified EST sequences using phrap. In addition, the consensus DNA sequences obtained were often (but not always) extended using repeated cycles of BLAST and phrap to extend the consensus sequence as far as possible using the sources of EST sequences discussed above. To obtain the full-length clone, preferred oligonucleotide primers are as follows:

forward PCR primer 5' atggtggagt ggaggacctg (SEQ ID NO:28)

reverse PCR primer 5' ctccaacacc aagtactctt ga 3' (SEQ ID NO:29)

- 25 The probe can be a unique region preferably from about 19 to about 100 base pairs.

The cDNA libraries used to isolate the cDNA clones are constructed by standard methods using commercially available reagents, such as those from Invitrogen, San Diego, CA. The cDNA is primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

- 30 **PRO-C-MG.72.** The initial GeneCall assembled fragment sequence was SEQ ID No. 5. This was similar to EST accession # aa771960. To obtain the full-length clone, oligonucleotide probes based on this sequence were as follows:

forward PCR primer 5' gctgcttcttggtggaagattctgg 3' (SEQ ID NO:20)

reverse PCR primer 5' ccagaatctccaaccaagaagcagc 3' (SEQ ID NO:21)

hybridization probe 5' gcatcatgctgtttgacactttccaattaaaagtccttcataaaactt tgc 3' (SEQ ID NO:22),

and a reverse hybridization probe was ggagctgccattagaatcaagaatctttgc (SEQ ID NO:23)

A full length clone was identified that contained a single open reading frame with an apparent translational initiation site at nucleotide positions 71-73 and a stop signal at nucleotide positions 2060-2062 (SEQ ID NO:13). The predicted polypeptide is 663 amino acids long. Analysis of the full-length PRO-C-MG.72 sequence shown in SEQ ID NO:14 evidences the presence of a variety of important polypeptide domains. The most interesting of which is the RhoGAP domain approximate from about amino acid 343 to about 494, as determined by the Pfam algorithm, giving a very significant E-value of 8.2×10^{-28} and a score of 105.8. Accordingly, PRO-C-MG.72 is believed to have activity as a GTPase-activating protein, preferably of the Rho-type. In this regard a particularly important regions, that contain GTPase-activating active domains, are from about amino acid position 206 to about 553, to about 307 to about 500, and to about 341 to about 533, in SEQ ID NO:14.

EXAMPLE 2

Use of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 as a hybridization probe

The following method describes use of a nucleotide sequence encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 as a hybridization probe.

DNA comprising the coding sequence of full-length or mature PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 is employed as a probe to screen for homologous DNAs (such as those encoding naturally-occurring variants of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72) in human tissue cDNA libraries or human tissue genomic libraries.

Hybridization and washing of filters containing either library DNAs is performed under the following high stringency conditions. Hybridization of radiolabeled PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-derived probe to the filters is performed in a solution of 50% formamide, 5x SSC, 0.1% SDS, 0.1% sodium pyrophosphate, 50 mM sodium phosphate, pH 6.8, 2x Denhardt's solution, and 10% dextran sulfate at 42°C for 20 hours. Washing of the filters is performed in an aqueous solution of 0.1x SSC and 0.1% SDS at 42°C.

DNAs having a desired sequence identity with the DNA encoding full-length native sequence PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can then be identified using standard techniques known in the art.

EXAMPLE 3

Expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 in *E. coli*

This example illustrates preparation of an unglycosylated form of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 by recombinant expression in *E. coli*.

The DNA sequence encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 is initially amplified using selected PCR primers. The primers should contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector. A variety of expression vectors can be employed. An example of a suitable vector is pBR322 (derived from *E. coli*; see Bolivar et al., *Gene*, 2:95 (1977)) which contains genes for ampicillin and tetracycline resistance. The vector is digested with restriction enzyme and dephosphorylated. The PCR amplified sequences are then ligated into the vector. The

vector will preferably include sequences which encode for an antibiotic resistance gene, a *trp* promoter, a polyhis leader (including the first six STII codons, polyhis sequence, and enterokinase cleavage site), the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 coding region, lambda transcriptional terminator, and an *argU* gene.

The ligation mixture is then used to transform a selected *E. coli* strain using the methods described in Sambrook et al., *supra*. Transformants are identified by their ability to grow on LB plates and antibiotic resistant colonies are then selected. Plasmid DNA can be isolated and confirmed by restriction analysis and DNA sequencing.

Selected clones can be grown overnight in liquid culture medium such as LB broth supplemented with antibiotics. The overnight culture can subsequently be used to inoculate a larger scale culture. The cells are then grown to a desired optical density, during which the expression promoter is turned on.

After culturing the cells for several more hours, the cells can be harvested by centrifugation. The cell pellet obtained by the centrifugation can be solubilized using various agents known in the art, and the solubilized PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 protein can then be purified using a metal chelating column under conditions that allow tight binding of the protein.

PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be expressed in *E. coli* in a poly-His tagged form, using the following procedure. The DNA encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 is initially amplified using selected PCR primers. The primers will contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector, and other useful sequences providing for efficient and reliable translation initiation, rapid purification on a metal chelation column, and proteolytic removal with enterokinase. The PCR-amplified, poly-His tagged sequences are then ligated into an expression vector, which is used to transform an *E. coli* host based on strain 52 (W3110 *fuhA(tonA) lon galE rpoHts(htpRts) clpP(lacIq)*). Transformants are first grown in LB containing 50 mg/ml carbenicillin at 30°C with shaking until an O.D.600 of 3-5 is reached. Cultures are then diluted 50-100 fold into CRAP media (prepared by mixing 3.57 g (NH₄)₂ SO₄, 0.71 g sodium citrate•2H₂O, 1.07 g KCl, 5.36 g Difco yeast extract, 5.36 g Sheffield hycase SF in 500 mL water, as well as 110 mM MPOS, pH 7.3, 0.55% (w/v) glucose and 7 mM MgSO₄) and grown for approximately 20-30 hours at 30°C with shaking. Samples are removed to verify expression by SDS-PAGE analysis, and the bulk culture is centrifuged to pellet the cells. Cell pellets are frozen until purification and refolding.

E. coli paste from 0.5 to 1 L fermentations (6-10 g pellets) is resuspended in 10 volumes (w/v) in 7 M guanidine, 20 mM Tris, pH 8 buffer. Solid sodium sulfite and sodium tetrathionate is added to make final concentrations of 0.1M and 0.02 M, respectively, and the solution is stirred overnight at 4°C. This step results in a denatured protein with all cysteine residues blocked by sulfitolization. The solution is centrifuged at 40,000 rpm in a Beckman Ultracentrifuge for 30 min. The supernatant is diluted with 3-5 volumes of metal chelate column buffer (6 M guanidine, 20 mM Tris, pH 7.4) and filtered through 0.22 micron filters to clarify. The clarified extract is loaded onto a 5 ml Qiagen Ni-NTA metal chelate column equilibrated in the metal chelate column buffer. The column is washed with additional buffer containing 50 mM imidazole (Calbiochem, Utrol

grade), pH 7.4. The protein is eluted with buffer containing 250 mM imidazole. Fractions containing the desired protein are pooled and stored at 4°C. Protein concentration is estimated by its absorbance at 280 nm using the calculated extinction coefficient based on its amino acid sequence.

The proteins are refolded by diluting the sample slowly into freshly prepared refolding buffer consisting of: 20 mM Tris, pH 8.6, 0.3 M NaCl, 2.5 M urea, 5 mM cysteine, 20 mM glycine and 1 mM EDTA. Refolding volumes are chosen so that the final protein concentration is between 50 to 100 micrograms/ml. The refolding solution is stirred gently at 4°C for 12-36 hours. The refolding reaction is quenched by the addition of TFA to a final concentration of 0.4% (pH of approximately 3). Before further purification of the protein, the solution is filtered through a 0.22 micron filter and acetonitrile is added to 2-10% final concentration. The refolded protein is chromatographed on a Poros R1/H reversed phase column using a mobile buffer of 0.1% TFA with elution with a gradient of acetonitrile from 10 to 80%. Aliquots of fractions with A280 absorbance are analyzed on SDS polyacrylamide gels and fractions containing homogeneous refolded protein are pooled. Generally, the properly refolded species of most proteins are eluted at the lowest concentrations of acetonitrile since those species are the most compact with their hydrophobic interiors shielded from interaction with the reversed phase resin. Aggregated species are usually eluted at higher acetonitrile concentrations. In addition to resolving misfolded forms of proteins from the desired form, the reversed phase step also removes endotoxin from the samples.

Fractions containing the desired folded PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide are pooled and the acetonitrile removed using a gentle stream of nitrogen directed at the solution. Proteins are formulated into 20 mM Hepes, pH 6.8 with 0.14 M sodium chloride and 4% mannitol by dialysis or by gel filtration using G25 Superfine (Pharmacia) resins equilibrated in the formulation buffer and sterile filtered.

EXAMPLE 4

Expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 in mammalian cells

This example illustrates preparation of a potentially glycosylated form of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 by recombinant expression in mammalian cells.

The vector, pRK5 (see EP 307,247, published March 15, 1989), is employed as the expression vector. Optionally, the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 DNA is ligated into pRK5 with selected restriction enzymes to allow insertion of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 DNA using ligation methods such as described in Sambrook et al., *supra*. The resulting vector is called pRK5-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72.

In one embodiment, the selected host cells can be HUVEC cells as described above, using the vectors and transfection methods described herein for other mammalian cells. Transfected HUVEC cells over-expressing PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 or expressing PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antisense are tested, for example, in the tube formation assay.

In one embodiment, the selected host cells can be 293 cells. Human 293 cells (ATCC CCL 1573) are grown to confluence in tissue culture plates in medium such as DMEM supplemented with fetal calf serum and optionally, nutrient components and/or antibiotics. About 10 µg pRK5-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 DNA is mixed with about 1 µg DNA encoding the VA RNA gene [Thimmapaya et al., *Cell*, 31:543 (1982)] and dissolved in 500 µl of 1 mM Tris-HCl, 0.1 mM EDTA, 0.227 M CaCl₂. To this mixture is added, dropwise, 500 µl of 50 mM HEPES (pH 7.35), 280 mM NaCl, 1.5 mM NaPO₄, and a precipitate is allowed to form for 10 minutes at 25°C. The precipitate is suspended and added to the 293 cells and allowed to settle for about four hours at 37°C. The culture medium is aspirated off and 2 ml of 20% glycerol in PBS is added for 30 seconds. The 293 cells are then washed with serum free medium, fresh medium is added and the cells are incubated for about 5 days.

Approximately 24 hours after the transfections, the culture medium is removed and replaced with culture medium (alone) or culture medium containing 200 µCi/ml ³⁵S-cysteine and 200 µCi/ml ³⁵S-methionine. After a 12 hour incubation, the conditioned medium is collected, concentrated on a spin filter, and loaded onto a 15% SDS gel. The processed gel can be dried and exposed to film for a selected period of time to reveal the presence of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide. The cultures containing transfected cells can undergo further incubation (in serum free medium) and the medium is tested in selected bioassays.

In an alternative technique, PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be introduced into 293 cells transiently using the dextran sulfate method described by Sompanyrac et al., *Proc. Natl. Acad. Sci.*, 12:7575 (1981). 293 cells are grown to maximal density in a spinner flask and 700 µg pRK5-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 DNA is added. The cells are first concentrated from the spinner flask by centrifugation and washed with PBS. The DNA-dextran precipitate is incubated on the cell pellet for four hours. The cells are treated with 20% glycerol for 90 seconds, washed with tissue culture medium, and re-introduced into the spinner flask containing tissue culture medium, 5 µg/ml bovine insulin and 0.1 µg/ml bovine transferrin. After about four days, the conditioned media is centrifuged and filtered to remove cells and debris. The sample containing expressed PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can then be concentrated and purified by any selected method, such as dialysis and/or column chromatography.

In another embodiment, PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be expressed in CHO cells. The pRK5-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be transfected into CHO cells using known reagents such as CaPO₄ or DEAE-dextran. As described above, the cell cultures can be incubated, and the medium replaced with culture medium (alone) or medium containing a radiolabel such as ³⁵S-methionine. After determining the presence of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, the culture medium can be replaced with serum free medium. Preferably, the cultures are incubated for about 6 days, and then the conditioned medium is harvested. The medium containing the expressed PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can then be concentrated and purified by any selected method.

Epitope-tagged PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can also be expressed in host CHO cells. The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be subcloned out of the pRK5 vector. The subclone insert can undergo PCR to fuse in frame with a selected epitope tag such as a poly-his tag into a Baculovirus expression vector. The poly-his tagged PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 insert can then be subcloned into a SV40 driven vector containing a selection marker such as DHFR for selection of stable clones. Finally, the CHO cells can be transfected (as described above) with the SV40 driven vector. Labeling can be performed, as described above, to verify expression. The culture medium containing the expressed poly-His tagged PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can then be concentrated and purified by any selected method, such as by Ni^{2+} -chelate affinity chromatography.

PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can also be expressed in CHO and/or COS cells by a transient expression procedure or in CHO cells by another stable expression procedure.

Stable expression in CHO cells is performed using the following procedure. The proteins are expressed as an IgG construct (immunoadhesin), in which the coding sequences for the soluble forms (e.g. extracellular domains) of the respective proteins are fused to an IgG1 constant region sequence containing the hinge, CH2 and CH2 domains and/or is a poly-His tagged form.

Following PCR amplification, the respective DNAs are subcloned in a CHO expression vector using standard techniques as described in Ausubel et al., Current Protocols of Molecular Biology, Unit 3.16, John Wiley and Sons (1997). CHO expression vectors are constructed to have compatible restriction sites 5' and 3' of the DNA of interest to allow the convenient shuttling of cDNA's. The vector used expression in CHO cells is as described in Lucas et al., Nucl. Acids Res. 24:9 (1774-1779 (1996), and uses the SV40 early promoter/enhancer to drive expression of the cDNA of interest and dihydrofolate reductase (DHFR). DHFR expression permits selection for stable maintenance of the plasmid following transfection.

Twelve micrograms of the desired plasmid DNA is introduced into approximately 10 million CHO cells using commercially available transfection reagents Superfect® (Quiagen), Dospers® or Fugene® (Boehringer Mannheim). The cells are grown as described in Lucas et al., *supra*. Approximately 3×10^7 cells are frozen in an ampule for further growth and production as described herein.

The ampules containing the plasmid DNA are thawed by placement into water bath and mixed by vortexing. The contents are pipetted into a centrifuge tube containing 10 mLs of media and centrifuged at 1000 rpm for 5 minutes. The supernatant is aspirated and the cells are resuspended in 10 mL of selective media (0.2 μm filtered PS20 with 5% 0.2 μm diafiltered fetal bovine serum). The cells are then aliquoted into a 100 mL spinner containing 90 mL of selective media. After 1-2 days, the cells are transferred into a 250 mL spinner filled with 150 mL selective growth medium and incubated at 37°C. After another 2-3 days, 250 mL, 500 mL and 2000 mL spinners are seeded with 3×10^5 cells/mL. The cell media is exchanged with fresh media by centrifugation and resuspension in production medium. Although any suitable CHO media can be employed, a production medium described in U.S. Patent No. 5,122,469, issued June 16, 1992 can actually be used. A 3L production spinner is seeded at 1.2×10^6 cells/mL. On day 0, the cell number pH is determined. On day 1, the

spinner is sampled and sparging with filtered air is commenced. On day 2, the spinner is sampled, the temperature shifted to 33°C, and 30 mL of 500 g/L glucose and 0.6 mL of 10% antifoam (e.g., 35% polydimethylsiloxane emulsion, Dow Corning 365 Medical Grade Emulsion) taken. Throughout the production, the pH is adjusted as necessary to keep it at around 7.2. After 10 days, or until the viability dropped below 70%, the cell culture is harvested by centrifugation and filtering through a 0.22 μ m filter. The filtrate was either stored at 4°C or immediately loaded onto columns for purification.

For the poly-His tagged constructs, the proteins are purified using a Ni-NTA column (Qiagen). Before purification, imidazole is added to the conditioned media to a concentration of 5 mM. The conditioned media is pumped onto a 6 ml Ni-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 4°C. After loading, the column is washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein is subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80°C.

Immunoadhesin (Fc-containing) constructs are purified from the conditioned media as follows. The conditioned medium is pumped onto a 5 ml Protein A column (Pharmacia) which had been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column is washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein is immediately neutralized by collecting 1 ml fractions into tubes containing 275 μ L of 1 M Tris buffer, pH 9. The highly purified protein is subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity is assessed by SDS polyacrylamide gels and by N-terminal amino acid sequencing by Edman degradation.

EXAMPLE 5

Expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 in Yeast

The following method describes recombinant expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 in yeast.

First, yeast expression vectors are constructed for intracellular production or secretion of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 from the ADH2/GAPDH promoter. DNA encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 and the promoter is inserted into suitable restriction enzyme sites in the selected plasmid to direct intracellular expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. For secretion, DNA encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be cloned into the selected plasmid, together with DNA encoding the ADH2/GAPDH promoter, a mammalian signal peptide, or, for example, a yeast alpha-factor or invertase secretory signal/leader sequence, and linker sequences (if needed) for expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72.

Yeast cells, such as yeast strain AB110, can then be transformed with the expression plasmids described above and cultured in selected fermentation media. The transformed yeast supernatants can be analyzed by precipitation with 10% trichloroacetic acid and separation by SDS-PAGE, followed by staining of the gels with Coomassie Blue stain.

Recombinant PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can

5

Expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 in Baculovirus-Infected Insect Cells

10

15

25

30

analyzed by SDS-PAGE and silver staining or Western blot with Ni^{2+} -NTA-conjugated to alkaline phosphatase (Qiagen). Fractions containing the eluted His₁₀-tagged PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 are pooled and dialyzed against loading buffer.

Alternatively, purification of the IgG tagged (or Fc tagged) PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 can be performed using known chromatography techniques, including for instance, Protein A or protein G column chromatography.

EXAMPLE 7

Preparation of Antibodies that Bind PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72

This example illustrates preparation of monoclonal antibodies which can specifically bind PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72.

Techniques for producing the monoclonal antibodies are known in the art and are described, for instance, in Goding, *supra*. Immunogens that can be employed include purified PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, fusion proteins containing PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, and cells expressing recombinant PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 on the cell surface. Selection of the immunogen can be made by the skilled artisan without undue experimentation.

Mice, such as Balb/c, are immunized with the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 immunogen emulsified in complete Freund's adjuvant and injected subcutaneously or intraperitoneally in an amount from 1-100 micrograms. Alternatively, the immunogen is emulsified in MPL-TDM adjuvant (Ribi Immunochemical Research, Hamilton, MT) and injected into the animal's hind foot pads. The immunized mice are then boosted 10 to 12 days later with additional immunogen emulsified in the selected adjuvant. Thereafter, for several weeks, the mice can also be boosted with additional immunization injections. Serum samples can be periodically obtained from the mice by retro-orbital bleeding for testing in ELISA assays to detect anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibodies.

After a suitable antibody titer has been detected, the animals "positive" for antibodies can be injected with a final intravenous injection of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. Three to four days later, the mice are sacrificed and the spleen cells are harvested. The spleen cells are then fused (using 35% polyethylene glycol) to a selected murine myeloma cell line such as P3X63AgU.1, available from ATCC, No. CRL 1597. The fusions generate hybridoma cells which can then be plated in 96 well tissue culture plates containing HAT (hypoxanthine, aminopterin, and thymidine) medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

The hybridoma cells will be screened in an ELISA for reactivity against PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72. Determination of "positive" hybridoma cells secreting the desired monoclonal antibodies against PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 is within the skill in the art.

The positive hybridoma cells can be injected intraperitoneally into syngeneic Balb/c mice to produce

ascites containing the anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 monoclonal antibodies. Alternatively, the hybridoma cells can be grown in tissue culture flasks or roller bottles. Purification of the monoclonal antibodies produced in the ascites can be accomplished using ammonium sulfate precipitation, followed by gel exclusion chromatography. Alternatively, affinity chromatography based upon binding of antibody to protein A or protein G can be employed.

EXAMPLE 8

Purification of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 Polypeptides

Using Specific Antibodies

Native or recombinant PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides can be purified by a variety of standard techniques in the art of protein purification. For example, pro-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, mature PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or pre-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide is purified by immunoaffinity chromatography using antibodies specific for the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide of interest. In general, an immunoaffinity column is constructed by covalently coupling the anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide antibody to an activated chromatographic resin.

Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway, N.J.). Likewise, monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated SEPHAROSE™ (Pharmacia LKB Biotechnology). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

Such an immunoaffinity column is utilized in the purification of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide by preparing a fraction from cells containing PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide in a soluble form. This preparation is derived by solubilization of the whole cell or of a subcellular fraction obtained via differential centrifugation by the addition of detergent or by other methods well known in the art. Alternatively, soluble PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide can be secreted in useful quantity into the medium in which the cells are grown, by employing a heterologous secretion signal peptide.

A soluble PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide (e.g., high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide binding (e.g., a low pH buffer such as approximately pH 2-3, or a high

concentration of a chaotrope such as urea or thiocyanate ion), and PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide is collected.

EXAMPLE 9

Drug Screening

5 This invention is particularly useful for screening compounds by using PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides or binding fragment thereof in any of a variety of drug screening techniques. The PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or fragment employed in such a test can either be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. One method of drug screening utilizes eukaryotic or
10 prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One can measure, for example, the formation of complexes between PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or a
15 fragment and the agent being tested. Alternatively, one can examine the diminution in complex formation between the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide and its target cell or target receptors caused by the agent being tested.

Thus, the present invention provides methods of screening for drugs or any other agents which can affect a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide-
20 associated disease or disorder. These methods comprise contacting such an agent with an PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or fragment thereof and assaying (i) for the presence of a complex between the agent and the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or fragment, or (ii) for the presence of a complex between the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or fragment and the cell,
25 by methods well known in the art. In such competitive binding assays, the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or fragment is typically labeled. After suitable incubation, free PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or fragment is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of the particular agent to bind to PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or to interfere with the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide/cell complex.
30

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to a polypeptide and is described in detail in WO 84/03564, published on September 13, 1984. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid
35 substrate, such as plastic pins or some other surface. As applied to a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, the peptide test compounds are reacted with PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide and washed. Bound PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide is detected by

methods well known in the art. Purified PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide can also be coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies can be used to capture the peptide and immobilize it on the solid support.

5 This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide specifically compete with a test compound for binding to PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or fragments thereof. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

EXAMPLE 10

Rational Drug Design

The goal of rational drug design is to produce structural analogs of biologically active polypeptide of interest (*i.e.*, a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide) or of small molecules with which they interact, *e.g.*, agonists, antagonists, or inhibitors. Any of these examples can be used to fashion drugs which are more active or stable forms of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or which enhance or interfere with the function of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide *in vivo* (*c.f.*, Hodgson, Bio/Technology, 9: 19-21 (1991)).

20 In one approach, the three-dimensional structure of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or of an PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide-inhibitor complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide must be ascertained to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide can be gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design analogous PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide-like molecules or to identify efficient inhibitors. Useful examples of rational drug design can include molecules which have improved activity or stability as shown by Braxton and Wells, Biochemistry, 31:7796-7801 (1992) or which act as inhibitors, agonists, or antagonists of native peptides as shown by Athauda *et al.*, J. Biochem., 113:742-746 (1993).

35 It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore upon which subsequent drug design can be based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids would be expected to be an analog of the original receptor. The anti-id could

then be used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides would then act as the pharmacore.

By virtue of the present invention, sufficient amounts of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide can be made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide amino acid sequence provided herein will provide guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

EXAMPLE 11

Preparation of Antisense Oligonucleotides

Oligonucleotide Synthesis. Unsubstituted and substituted phosphodiester (P = O) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard phosphoramidite chemistry with oxidation by iodine. Phosphorothioates (P = S) are synthesized as for the phosphodiester oligonucleotides except the standard oxidation bottle is replaced by 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation wait step is increased to 68 sec and is followed by the capping step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at 55° C (18 hr), the oligonucleotides are purified by precipitating twice with 2.5 volumes of ethanol from a 0.5 M NaCl solution. Phosphinate oligonucleotides are prepared as described in U.S. Pat. No. 5,508,270, herein incorporated by reference. Alkyl phosphonate oligonucleotides are prepared as described in U.S. Pat. No. 4,469,863, herein incorporated by reference. 3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Pat. Nos. 5,610,289 or 5,625,050, herein incorporated by reference. Phosphoramidite oligonucleotides are prepared as described in U.S. Pat. No., 5,256,775 or U.S. Pat. No. 5,366,878, herein incorporated by reference. Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively), herein incorporated by reference. 3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Pat. No. 5,476,925, herein incorporated by reference. Phosphotriester oligonucleotides are prepared as described in U.S. Pat. No. 5,023,243, herein incorporated by reference. Borano phosphate oligonucleotides are prepared as described in U.S. Pat. Nos. 5,130,302 and 5,177,198, both herein incorporated by reference.

Oligonucleoside Synthesis. Methylenemethylimino linked oligonucleosides (MMI linked oligonucleosides), methylenedimethylhydrazo linked oligonucleosides (MDH linked oligonucleosides), and methylenecarbonylamino linked oligonucleosides (amide-3 linked oligonucleosides), and methyleneaminocarbonyl linked oligonucleosides (amide-4 linked oligonucleosides), as well as mixed backbone compounds having, for instance, alternating MMI and P = O or P = S linkages are prepared as described in U.S. Pat. Nos. 5,378,825, 5,386,023, 5,489,677, 5,602,240 and 5,610,289, all of which are herein incorporated by reference. Formacetal and thioformacetal linked oligonucleosides are prepared as described in U.S. Pat. Nos. 5,264,562 and 5,264,564, herein incorporated by reference. Ethylene oxide linked oligonucleosides are prepared as described in U.S. Pat. No. 5,223,618, herein incorporated by reference.

PNA Synthesis. Peptide nucleic acids (PNAs) are prepared in accordance with any of the various

procedures referred to in Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications, Bioorganic & Medicinal Chemistry, 4:5-23 (1996). They are also prepared in accordance with U.S. Pat. Nos. 5,539,082, 5,700,922, and 5,719,262, herein incorporated by reference.

Synthesis of Chimeric Oligonucleotides. [2'-O-Me]-[2'-deoxy]-[2'-O-Me] Chimeric Phosphorothioate

- 5 Oligonucleotides: Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligonucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 380B, as above. Oligonucleotides are synthesized using the automated synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphoramidite for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite for 5' and 3' wings. The standard synthesis cycle is
- 10 modified by increasing the wait step after the delivery of tetrazole and base to 600 sec repeated four times for RNA and twice for 2'-O-methyl. The fully protected oligonucleotide is cleaved from the support and the phosphate group is deprotected in 3:1 ammonia/ethanol at room temperature overnight then lyophilized to dryness. Treatment in methanolic ammonia for 24 hrs at room temperature is then done to deprotect all bases and sample is again lyophilized to dryness. The pellet is resuspended in 1M TBAF in THF for 24 hrs at room
- 15 temperature to deprotect the 2' positions. The reaction is then quenched with 1M TEAA and the sample is then reduced to 1/2 volume by rotovac before being desalted on a G25 size exclusion column. The oligo recovered is then analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry. [2'-O-(2-Methoxyethyl)]-[2'-deoxy]-[2'-O-(Methoxyethyl)] Chimeric Phosphorothioate Oligonucleotides: [2'-O-(2-methoxyethyl)]-[2'-deoxy]-[2'-O-(methoxyethyl)] chimeric phosphorothioate
- 20 oligonucleotides are prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites. [2'-O-(2-Methoxyethyl)Phosphodiester]-[2'-deoxy Phosphorothioate]-[2'-O-(2-Methoxyethyl) Phosphodiester] Chimeric Oligonucleotides: [2'-O-(2-methoxyethyl phosphodiester)]-[2'-deoxy
- 25 phosphorothioate]-[2'-O-(methoxyethyl) phosphodiester] chimeric oligonucleotides are prepared as per the above procedure for the 2'-O-methyl chimeric oligonucleotide with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites, oxidation with iodine to generate the phosphodiester internucleotide linkages within the wing portions of the chimeric structures and sulfurization utilizing 3H-I,2 benzodithiole-3-one 1,1 dioxide (Beaucage Reagent) to generate the phosphorothioate internucleotide linkages for the center gap. Other chimeric oligonucleotides, chimeric oligonucleosides and mixed chimeric
- 30 oligonucleotides/oligonucleosides are synthesized according to U.S. Pat. No. 5,623,065, herein incorporated by reference.

Example 12

Corneal Angiogenesis Assay

- 35 A corneal angiogenesis assay, using VEGF as an inducer of angiogenesis, can be used to test molecules as an antagonist of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 for anti-angiogenic activity, or as an agonist of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 for enhancing angiogenesis. A 1.5 mm incision is made approximately 1 mm from the center of the cornea of isoflurane-ketamine (60-80 mg/kg) xylazine (10-15 mg/kg) anesthetized Sprague-Dawley rats.

Using a curved spatula, the incision is bluntly dissected through the stroma towards the outer canthus of the eye. A Hydron pellet (2 x 20 mm) containing VEGF (200ng), sucralfate (100 µg) with or without (control) test molecule, at various amounts, is inserted into the base of the pocket. After surgery, the eyes are coated with gentamicin ointment. Animals are observed at 24-48 hr for the occurrence of nonspecific inflammation and then daily thereafter. At day 6, the animals are euthanized and injected with FITC-dextran to allow for visualization of the vasculature. Corneal whole mounts are made of the enucleated eyes and analyzed for neovascular area using the computer assisted image analysis.

The *in vivo* anti-angiogenic effects of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antagonist are shown using Hydron pellets containing 200 ng recombinant VEGF, with or without various amounts of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antagonist, implanted into the corneas of Sprague-Dawley rats as described above. In vivo enhancement of angiogenesis can also be studied in this system. Data from this experiment will show that pellets containing the combination PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antagonist and VEGF can produce a significant reduction in vessel length compared to the VEGF only (positive) controls. These *in vivo* data are consistent with the *in vitro* results, i.e., antagonism of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 produces strong inhibition of angiogenesis in endothelial tissue. Conversely, agonism of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 produces enhancement of angiogenesis in endothelial tissue.

Example 13

Endothelial Proliferation Assay:

HUVEC are seeded on collagen-coated 96-well plates at 6,000 cells/cm² in Clonetics EGM supplemented with 10% FBS, 2 mM L-glutamine, 100U/ml penicillin and 100 U/ml streptomycin and allowed to attach for 4 hr. Medium is then replaced with 1X basal medium consisting of M199 supplemented with 1%FBS, 1X ITS, 2 mM L-glutamine, 50 mg/ml ascorbic acid, 26.5 mM NaHCO₃, 100U/ml penicillin and 100 U/ml streptomycin supplemented with 40 ng/ml bFGF, 40 ng/ml VEGF and 80 nM PMA. Cells are cultured in above medium in the presence of test drugs or vehicle for 4 hr. Then 5 ml (100 mM) of 5'-bromo-2'-deoxyuridine (BrdU) is added in a final volume of 100 ml/well and cells are incubated for another 20 hr. BrdU incorporation is evaluated by an ELISA kit from Boehringer Mannheim (Indianapolis, IN).

Data are expressed as the mean ± standard error. Statistic analysis is performed using one-way ANOVA (INSTAT, Graph Pad Software, Sorrento Valley, CA). Multiple comparisons against the control are analyzed using Bonferroni modification of Student's t-test to determine differences between groups. A p value < 0.05 is accepted as significant.

This assay can indicate that PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antagonists can repress or inhibit growth factor-induced endothelial cell proliferation. BrdU incorporation assays are performed to detect proliferation of endothelial cells cultured on type I collagen-coated surface in medium containing the growth factors, e.g. VEGF, bFGF and PMA, in the presence of vehicle or PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antagonists. The present invention, therefore, is also directed to a method of inhibiting growth factor induced endothelial cell

proliferation by contacting endothelial cells with an effective amount of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antagonist. Conditions associated with undesired vascularization and angiogenesis resulting from growth factor induced endothelial cell proliferation can be treated by administering the antagonists in the manner described herein.

PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 also regulate gene expression events associated with angiogenesis. Flk/KDR and Flt-1 are two structurally related endothelial cell tyrosine kinase receptors for VEGF. The importance of these two receptors during angiogenesis has been clearly demonstrated by the findings that KDR functions as a transducer to signal endothelial cell proliferation and differentiation and that Flt-1 is a critical survival factor involved in endothelial cell morphogenesis (Fong, G.H. et al., (1995) *Nature* (London) 376:66-70; Ferrara, N. et al., (1997) *Endocr. Rev.* 18:4-25; Ilan, N. et al., (1998) *J. Cell Sci.* 111:3621-3631). Whether enhancing or blocking PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 activity alters Flt-1 and KDR gene expression can be readily determined using real time quantitative RT-PCR, in the system using a mixture of growth factors in HUVEC grown in three dimensional collagen gels. It is also well known that the production of proteases (e.g. plasminogen activators) and their inhibitors (e.g. plasminogen activator inhibitor I, PAI-1) is correlated with endothelial cell degradation of extracellular matrix and migration, the two critical steps of the angiogenic processes. Consequently, the effects of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 agonism or antagonism on gene expression of urokinase type plasminogen (uPA) and PAI-1 in three dimensional collagen gels can be readily determined. Treatment of HUVEC with test drug can either reduced or enhance uPA mRNA at about 4 hr and reduce or enhance PAI-1 gene expression at about 24 hr.

EXAMPLE 14

Stimulation of Endothelial Cell Proliferation

This assay is designed to determine whether PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 shows the ability to stimulate adrenal cortical capillary endothelial cell (ACE) growth.

Bovine adrenal cortical capillary endothelial cells (ACE) (from primary culture, maximum of 12-14 passages) are plated in 96-well plates at 500 cells/well per 100 microliter. Assay media included low glucose DMEM, 10% calf serum, 2 mM glutamine, and 1X penicillin/streptomycin/fungizone. Control wells included the following: (1) no ACE cells added; (2) ACE cells alone; (3) ACE cells plus VEGF (5 ng/ml); and (4) ACE cells plus FGF (5ng/ml). The control or test sample, (in 100 microliter volumes), is then added to the wells (at dilutions of 1%, 0.1% and 0.01%, respectively). The cell cultures are incubated for 6-7 days at 37°C/5%CO₂. After the incubation, the media in the wells is aspirated, and the cells are washed 1X with PBS. An acid phosphatase reaction mixture (100 microliter; 0.1M sodium acetate, pH 5.5, 0.1% Triton X-100, 10 mM p-nitrophenyl phosphate) is then added to each well. After a 2 hour incubation at 37°C, the reaction is stopped by addition of 10 microliters 1N NaOH. Optical density (OD) is measured on a microplate reader at 405 nm.

The activity of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 is calculated as the fold increase in proliferation (as determined by the acid phosphatase activity, OD 405 nm) relative to (1) cell only background, and (2) relative to maximum stimulation by VEGF. VEGF (at 3-10 ng/ml) and FGF (at 1-5 ng/ml) are employed as an activity reference for maximum stimulation. Results of

the assay are considered "positive" if the observed stimulation is $\geq 50\%$ increase over background.

EXAMPLE 15

Inhibition of Vascular Endothelial Growth Factor (VEGF)

Stimulated Proliferation of Endothelial Cell Growth

5 The ability of various PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides to inhibit VEGF stimulated proliferation of endothelial cells can be tested. Specifically, bovine adrenal cortical capillary endothelial cells (ACE) (from primary culture, maximum of 12-14 passages) are plated in 96-well plates at 500 cells/well per 100 microliter. Assay media include low glucose DMEM, 10% calf serum, 2 mM glutamine, and 1X penicillin/streptomycin/fungizone. Control wells include the
10 following: (1) no ACE cells added; (2) ACE cells alone; (3) ACE cells plus 5 ng/ml FGF; (4) ACE cells plus 3 ng/ml VEGF; (5) ACE cells plus 3 ng/ml VEGF plus 1 ng/ml TGF-beta; and (6) ACE cells plus 3 ng/ml VEGF plus 5 ng/ml LIF. The test samples, poly-his tagged PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides (in 100 microliter volumes), are then added to the wells (at dilutions of 1%, 0.1% and 0.01%, respectively). The cell cultures are incubated for 6-7 days at 37°C/5%CO₂. After the
15 incubation, the media in the wells is aspirated, and the cells are washed 1X with PBS. An acid phosphatase reaction mixture (100 microliter; 0.1M sodium acetate, pH 5.5, 0.1% Triton X-100, 10 mM p-nitrophenyl phosphate) is then added to each well. After a 2 hour incubation at 37°C, the reaction is stopped by addition of 10 microliters 1N NaOH. Optical density (OD) is measured on a microplate reader at 405 nm.

The activity of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72
20 polypeptides is calculated as the percent inhibition of VEGF (3 ng/ml) stimulated proliferation (as determined by measuring acid phosphatase activity at OD 405 nm) relative to the cells without stimulation. TGF-beta is employed as an activity reference at 1 ng/ml, since TGF-beta blocks 70-90% of VEGF-stimulated ACE cell proliferation. The results are indicative of the utility of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides in cancer therapy and specifically in inhibiting tumor
25 angiogenesis. The results are considered positive if the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide exhibits 30% or greater inhibition of VEGF stimulation of endothelial cell growth (relative inhibition 30% or greater).

EXAMPLE 16

Induction of c-fos in Endothelial Cells

30 This assay is designed to determine whether PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides show the ability to induce c-fos in endothelial cells.

Human venous umbilical vein endothelial cells (HUVEC, Cell Systems) in growth media (50% Ham's F12 w/o GHT: low glucose, and 50% DMEM without glycine: with NaHCO₃, 1% glutamine, 10 mM HEPES, 10% FBS, 10 ng/ml bFGF) are plated on 96-well microtiter plates at a cell density of 1x10⁴ cells/well. The day
35 after plating, the cells are starved by removing the growth media and treating the cells with 100 μ l/well test samples and controls (positive control: growth media; negative control: 10 mM HEPES, 140 mM NaCl, 4% (w/v) mannitol, pH 6.8). The cells are incubated for 30 minutes at 37°C, in 5%CO₂. The samples are removed, and the first part of the bDNA kit protocol (Chiron Diagnostics, cat. #6005-037) is followed, where each

capitalized reagent/buffer listed below is available from the kit.

Briefly, the amounts of the TM Lysis Buffer and Probes needed for the tests are calculated based on information provided by the manufacturer. The appropriate amounts of thawed Probes are added to the TM Lysis Buffer. The Capture Hybridization Buffer is warmed to room temperature. The bDNA strips are set up in the metal strip holders, and 100 μ l of Capture Hybridization Buffer is added to each b-DNA well needed, followed by incubation for at least 30 minutes. The test plates with the cells are removed from the incubator, and the media is gently removed using the vacuum manifold. 100 μ l of Lysis Hybridization Buffer with Probes are quickly pipetted into each well of the microtiter plates. The plates are then incubated at 55°C for 15 minutes. Upon removal from the incubator, the plates are placed on the vortex mixer with the microtiter adapter head and vortexed on the #2 setting for one minute. 80 μ l of the lysate is removed and added to the bDNA wells containing the Capture Hybridization Buffer, and pipetted up and down to mix. The plates are incubated at 53°C for at least 16 hours.

On the next day, the second part of the bDNA kit protocol is followed. Specifically, the plates are removed from the incubator and placed on the bench to cool for 10 minutes. The volumes of additions needed are calculated based upon information provided by the manufacturer. An Amplifier Working Solution is prepared by making a 1:100 dilution of the Amplifier Concentrate (20 fm/ μ l) in AL Hybridization Buffer. The hybridization mixture is removed from the plates and washed twice with Wash A. 50 μ l of Amplifier Working Solution is added to each well and the wells are incubated at 53°C for 30 minutes. The plates are then removed from the incubator and allowed to cool for 10 minutes. The Label Probe Working Solution is prepared by making a 1:100 dilution of Label Concentrate (40 pmoles/ μ l) in AL Hybridization Buffer. After the 10-minute cool-down period, the amplifier hybridization mixture is removed and the plates are washed twice with Wash A. 50 μ l of Label Probe Working Solution is added to each well and the wells are incubated at 53°C for 15 minutes. After cooling for 10 minutes, the Substrate is warmed to room temperature. Upon addition of 3 μ l of Substrate Enhancer to each ml of Substrate needed for the assay, the plates are allowed to cool for 10 minutes, the label hybridization mixture is removed, and the plates are washed twice with Wash A and three times with Wash D. 50 μ l of the Substrate Solution with Enhancer is added to each well. The plates are incubated for 30 minutes at 37°C and RLU is read in an appropriate luminometer.

The replicates are averaged and the coefficient of variation is determined. The measure of activity of the fold increase over the negative control (HEPES buffer) value is indicated by chemiluminescence units (RLU). The results are considered positive if the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide exhibits at least a two-fold value over the negative control. Typically a negative control = about 1.00 RLU at 1.00% dilution, and a Positive control = about 8.39 RLU at 1.00% dilution.

EXAMPLE 17

Human Venous Endothelial Cell Calcium Flux Assay

This assay is designed to determine whether PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptides show the ability to stimulate calcium flux in human umbilical vein endothelial cells (HUVEC, Cell Systems). Calcium influx is a well documented response upon binding of certain

ligands to their receptors. A test compound that results in a positive response in the present Ca influx assay can be said to bind to a specific receptor and activate a biological signaling pathway in human endothelial cells. This could ultimately lead, for example to cell division, inhibition of cell proliferation, endothelial tube formation, cell migration, apoptosis, etc.

Human venous umbilical vein endothelial cells (HUVEC, Cell Systems) in growth media (50:50 without glycine, 1% glutamine, 10mM Hepes, 10% FBS, 10 ng/ml Bfgf), are plated on 96-well microtiter ViewPlates-96 (Packard Instrument Company Part #6005182) microtiter plates at a cell density of 2×10^4 cells/well. The day after plating, the cells are washed three times with buffer (HBSS plus 10 mM Hepes), leaving 100 μ l/well. Then 100 μ l/well of 8 μ M Fluo-3 (2x) is added. The cells are incubated for 1.5 hours at 37°C/5%CO₂. After incubation, the cells are then washed 3x with buffer (described above) leaving 100 μ l/well. Test drugs are prepared on different 96-well plates at 5x concentration in buffer. The positive control corresponded to 50 μ M ionomycin (5x); the negative control corresponded to Protein 32. Cell plate and sample plates are run on a FLIPR (Molecular Devices) machine. The FLIPR machine added 25 μ l of test sample to the cells, and readings are taken every second for one minute, then every 3 seconds for the next three minutes.

The fluorescence change from baseline to the maximum rise of the curve (Δ change) is calculated, and replicates averaged. The rate of fluorescence increase is monitored, and only those samples which had a Δ change greater than 1000 and a rise within 60 seconds, are considered positive. Results are expressed relative to the positive control.

EXAMPLE 18

Endothelial Cell Tube Formation Assay

As an alternative to the tube formation assay described in Example 1, either of the following tube formation assays can be used. In the tube formation assay, agents that stimulate or inhibit endothelial tube formation, including agents involved in stimulating or inhibiting tracking, chemotaxis, and/or endothelial shape change, in the 3-dimensional matrix, and in particular those that stimulate endothelial cells to differentiate into a tube-like structure in a 3-dimensional matrix in the presence of an exogenous growth factor (e.g., VEGF, bFGF), can be readily identified. These agents can be agonists or antagonists as described herein.

Matrigel Tube Formation Assay. 0.5 ml of Matrigel (Becton Dickinson #4023) is pipeted on to the surface of each well of a 24 well tissue culture plate. The matrigel is allowed to solidify by incubation at 37°C for 20 min. Human umbilical vein endothelial cells are resuspended in culture medium (Medium 199, supplemented with 1% fetal bovine serum, 1x insulin-transferrin-selenium (ITS) solution, 2 mmol/L L-glutamine, 26.5 NaHCO₃, 100 U/ml penicillin, 100 U/ml streptomycin) at 2×10^5 cells per ml. One of the following can be added: (a) no additives; (b) basic fibroblast growth factor (40 ng/ml); (c) vascular endothelial cell growth factor (40 ng/ml); (d) phorbol myristate acetate (80 nM); (e) the combination of (b), (c) and (d); and, anyone or more of (a), (b), (c), (d) or (e) in combination with added test agent at various concentrations. Then 200 μ l per well of the cell suspension is added to top of the solidified Matrigel (about 40,000 cells/well). The cells are then cultured in a humidified 5% CO₂ incubator and observed at 4, 8 and 24 hrs. Activity in the assay is measured as an alteration in the formation of tube-like structures, which can be quantitated by measurement of the length of tube-like structures formed and/or the area of the culture covered by the tube-like

network.

Fibrin gel tube formation assay. A thrombin solution is prepared by the addition of 2 ml of basal medium (Medium 199 supplemented with 1% fetal bovine serum, 1x insulin-transferrin-selenium (ITS), 2 mmol/L L-glutamine, 26.5 NaHCO₃, 100 U/ml penicillin, 100 U/ml streptomycin) to 100U thrombin (Sigma Chemical Company, Catalog 6634). This is kept on ice. A fibrinogen solution is prepared by dissolving 112 mg of fibrinogen (Sigma Chemical Company, Catalog #F-4883) in 44 ml of basal medium. Human umbilical endothelial cells are then suspended at a final concentration of 4×10^5 cells per ml in the fibrinogen solution. The thrombin solution is then aliquoted (10 ul/well) to wells of a 48 well tissue culture plate on ice. Then 300 ul of the HUVEC/Fibrinogen mix is added to each thrombin containing well and mixed by pipetting up and down 3 to 5 times. The plates are incubated at 37°C for 20 minutes to allow solidification of the fibrin gel. Basal media is then added with one of the following: (a) no additives; (b) basic fibroblast growth factor (40 ng/ml); (c) vascular endothelial cell growth factor (40 ng/ml); (d) phorbol myristate acetate (80 nM); (e) the combination of (b), (c) and (d); and, anyone or more of (a), (b), (c), (d) or (e) in combination with added test agent at various concentrations. Activity in the assay is measured as an alteration in the formation of tube-like structures, which can be quantitated by measurement of the length of tube-like structures formed and/or the area of the culture covered by the tube-like network.

Agents, either agonists or antagonists (e.g., those that block PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 expression), can also be quickly screened with the following scoring in which a positive result is equal to or greater than 2: score of 1, cells are all round; score of 2, cells are elongated; score of 3, cells are forming tubes with some connections; and score of 4, cells are forming complex tubular networks.

Optionally, one can add to the cell solutions about 1 μ M 6-FAM-FITC dye to stain vacuoles while they are forming. And, after incubation, cells can be fixed with 3.7% formalin at room temperature for 10 minutes, washed with PBS five times, then stained with Rh-Phalloidin at 4°C overnight followed by nuclear staining with 4 μ M DAPI. Subsequently, the cells can be scored for the effect of agent on apoptosis, allowing one to identify factors that facilitate or reduce cell survival in the 3-dimensional matrix, particularly in the presence of exogenous growth factors (e.g., VEGF, bFGF). In this apoptosis assay, a positive result is equal to or less than about 1, where 0 = no apoptosis, 1 = less than about 20% cells are apoptotic, 2 = less than about 50% cells are apoptotic, and 3 = greater than about 50% cells are apoptotic. In addition, with the addition of vacuole stain, one can identify factors that stimulate or reduce endothelial vacuole formation and lumen formation in the presence of bFGF or VEGF (40 ng/ml). along with 1 μ M 6-FAM-FITC dye to stain vacuoles while they are forming. Cells are incubated at 37°C/5% CO₂ for 48 hr, fixed with 3.7% formalin at room temperature for 10 minutes, washed with PBS five times, then stained with Rh-Phalloidin at 4°C overnight followed by nuclear staining with 4 μ M DAPI. A positive result is equal to or greater than 2: 1 = vacuoles present in less than 20% of cells, 2 = vacuoles present in 20-50% of cells, 3 = vacuoles present in greater than 50% of cells. This assay is designed to identify factors that are involved in stimulating pinocytosis, ion pumping, permeability, and junction formation.

Deposit of Material

The following materials have been deposited with the American Type Culture Collection, 10801

University Blvd., Manassas, VA 20110-2209, USA (ATCC):

<u>Material</u>	<u>ATCC Dep. No.</u>	<u>Deposit Date</u>
DNA-C-MG.2-1776	PTA-799	September 28, 1999
DNA-C-MG.12-1776	PTA-798	September 28, 1999

5 This deposit was made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of a viable culture of the deposit for 30 years from the date of deposit. The deposit will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the progeny of the culture of the deposit to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC §122 and the Commissioner's rules pursuant thereto (including 37 CFR §1.14 with particular reference to 886 OG 638).

15 The assignee of the present application has agreed that if a culture of the materials on deposit should die or be lost or destroyed when cultivated under suitable conditions, the materials will be promptly replaced on notification with another of the same. Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

20 The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the construct deposited, which is intended as a single illustration of certain embodiments of the invention and any constructs that are functionally equivalent are within the scope of this invention. The deposit of material herein does not constitute an admission that the written description herein contained is inadequate to enable the practice of any embodiment of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims to the specific illustrations that it represents. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule that comprises a nucleotide sequence having at least about 80% sequence identity to (a) a nucleotide sequence encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide comprising the sequence of amino acid residues from about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively, or (b) the complement of the nucleotide sequence of (a).
2. The isolated nucleic acid molecule of Claim 1, comprising the nucleotide sequence from about 66 to about 1796 of SEQ ID NO:1, about 465 to about 1886 of SEQ ID NO:3, about 271 to about 1788 of SEQ ID NO:17, about 267 to about 1298 of SEQ ID NO:15, or about 71 to about 2059 of SEQ ID NO:13, respectively.
3. The isolated nucleic acid molecule of Claim 1, comprising the nucleotide sequence of SEQ ID NO:1 or SEQ ID NO:3, respectively.
4. The isolated nucleic acid molecule of Claim 1, comprising a nucleotide sequence that encodes the sequence of amino acid residues from about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively.
5. An isolated nucleic acid molecule comprising a nucleotide sequence that comprises at least about 80% sequence identity to (a) a nucleotide sequence encoding the polypeptide encoded by the human protein cDNA deposited with the ATCC on September 28, 1999, under ATCC Deposit No. PTA-799, on September 28, 1999, under ATCC Deposit No. PTA-798, (DNA-C-MG.2-177 and DNA-C-MG.12-1776, respectively), or (b) the complement of the DNA molecule of (a).
6. The isolated nucleic acid molecule of Claim 5, comprising a nucleotide sequence encoding the polypeptide encoded by the human protein cDNA deposited with the ATCC on September 28, 1999, under ATCC Deposit No. PTA-799, on September 28, 1999, under ATCC Deposit No. PTA-798, (DNA-C-MG.2-1776 and DNA-C-MG.12-1776, respectively).
7. An isolated nucleic acid molecule comprising a nucleotide sequence that comprises at least about 80% sequence identity to (a) the full-length polypeptide coding sequence of the human protein cDNA deposited with the ATCC on September 28, 1999, under ATCC Deposit No. PTA-799, on September 28, 1999, under ATCC Deposit No. PTA-798, (DNA-C-MG.2-1776 and, DNA-C-MG.12-1776, respectively), or (b) the complement of the coding sequence of (a).

8. The isolated nucleic acid molecule of Claim 7 comprising the full-length polypeptide coding sequence of the human protein cDNA deposited with the ATCC on September 28, 1999, under ATCC Deposit No. PTA-799, on September 28, 1999, under ATCC Deposit No. PTA-798, (DNA-C-MG.2-1776 and DNA-C-MG.12-1776, respectively).

5

9. An isolated nucleic acid molecule encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide comprising nucleic acid that hybridizes to the complement of the nucleic acid sequence that encodes amino acids about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively.

10

10. The isolated nucleic acid molecule of Claim 9, wherein the nucleic acid that encodes amino acids about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively, comprises nucleotides about 66 to about 1796 of SEQ ID NO:1, about 465 to about 1886 of SEQ ID NO:3, about 271 to about 1788 of SEQ ID NO:17, about 267 to about 1298 of SEQ ID NO:15, or about 71 to about 2059 of SEQ ID NO:13, respectively.

15

11. The isolated nucleic acid molecule of Claim 9, wherein the hybridization occurs under stringent hybridization or wash conditions.

20

12. An isolated nucleic acid molecule comprising (a) a nucleic acid sequence encoding a polypeptide scoring at least 80% positives when compared to the amino acid sequence from about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively, or (b) the complement of the DNA of (a).

25

13. An isolated nucleic acid molecule comprising at least about 596 or 1535 nucleotides, for PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 respectively, and which is produced by hybridizing a test DNA molecule under stringent hybridization conditions with (a) a DNA molecule which encodes a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide comprising a sequence of amino acid residues from about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively, or (b) the complement of the DNA molecule of (a), and isolating the test DNA molecule.

30

35

14. The isolated nucleic acid molecule of Claim 13, which has at least about 80% sequence identity to (a) or (b).

15. A vector comprising the nucleic acid molecule of Claim 1.

16. The vector of Claim 15, wherein the nucleic acid molecule is operably linked to control sequences recognized by a host cell transformed with the vector.

5

17. A nucleic acid molecule deposited with the ATCC on September 28, 1999, under ATCC Deposit No. PTA-799, on September 28, 1999, under ATCC Deposit No. PTA-798, (DNA-C-MG.2-1776 and DNA-C-MG.12-1776, respectively).

10

18. A host cell comprising the vector of Claim 15.

19. The host cell of Claim 18, wherein the cell is a CHO cell.

20. The host cell of Claim 18, wherein the cell is an *E. coli*.

15

21. The host cell of Claim 18, wherein the cell is a yeast cell.

20

22. A process for producing a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide comprising culturing the host cell of Claim 18 under conditions suitable for expression of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, wherein the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide is produced.

25

23. The process of claim 22, further comprising the step of recovering the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide from the cell culture.

30

24. An isolated PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide comprising an amino acid sequence comprising at least about 80% sequence identity to the sequence of amino acid residues from about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively.

35

25. The isolated PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide of Claim 24 comprising amino acid residues about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively.

26. An isolated PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72

polypeptide having at least about 80% sequence identity to the polypeptide encoded by the cDNA insert of the vector deposited with the ATCC on September 28, 1999, under ATCC Deposit No. PTA-799, on September 28, 1999, under ATCC Deposit No. PTA-798, (DNA-C-MG.2-1776 and DNA-C-MG.12-1776, respectively).

5 27. The isolated PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide of Claim 26 which is encoded by the cDNA insert of the vector deposited with the ATCC on September 28, 1999, under ATCC Deposit No. PTA-799, on September 28, 1999, under ATCC Deposit No. PTA-798, (DNA-C-MG.2-1776 and DNA-C-MG.12-1776, respectively).

10 28. An isolated PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide scoring at least 80% positives when compared to the sequence of amino acid residues from about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively.

15 29. An isolated PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide comprising the sequence of amino acid residues from about 1 to about 577 of SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively, or a fragment thereof sufficient to provide a
20 antibody.

 30. An isolated polypeptide produced by (i) hybridizing a test DNA molecule under stringent conditions with (a) a DNA molecule encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide comprising the sequence of amino acid residues from about 1 to about 577 of
25 SEQ ID NO:2, about 1 to about 474 of SEQ ID NO: 4, about 1 to about 506 of SEQ ID NO: 18, about 1 to about 344 of SEQ ID NO: 16, or about 1 to about 633 of SEQ ID NO: 14, respectively, or (b) the complement of the DNA molecule of (a), (ii) culturing a host cell comprising the test DNA molecule under conditions suitable for the expression of the polypeptide, and (iii) recovering the polypeptide from the cell culture.

30 31. The isolated polypeptide of Claim 30, wherein the test DNA has at least about 80% sequence identity to (a) or (b).

 32. A chimeric molecule comprising a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide fused to a heterologous amino acid sequence.

35 33. The chimeric molecule of Claim 32, wherein the heterologous amino acid sequence is an epitope tag sequence.

34. The chimeric molecule of claim 32, wherein the heterologous amino acid sequence is a secretion signal peptide.

35. The chimeric molecule of Claim 32, wherein the heterologous amino acid sequence is a Fc region of an immunoglobulin.

36. An antibody which specifically binds to a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

37. The antibody of Claim 36, wherein the antibody is a monoclonal antibody.

38. The antibody of Claim 36, wherein the antibody is a humanized antibody.

39. The antibody of Claim 36, wherein the antibody is an antibody fragment.

40. An antigen compound targeted to a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 gene sequence, wherein said antigen compound inhibits the expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72.

41. The antigen compound of claim 40, which is an antisense oligonucleotide comprising 5 to 60 nucleobases.

42. The antigen compound of claim 41 wherein the antisense oligonucleotide comprises at least one modified internucleoside linkage.

43. The antigen compound of claim 41 wherein the modified internucleoside linkage is a phosphorothioate linkage.

44. The antigen compound of claim 41 wherein the antisense oligonucleotide comprises at least one modified sugar moiety.

45. The antigen compound of claim 44 wherein the modified sugar moiety is a 2'-O-methoxyethyl sugar moiety.

46. The antigen compound of claim 41 wherein the antisense oligonucleotide comprises at least one modified nucleobase.

47. The antigen compound of claim 46 wherein the modified nucleobase is a 5-methylcytosine.

48. The antigene compound of claim 41 wherein the antisense oligonucleotide is a chimeric oligonucleotide.

49. The antigene compound of claim 41 that is a peptide nucleic acid or a ribozyme.

50. The antigene compound of claim 40, wherein the 5 to 60 nucleobases comprises a sequence from SEQ ID NO. 1.

51. A vector comprising a DNA which encodes the antigene oligonucleotide of claim 40.

52. A host cell comprising the vector of claim 51.

53. A method of inhibiting the expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 in human cells or tissues in vitro comprising contacting the cells or tissues with the antigene compound of claim 40 so that expression of PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 is inhibited.

54. A pharmaceutical preparation for treating a condition related to PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 expression by a cell, comprising a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene which enters the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-expressing cell and binds specifically to a nucleic acid molecule encoding PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72, said antigene being present in a biologically acceptable carrier.

55. The pharmaceutical preparation according to claim 54, which further comprises at least one agent for improving delivery of said antigene to the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-expressing cell.

56. The pharmaceutical preparation according to claim 55, wherein the at least one agent comprises a lipid.

57. The pharmaceutical preparation according to claim 55, which further comprises at least one targeting agent for improving cell-type specific delivery of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antigene.

58. An agonist to a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

59. An antagonist to a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

60. A composition comprising (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) an agonist to a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (c) an antagonist to a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (d) an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody in admixture with a pharmaceutically acceptable carrier.

61. The composition of Claim 60 comprising a therapeutically effective amount of (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (c) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, thereof.

62. The composition of Claim 60, further comprising a cardiovascular, endothelial, angiogenic, or angiostatic agent.

63. The composition of Claim 62, wherein the polypeptide or antibody and the cardiovascular, endothelial, angiogenic or angiostatic agent are present in therapeutically effective amounts.

64. The composition of Claim 60, wherein the agonist is a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72-encoding nucleic acid.

65. The composition of Claim 60, wherein the antagonist is an PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antisense molecule or antibody.

66. The composition of Claim 60 further comprising a cardiovascular, endothelial, angiogenic or angiostatic agent.

67. A method of preparing the composition of Claim 60 comprising admixing (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (c) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, with a pharmaceutically acceptable carrier.

68. An article of manufacture comprising:

(a) a composition comprising (i) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (ii) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45,

PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (iii) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, in admixture with a pharmaceutically acceptable carrier;

(b) a container containing the composition; and

(c) a label affixed to said container, or a package insert included in said pharmaceutical product referring to the use of (a) the treatment of an angiogenic disorder.

69. The article of manufacture of Claim 68, wherein the agonist is an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody or antigene compound.

70. The article of manufacture of Claim 68, wherein said antagonist is an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody or antigene compound.

71. The article of manufacture of Claim 68, wherein the composition comprises a therapeutically effective amount of said (a) PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or (c) antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, in admixture with said pharmaceutically acceptable carrier.

72. A method for identifying an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide comprising:

(a) contacting cells and a test compound to be screened under conditions suitable for the induction of a cellular response normally enhanced by a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide; and

(b) determining the induction of said cellular response to determine if the test compound is an effective agonist, wherein the induction of said cellular response is indicative of said test compound being an effective agonist.

73. The method of Claim 72, wherein the cellular response normally induced by said PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide is stimulation of tube formation.

74. A method for identifying a compound that inhibits an activity of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide comprising contacting a test compound with a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide under conditions and for a time sufficient to allow the test compound and polypeptide to interact and determining whether the activity of said PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide is inhibited.

75. A method for identifying a compound that inhibits the expression of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or gene in cells that normally expresses the polypeptide, wherein the method comprises contacting the cells with a test compound under conditions suitable for allowing expression of said PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide and determining whether the expression of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or gene is inhibited.

76. The method of claim 75 wherein the compound is an antigene compound.

77. The method of Claim 76, wherein said antigene compound is an antisense oligonucleotide.

78. A method for diagnosing a disease or susceptibility to a disease which is related to a mutation in a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide-encoding nucleic acid sequence comprising:

(a) isolating a nucleic acid sequence encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide from a sample derived from a host; and

(b) determining the presence or absence of said mutation in the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide nucleic acid sequence, wherein the presence or absence of said mutation is indicative of the presence of said disease or susceptibility to said disease.

79. A method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal which comprises analyzing the level of expression of a gene encoding a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide (a) in a test sample of tissue cells obtained from said mammal, and (b) in a control sample of known normal tissue cells of the same cell type, wherein a higher or lower expression level in the test sample as compared to the control sample is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in said mammal.

80. A method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal which comprises detecting the presence or absence of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide in a test sample of tissue cells obtained from said mammal, wherein the presence or absence of said PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide in said test sample is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in said mammal.

81. A method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal comprising (a) contacting an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody with a test sample of tissue cells obtained from the mammal, and (b) detecting the formation of a complex between the anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72

antibody and a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide in the test sample, wherein the formation of said complex is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in the mammal.

5 82. A method for determining the presence of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide in a sample comprising contacting a sample suspected of containing a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide with an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody and determining binding of said antibody to a component of said sample.

10

 83. A cardiovascular, endothelial or angiogenic disorder diagnostic kit comprising an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody or antigene and a carrier.

 84. The kit of Claim 83 further comprising instructions for using said antibody to detect the presence of the PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide or antigene.

15

 85. A method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal a therapeutically effective amount of (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (c) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

20

 86. The method of Claim 85, wherein the disorder is vascular trauma or cancer.

25

 87. The method of Claim 85, wherein the amount of (a), (b) or (c) is administered together with a cardiovascular, endothelial, angiogenic, or angiostatic agent.

 88. The method of Claim 85, wherein the amount of (a), (b) or (c) is administered in combination with a chemotherapeutic agent, a growth inhibitory agent, a cytotoxic agent, or an angiostatic agent.

30

 89. The method of Claim 85 wherein said agonist is an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody or antigene.

 90. The method of Claim 85 wherein said antagonist is an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody or antigene.

35

 91. A method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal

comprising administering to the mammal a nucleic acid molecule that encodes (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (c) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

5

92. The method of Claim 91 wherein said agonist is an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody or antigene.

93. The method of Claim 91 wherein said antagonist is an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody or antigene.

10

94. The method of Claim 91, wherein the nucleic acid molecule is administered via *ex vivo* gene therapy.

95. The method of Claim 91, wherein the cardiovascular, endothelial or angiogenic disorder is vascular trauma or a cancer.

15

96. A recombinant retroviral particle comprising a retroviral vector consisting essentially of a promoter, nucleic acid encoding (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) an agonist polypeptide of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (c) an antagonist polypeptide of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, and a signal sequence for cellular secretion of the polypeptide, wherein the retroviral vector is in association with retroviral structural proteins.

20

97. An *ex vivo* producer cell comprising a nucleic acid construct that expresses retroviral structural proteins and also comprises a retroviral vector consisting essentially of a promoter, nucleic acid encoding (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) an agonist polypeptide of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (c) an antagonist polypeptide of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, and a signal sequence for cellular secretion of the polypeptide, wherein said producer cell packages the retroviral vector in association with the structural proteins to produce recombinant retroviral particles.

25

30

98. A method for inhibiting endothelial cell growth in a mammal comprising administering to the mammal an effective amount of (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (c) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (d) an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-

35

MG.64 or PRO-C-MG.72 antibody, wherein endothelial cell growth in said mammal is inhibited.

99. A method for stimulating endothelial cell growth in a mammal comprising administering to the mammal an effective amount of (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (c) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (d) an anti-PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 antibody, wherein endothelial cell growth in said mammal is stimulated.

100. A method for inhibiting angiogenesis comprising administering an effective amount of an (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (c) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, to the mammal, wherein said angiogenesis is inhibited.

101. A method for stimulating angiogenesis comprising administering to a mammal an effective amount of (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (c) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, wherein angiogenesis is stimulated.

102. A method for inhibiting endothelial tube formation comprising administering an tube-formation-inducing effective amount of (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (c) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide to the mammal, wherein said tube formation is inhibited.

103. A method for stimulating endothelial tube formation comprising administering to a mammal an tube-formation-inducing effective amount (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (c) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, wherein tube formation is stimulated.

104. A method for treating a tumor, reducing the size of a tumor, reducing the vasculature supporting a tumor, or reducing the tumor burden of a mammal, comprising administering to a mammal in need thereof a therapeutically effective amount of (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (c) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-

MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

105. A method for treating a disease or disorder characterized by undesirable excessive neovascularization, comprising administering to a mammal in need thereof a therapeutically effective amount of (a) a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, (b) an agonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide, or (c) an antagonist of a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 polypeptide.

106. The method of claim 105, wherein the disease or disorder is selected from the group consisting of rheumatoid arthritis, psoriasis, atherosclerosis, retinopathy, retrolental fibroplasia, neovascular glaucoma, age-related macular degeneration, thyroid hyperplasias, Grave's disease, tissue transplantation, chronic inflammation, lung inflammation, and obesity

107. A microarray, the microarray comprising a solid support and a plurality of different oligonucleotides attached to the support, wherein each of the different oligonucleotides is attached to the surface of the solid support in a different predefined region, has a different determinable sequence, and is at least 6 nucleotides in length, wherein at least one of the different oligonucleotides is a PRO-C-MG.2, PRO-C-MG.12, PRO-C-MG.45, PRO-C-MG.64 or PRO-C-MG.72 sequence.

108. The array of claim 107, wherein each different oligonucleotides is from about 5 to about 60 nucleotides in length.

109. The array of claim 107, wherein the array comprises 2 to 1,000,000 different oligonucleotides attached to the solid support.

110. The array of claim 107, wherein the solid support is glass.

111. The array of claim 107, wherein the oligonucleotides are attached to the surface of the solid support through a linker group.

112. The array of claim 107, wherein the oligonucleotides are at least 20% pure.

**NOVEL POLYPEPTIDES, THEIR NUCLEIC ACIDS, AND METHODS FOR THEIR USE IN
ANGIOGENESIS AND VASCULARIZATION**

5

Abstract of the Disclosure

The present invention is directed to novel polypeptides critical for angiogenesis and vascularization, and to nucleic acid molecules encoding those polypeptides. Also provided herein are vectors and host cells comprising those nucleic acid sequences, chimeric polypeptide molecules comprising the polypeptides of the present invention fused to heterologous polypeptide sequences, antibodies which bind to the polypeptides of the present invention and to methods for producing the polypeptides of the present invention. Compositions and methods are disclosed for stimulating or inhibiting angiogenesis and/or neo- or cardio-vascularization in mammals, including humans. Pharmaceutical compositions are based on polypeptides or antagonists thereto that have been identified for one or more of these uses. Disorders that can be diagnosed, prevented, or treated by the compositions herein include trauma such as wounds, various cancers, and disorders of the vessels including atherosclerosis.

005007" 85448960

Sequence Listing

5 <110> Genentech, Inc. and CuraGen Corporation
 <120> NOVEL POLYPEPTIDES, THEIR NUCLEIC ACIDS, AND METHODS
 FOR THEIR USE IN ANGIOGENESIS AND VASCULARIZATION
 10 <130> P1776R2PCT
 <141> 2000-10-05
 <150> US 60/158,587
 15 <151> 1999-10-07
 <150> US 60/162,611
 <151> 1999-10-28
 20 <160> 29
 <210> 1
 <211> 2891
 <212> DNA
 <213> Homo sapiens
 25 <400> 1
 ggcgaggagtc ggcgccctogg gttcctacct cgcgtcccta ggcgggggcg 50
 gccggggcgtc ccggg atg gcc ttc atg gag aag ccg cca 89
 30 Met Ala Phe Met Glu Lys Pro Pro
 1 5
 gcc gcc aag gtg ctg ctg gac gac acg gtg ccg ctg aca 128
 35 Ala Gly Lys Val Leu Leu Asp Asp Thr Val Pro Leu Thr
 10 15 20
 gca gcc atc gag gcg agc cag agc ctg cag tcc cac acg 167
 40 Ala Ala Ile Glu Ala Ser Gln Ser Leu Gln Ser His Thr
 25 30
 gaa tat att att cga gtg caa aga gga att tct gtg gaa 206
 45 Glu Tyr Ile Ile Arg Val Gln Arg Gly Ile Ser Val Glu
 35 40 45
 aac agc tgg cag att gtt aga aga tac agt gac ttt gat 245
 50 Asn Ser Trp Gln Ile Val Arg Arg Tyr Ser Asp Phe Asp
 50 55 60
 ttg ctt aac aac agc tta cag att gca ggc cta agt cta 284
 55 Leu Leu Asn Asn Ser Leu Gln Ile Ala Gly Leu Ser Leu
 65 70
 cct ctt cct ccc aaa aaa ttg att ggt aac atg gat cgt 323
 60 Pro Leu Pro Pro Lys Lys Leu Ile Gly Asn Met Asp Arg
 75 80 85
 gaa ttc ata gct gaa agg cag aaa ggt ctt cag aac tat 362
 65 Glu Phe Ile Ala Glu Arg Gln Lys Gly Leu Gln Asn Tyr
 90 95
 ctc aac gtg atc aca aca aat cat atc ttg tct aat tgt 401
 70 Leu Asn Val Ile Thr Thr Asn His Ile Leu Ser Asn Cys
 100 105 110

gag ctg gtt aag aag ttt tta gat cca aac aac tat tcc 440
 Glu Leu Val Lys Lys Phe Leu Asp Pro Asn Asn Tyr Ser
 115 120 125

5 gca aac tat act gag att gcc ttg caa cag gtt tcc atg 479
 Ala Asn Tyr Thr Glu Ile Ala Leu Gln Gln Val Ser Met
 130 135

10 ttc ttc cga tca gaa cca aag tgg gag gtg gtg gaa cct 518
 Phe Phe Arg Ser Glu Pro Lys Trp Glu Val Val Glu Pro
 140 145 150

15 ttg aaa gac ata ggt tgg aga ata agg aag aaa tat ttc 557
 Leu Lys Asp Ile Gly Trp Arg Ile Arg Lys Lys Tyr Phe
 155 160

20 ttg atg aag att aaa aat cag cca aag gaa cgg cta gtg 596
 Leu Met Lys Ile Lys Asn Gln Pro Lys Glu Arg Leu Val
 165 170 175

25 tta agc tgg gct gac ctt ggc cca gac aag tat ttg tca 635
 Leu Ser Trp Ala Asp Leu Gly Pro Asp Lys Tyr Leu Ser
 180 185 190

30 gat aaa gat ttt cag tgt cta atc aaa ctt ctg cct tct 674
 Asp Lys Asp Phe Gln Cys Leu Ile Lys Leu Leu Pro Ser
 195 200

35 tgt ttg cac cct tac atc tat cgg gtt acc ttt gcc aca 713
 Cys Leu His Pro Tyr Ile Tyr Arg Val Thr Phe Ala Thr
 205 210 215

40 gct aat gaa tcc tca gcg ttg cta att agg atg ttt aac 752
 Ala Asn Glu Ser Ser Ala Leu Leu Ile Arg Met Phe Asn
 220 225

45 gaa aag gga aca ttg aag gat ctg atc tac aag gca aaa 791
 Glu Lys Gly Thr Leu Lys Asp Leu Ile Tyr Lys Ala Lys
 230 235 240

50 cca aaa gac cca ttt cta aag aag tac tgc aac cct aag 830
 Pro Lys Asp Pro Phe Leu Lys Lys Tyr Cys Asn Pro Lys
 245 250 255

55 aag att cag ggc ctg gaa ctc cag caa ata aaa aca tat 869
 Lys Ile Gln Gly Leu Glu Leu Gln Gln Ile Lys Thr Tyr
 260 265

60 gga cgg caa ata tta gag gta ctg aag ttt ctt cat gac 908
 Gly Arg Gln Ile Leu Glu Val Leu Lys Phe Leu His Asp
 270 275 280

aag gga ttc cct tat ggg cat ctt cac gcc tcc aat gtg 947
 Lys Gly Phe Pro Tyr Gly His Leu His Ala Ser Asn Val
 285 290

atg ctc gat ggg gac act tgt cgg ctg ctg gac ctt gag 986
 Met Leu Asp Gly Asp Thr Cys Arg Leu Leu Asp Leu Glu
 295 300 305

aat tcc tta ttg ggc ctg cct tcc ttc tac cga tct tat 1025
 Asn Ser Leu Leu Gly Leu Pro Ser Phe Tyr Arg Ser Tyr
 310 315 320

ttt tca caa ttc agg aaa atc aat aca ttg gaa agt gtg 1064
 Phe Ser Gln Phe Arg Lys Ile Asn Thr Leu Glu Ser Val
 325 330

5 gat gtc cac tgc ttt ggc cac tta ctg tat gaa atg act 1103
 Asp Val His Cys Phe Gly His Leu Leu Tyr Glu Met Thr
 335 340 345

10 tat gga cga ccg cca gac tcg gtg cct gtg gac tcc ttc 1142
 Tyr Gly Arg Pro Pro Asp Ser Val Pro Val Asp Ser Phe
 350 355

15 cct cct gcc ccg tcc atg gct gtg gtg gcc gtg ttg gag 1181
 Pro Pro Ala Pro Ser Met Ala Val Val Ala Val Leu Glu
 360 365 370

20 tct acg ctg tct tgt gaa gcc tgt aaa aat ggc atg cct 1220
 Ser Thr Leu Ser Cys Glu Ala Cys Lys Asn Gly Met Pro
 375 380 385

acc atc tcc cgg ctc tta cag atg cca tta ttc agc gat 1259
 Thr Ile Ser Arg Leu Leu Gln Met Pro Leu Phe Ser Asp
 390 395

25 gtt tta cta acc act tct gaa aaa cca cag ttt aag atc 1298
 Val Leu Leu Thr Thr Ser Glu Lys Pro Gln Phe Lys Ile
 400 405 410

30 cct aca aag tta aaa gag gca ttg aga att gcc aaa gaa 1337
 Pro Thr Lys Leu Lys Glu Ala Leu Arg Ile Ala Lys Glu
 415 420

35 tgt ata gag aag aga cta att gag gaa cag aaa cag att 1376
 Cys Ile Glu Lys Arg Leu Ile Glu Glu Gln Lys Gln Ile
 425 430 435

40 cac cag cat cga aga ctg aca aga gct cag tcc cac cat 1415
 His Gln His Arg Arg Leu Thr Arg Ala Gln Ser His His
 440 445 450

gga tct gag gag gaa aga aaa aaa aga aag att tta gct 1454
 Gly Ser Glu Glu Glu Arg Lys Lys Arg Lys Ile Leu Ala
 455 460

45 cga aag aag tca aaa cga tct gct ctt gaa aat agt gaa 1493
 Arg Lys Lys Ser Lys Arg Ser Ala Leu Glu Asn Ser Glu
 465 470 475

50 gag cat tca gcg agg tac agc aac tcc aat aat tca gga 1532
 Glu His Ser Ala Arg Tyr Ser Asn Ser Asn Asn Ser Gly
 480 485

55 tct ggg gcc agc tca cct ctc acg tcc ccg tca tcg cca 1571
 Ser Gly Ala Ser Ser Pro Leu Thr Ser Pro Ser Ser Pro
 490 495 500

act cca ccc tct aca tca ggg ata tct gca tta cct cca 1610
 Thr Pro Pro Ser Thr Ser Gly Ile Ser Ala Leu Pro Pro
 505 510 515

60 cct cct cca cct cca cca cca cca gca gct ccc ttg cct 1649
 Pro Pro Pro Pro Pro Pro Pro Pro Ala Ala Pro Leu Pro
 520 525

0966453.103500

cct gcg agc acc gag gta cct gcc cag ctc tcg tct cag 1688
 Pro Ala Ser Thr Glu Val Pro Ala Gln Leu Ser Ser Gln
 530 535 540

5 gct gtg aat ggc atg agc cga ggg gcc ttg ctc agc tcc 1727
 Ala Val Asn Gly Met Ser Arg Gly Ala Leu Leu Ser Ser
 545 550

10 atc cag aat ttc caa aaa gga act ttg agg aaa gcc aaa 1766
 Ile Gln Asn Phe Gln Lys Gly Thr Leu Arg Lys Ala Lys
 555 560 565

15 acc tgt gat cac agt gct ccg aag atc ggc tg aa 1800
 Thr Cys Asp His Ser Ala Pro Lys Ile Gly
 570 575 577

gcttcctggtt tacacttgga gggaaaagtt ctttttttatt cctactcacc 1850

20 cctaccccc aaactaccct cttcctggga aagtaattgc tgagccagta 1900
 cagccacaaa cagtactatt ttgcagatgc tcatgtaagc agcttttcga 1950
 gagaaataat tctttaagca gaataaagtt aggctggcat tgctccctta 2000

25 agatcttgct cctttattaa ccctgtaaag gagtcttggt tatectctaa 2050
 tggccaggct tttgggacag cagcatattg aaatattttc accaactaaa 2100
 ggaaatagac agaaaaacaa tgacaatatt caatcacagc agtaaattggc 2150

30 ctttgtgttg caatcccttc taccatca gacagctcct agaaacattc 2200
 cttacagttc atttctctaa agcattttct gattcttaga taactccaat 2250
 ttttgctacc tttatcttag acattaacac tatagcccaa agcatagtta 2300

35 ctttgctaaa tcagaaagca actgagttct ttgttttctc ctcaaataga 2350
 atggggaacg ttcacaacat totcttaagt tctaacagga ataccattgt 2400

40 ggttatagaa ctgaggctg ctaaagcaac tactctagac ccatagttct 2450
 ttttagttag atgtattgaa acagacaaaa atattaacat cagaaaaagc 2500

45 tcttgccaat tagaggatct tottaatcct cagcaattaa gtttggggtt 2550
 tgaggggggc aggtcattgt tacaacagaa gttaaatttg catctataga 2600
 aatcaattat gatttttgaa agatttatct aaatatatca atatagcatc 2650

50 tctttaatgt tagtcattta ttagaaagat cctttatcct gatttgctta 2700
 aacctttcaa taaattgcac tttaaaggat tataaataat ccatttaaaa 2750

55 attcaagtac acacatcagt gttgggttact atgcagagaa tgtcattgtg 2800
 tatagtttca tgtaatctgt tatgtcagct gtatttttta ttaaaatcat 2850

60 gtcaagaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a 2891

<210> 2
 <211> 577
 <212> PRT
 <213> Homo sapiens

<400> 2

	Met	Ala	Phe	Met	Glu	Lys	Pro	Pro	Ala	Gly	Lys	Val	Leu	Leu	Asp	15
	1				5					10						
5	Asp	Thr	Val	Pro	Leu	Thr	Ala	Ala	Ile	Glu	Ala	Ser	Gln	Ser	Leu	30
					20					25						
	Gln	Ser	His	Thr	Glu	Tyr	Ile	Ile	Arg	Val	Gln	Arg	Gly	Ile	Ser	45
					35					40						
10	Val	Glu	Asn	Ser	Trp	Gln	Ile	Val	Arg	Arg	Tyr	Ser	Asp	Phe	Asp	60
					50					55						
	Leu	Leu	Asn	Asn	Ser	Leu	Gln	Ile	Ala	Gly	Leu	Ser	Leu	Pro	Leu	75
15					65					70						
	Pro	Pro	Lys	Lys	Leu	Ile	Gly	Asn	Met	Asp	Arg	Glu	Phe	Ile	Ala	90
					80					85						
20	Glu	Arg	Gln	Lys	Gly	Leu	Gln	Asn	Tyr	Leu	Asn	Val	Ile	Thr	Thr	105
					95					100						
	Asn	His	Ile	Leu	Ser	Asn	Cys	Glu	Leu	Val	Lys	Lys	Phe	Leu	Asp	120
25					110					115						
	Pro	Asn	Asn	Tyr	Ser	Ala	Asn	Tyr	Thr	Glu	Ile	Ala	Leu	Gln	Gln	135
					125					130						
	Val	Ser	Met	Phe	Phe	Arg	Ser	Glu	Pro	Lys	Trp	Glu	Val	Val	Glu	150
30					140					145						
	Pro	Leu	Lys	Asp	Ile	Gly	Trp	Arg	Ile	Arg	Lys	Lys	Tyr	Phe	Leu	165
					155					160						
35	Met	Lys	Ile	Lys	Asn	Gln	Pro	Lys	Glu	Arg	Leu	Val	Leu	Ser	Trp	180
					170					175						
	Ala	Asp	Leu	Gly	Pro	Asp	Lys	Tyr	Leu	Ser	Asp	Lys	Asp	Phe	Gln	195
40					185					190						
	Cys	Leu	Ile	Lys	Leu	Leu	Pro	Ser	Cys	Leu	His	Pro	Tyr	Ile	Tyr	210
					200					205						
	Arg	Val	Thr	Phe	Ala	Thr	Ala	Asn	Glu	Ser	Ser	Ala	Leu	Leu	Ile	225
45					215					220						
	Arg	Met	Phe	Asn	Glu	Lys	Gly	Thr	Leu	Lys	Asp	Leu	Ile	Tyr	Lys	240
					230					235						
50	Ala	Lys	Pro	Lys	Asp	Pro	Phe	Leu	Lys	Lys	Tyr	Cys	Asn	Pro	Lys	255
					245					250						
	Lys	Ile	Gln	Gly	Leu	Glu	Leu	Gln	Gln	Ile	Lys	Thr	Tyr	Gly	Arg	270
					260					265						
55	Gln	Ile	Leu	Glu	Val	Leu	Lys	Phe	Leu	His	Asp	Lys	Gly	Phe	Pro	285
					275					280						
	Tyr	Gly	His	Leu	His	Ala	Ser	Asn	Val	Met	Leu	Asp	Gly	Asp	Thr	300
60					290					295						
	Cys	Arg	Leu	Leu	Asp	Leu	Glu	Asn	Ser	Leu	Leu	Gly	Leu	Pro	Ser	315
					305					310						

5 Phe Tyr Arg Ser Tyr Phe Ser Gln Phe Arg Lys Ile Asn Thr Leu
 320 325 330
 Glu Ser Val Asp Val His Cys Phe Gly His Leu Leu Tyr Glu Met
 335 340 345
 10 Thr Tyr Gly Arg Pro Pro Asp Ser Val Pro Val Asp Ser Phe Pro
 350 355 360
 Pro Ala Pro Ser Met Ala Val Val Ala Val Leu Glu Ser Thr Leu
 365 370 375
 15 Ser Cys Glu Ala Cys Lys Asn Gly Met Pro Thr Ile Ser Arg Leu
 380 385 390
 Leu Gln Met Pro Leu Phe Ser Asp Val Leu Leu Thr Thr Ser Glu
 395 400 405
 20 Lys Pro Gln Phe Lys Ile Pro Thr Lys Leu Lys Glu Ala Leu Arg
 410 415 420
 Ile Ala Lys Glu Cys Ile Glu Lys Arg Leu Ile Glu Glu Gln Lys
 425 430 435
 25 Gln Ile His Gln His Arg Arg Leu Thr Arg Ala Gln Ser His His
 440 445 450
 Gly Ser Glu Glu Glu Arg Lys Lys Arg Lys Ile Leu Ala Arg Lys
 455 460 465
 30 Lys Ser Lys Arg Ser Ala Leu Glu Asn Ser Glu Glu His Ser Ala
 470 475 480
 Arg Tyr Ser Asn Ser Asn Asn Ser Gly Ser Gly Ala Ser Ser Pro
 485 490 495
 Leu Thr Ser Pro Ser Ser Pro Thr Pro Pro Ser Thr Ser Gly Ile
 500 505 510
 40 Ser Ala Leu Pro Pro Pro Pro Pro Pro Pro Pro Pro Ala Ala
 515 520 525
 Pro Leu Pro Pro Ala Ser Thr Glu Val Pro Ala Gln Leu Ser Ser
 530 535 540
 45 Gln Ala Val Asn Gly Met Ser Arg Gly Ala Leu Leu Ser Ser Ile
 545 550 555
 50 Gln Asn Phe Gln Lys Gly Thr Leu Arg Lys Ala Lys Thr Cys Asp
 560 565 570
 His Ser Ala Pro Lys Ile Gly
 575
 55 <210> 3
 <211> 2119
 <212> DNA
 <213> Homo sapiens
 60 <400> 3
 tgcacctcgg ttctatcgat tcgaattcgg ccacactggc cggatcctct 50
 agagatccct cgacctcgac ccacgcgtcc gctcaaggaa gttgaatgag 100

7

5

10

15

20

25

30

35

40

45

50

5.

6

365

370

5 gta gtg gaa gaa ggg gag gtt ctc tca cac tct cta gaa 1622
Val Val Glu Glu Gly Glu Val Leu Ser His Ser Leu Glu
375 380 385

10 gca gag cac agg tta ttg aaa gct atg ggt tgg cag gaa 1661
Ala Glu His Arg Leu Leu Lys Ala Met Gly Trp Gln Glu
390 395

15 tat cct gaa aat gat gag aat tgc ctt ccc ctc aca gag 1700
Tyr Pro Glu Asn Asp Glu Asn Cys Leu Pro Leu Thr Glu
400 405 410

20 gat gag ctc aaa gag ttc cac atg aag aca gag cag ctg 1739
Asp Glu Leu Lys Glu Phe His Met Lys Thr Glu Gln Leu
415 420 425

25 aga aga aat ggc ttt gga aag aat ggc ttc ttg cag agc 1778
Arg Arg Asn Gly Phe Gly Lys Asn Gly Phe Leu Gln Ser
430 435

30 cgc agt tcc agt ctg ttc tcc cct tgg aga agc act tgc 1817
Arg Ser Ser Ser Leu Phe Ser Pro Trp Arg Ser Thr Cys
440 445 450

35 aaa gca gag ttt gag gac tca gac acc gaa acc agt agc 1856
Lys Ala Glu Phe Glu Asp Ser Asp Thr Glu Thr Ser Ser
455 460

40 agt gaa aca tca gat gac gat gcc tgg aag t agg 1890
Ser Glu Thr Ser Asp Asp Asp Ala Trp Lys
465 470 474

45 catataaatg ctcacagtta aatctgaccc agtaaaactct gtgtgttttag 1940
ggagtataca aaagaaatcg ttcttttcct tttcttatgt tgttgaatac 1990
ttcattcaca agggaaataa tcatatccca aagagagaaa aaaaaaaaaa 2040
40 aaagggcggc cgcgactcta gagtcgacct gcagtaggga taacagggta 2090
ataagcttgg ccgccatggc ccaacttgt 2119

50 <210> 4
<211> 474
<212> PRT
<213> Homo sapiens

55 <400> 4
Met Ala Gln His Asp Phe Val Pro Ala Trp Leu Asn Phe Ser Thr
1 5 10 15
Pro Gln Ser Ala Lys Ser Pro Thr Ala Thr Phe Glu Lys His Gly
20 25 30
Glu His Leu Pro Arg Gly Glu Gly Arg Phe Gly Val Ser Arg Arg
35 40 45
60 Arg His Asn Ser Ser Asp Gly Phe Phe Asn Asn Gly Pro Leu Arg
50 55 60
Thr Ala Gly Asp Ser Trp His Gln Pro Ser Cys Ser Ala Met Ile
65 70 75

CCCT "AGT" 1000

	Leu	Trp	Thr	Leu	Val	Ser	Leu	Arg	Glu	His	Met	Leu	Glu	Ser	Gln	
					80					85					90	
5	Gly	Thr	His	Leu	Val	Gly	Ile	Ala	Leu	Pro	Arg	Gly	His	Asp	Gly	
					95					100					105	
	Met	Ser	Gln	Arg	Ser	Val	Gly	Gly	Thr	Gly	Asn	His	Arg	His	Trp	
					110					115					120	
10	Asn	Gly	Ser	Phe	His	Ser	Arg	Lys	Gly	Cys	Ala	Phe	Gln	Glu	Lys	
					125					130					135	
	Pro	Pro	Met	Glu	Ile	Arg	Glu	Glu	Lys	Lys	Glu	Asp	Lys	Val	Glu	
15					140					145					150	
	Lys	Leu	Gln	Phe	Glu	Glu	Glu	Asp	Phe	Pro	Ser	Leu	Asn	Pro	Glu	
					155					160					165	
	Ala	Gly	Lys	Gln	His	Gln	Pro	Cys	Arg	Pro	Ile	Gly	Thr	Pro	Ser	
20					170					175					180	
	Gly	Val	Trp	Glu	Asn	Pro	Pro	Ser	Ala	Lys	Gln	Pro	Ser	Lys	Met	
					185					190					195	
25	Leu	Val	Ile	Lys	Lys	Val	Ser	Lys	Glu	Asp	Pro	Ala	Ala	Ala	Phe	
					200					205					210	
	Ser	Ala	Ala	Phe	Thr	Ser	Pro	Gly	Ser	His	His	Ala	Asn	Gly	Asn	
30					215					220					225	
	Lys	Leu	Ser	Ser	Val	Val	Pro	Ser	Val	Tyr	Lys	Asn	Leu	Val	Pro	
					230					235					240	
	Lys	Pro	Val	Pro	Pro	Pro	Ser	Lys	Pro	Asn	Ala	Trp	Lys	Ala	Asn	
35					245					250					255	
	Arg	Met	Glu	His	Lys	Ser	Gly	Ser	Leu	Ser	Ser	Ser	Arg	Glu	Ser	
					260					265					270	
40	Ala	Phe	Thr	Ser	Pro	Ile	Ser	Val	Thr	Lys	Pro	Val	Val	Leu	Ala	
					275					280					285	
	Ser	Gly	Ala	Ala	Leu	Ser	Ser	Pro	Lys	Glu	Ser	Pro	Ser	Ser	Thr	
45					290					295					300	
	Thr	Pro	Pro	Ile	Glu	Ile	Ser	Ser	Ser	Arg	Leu	Thr	Lys	Leu	Thr	
					305					310					315	
	Arg	Arg	Thr	Thr	Asp	Arg	Lys	Ser	Glu	Phe	Leu	Lys	Thr	Leu	Lys	
50					320					325					330	
	Asp	Asp	Arg	Asn	Gly	Asp	Phe	Ser	Glu	Asn	Arg	Asp	Cys	Asp	Lys	
					335					340					345	
55	Leu	Glu	Asp	Leu	Glu	Asp	Asn	Ser	Thr	Pro	Glu	Pro	Lys	Glu	Asn	
					350					355					360	
	Gly	Glu	Glu	Gly	Cys	His	Gln	Asn	Gly	Leu	Ala	Leu	Pro	Val	Val	
					365					370					375	
60	Glu	Glu	Gly	Glu	Val	Leu	Ser	His	Ser	Leu	Glu	Ala	Glu	His	Arg	
					380					385					390	
	Leu	Leu	Lys	Ala	Met	Gly	Trp	Gln	Glu	Tyr	Pro	Glu	Asn	Asp	Glu	

```

                                395                                400                                405
Asn Cys Leu Pro Leu Thr Glu Asp Glu Leu Lys Glu Phe His Met
                                410                                415                                420
5
Lys Thr Glu Gln Leu Arg Arg Asn Gly Phe Gly Lys Asn Gly Phe
                                425                                430                                435
Leu Gln Ser Arg Ser Ser Ser Leu Phe Ser Pro Trp Arg Ser Thr
10
                                440                                445                                450
Cys Lys Ala Glu Phe Glu Asp Ser Asp Thr Glu Thr Ser Ser Ser
                                455                                460                                465
15
Glu Thr Ser Asp Asp Asp Ala Trp Lys
                                470

<210> 5
<211> 294
20
<212> DNA
<213> Homo sapiens

<220>
<221> unsure
25
<222> 162, 165, 183-184, 187, 204, 207, 211, 216, 219, 222-223, 225,
237, 240, 244, 250, 253, 262, 267, 269, 275, 279, 282, 289-290
<223> unknown base

<400> 5
30
cccgggatgg ccttcatgga gaagccgcca gctggcaagg tgctgctgga 50
cgacacggtg ccgctgacag cagccatcga ggcgagccag agcctgcagt 100
ccacacgga atatattatt cgagtgcaaa gaggaatttc tctggaaaac 150
35
agctggcaga tngtnagaag atacagtgac tttnatntgc ttaacaacag 200
cttnanatt ncaggncctna gnntncctct tcttcnnaan aaantgattn 250
40
ggnaacatgg ancgtgnant tcatngctng anaggcagnn aggt 294

<210> 6
<211> 24
45
<212> DNA
<213> Homo sapiens

<400> 6
ggacgacacg gtgccgctga cagc 24

50
<210> 7
<211> 51
<212> DNA
<213> Homo sapiens

55
<400> 7
gccatcgagg cgagccagag cctgcagtcc cacacggaat atattattcg 50
a 51

60
<210> 8
<211> 31
<212> DNA
<213> Homo sapiens

```

<400> 8
 gttttccaga gaaattcctc ttgactcg a 31

5 <210> 9
 <211> 793
 <212> DNA
 <213> Homo sapiens

10 <220>
 <221> unsure
 <222> 379, 433, 459, 492, 517, 541, 549, 561, 575, 579, 582, 710,
 742,
 774, 784
 <223> unknown base

15 <400> 9
 gggaagaaaa gaaagaagac aaggtggaaa agttgcagat ttgaagagga 50
 ggacttttct tccttgaatc cagaagctgg caaacagcat cagccatgca 100
 20 gacctattgg gacaccttct ggagtatggg aaaacccgcc tagtgccaag 150
 caaccctcca agatgctagt tatcaaaaaa gtttccaaag aggatcctgc 200
 25 tgctgcyttc tctgctgcat tcacctcacc aggatctcac catgcaaattg 250
 ggaacaaatt gtcattcgtg gttccaagtg totataagaa cctgggttct 300
 aagcctgtac cacctccttc caagcctaatt gcatggaaag ctaacaggat 350
 30 ggagcacaag tcaggatccc tttcctctna gccgggagtc tgcttttacc 400
 agtccaatct ctgttaccac accagtggta ctnggctagt ggtgcagctc 450
 35 tgagttctnc ccaaagagag tccctccagc accacccctc cnaattgaga 500
 tcagctcttc tcgtctngac caagtttgac ccgccgaacc naccgacang 550
 gaagragtga ngttcctgaa aactnctgna anggatgacc gggaatggga 600
 40 agacttitytc agaagaatag agactgtgac aagctggaag atttggagga 650
 caacagcaca ctgaacccaaa ggaaaatggg ggaggaaggc tgttcatcaa 700
 45 atggtctttn ctttctgtga gttggaagga ggggaggttt tntcacattt 750
 ttaggagcag agcacaggtt tttnaagttt tggntggcgg att 793

50 <210> 10
 <211> 24
 <212> DNA
 <213> Homo sapiens

55 <400> 10
 gacctattgg gacaccttct ggag 24

60 <210> 11
 <211> 43
 <212> DNA
 <213> Homo sapiens

<400> 11
 cccgcctagt gccaaagcaac cctccaagat gctagttatc aaa 43

<210> 12
 <211> 25
 <212> DNA
 <213> Homo sapiens

5

<400> 12
 cttggtcaga cgagaggagc tgatc 25

10

<210> 13
 <211> 2366
 <212> DNA
 <213> Homo sapiens

15

<220>
 <221> unsure
 <222> 103
 <223> unknown base

20

<400> 13
 tggccggatc ctctagagat ccctcgacct cgacccacgc gtccgcagaa 50

agtatgtccc ttctctcacc atg agc tgg ctc tcc agt tcc 91
 Met Ser Trp Leu Ser Ser Ser
 1 5

25

cag gga gtg gtn cta aca gcc tac cac ccc agc ggc aag 130
 Gln Gly Val Xaa Leu Thr Ala Tyr His Pro Ser Gly Lys
 10 15 20

30

gac cag acc gtc ggg aac agc cat gca aag gca ggg gag 169
 Asp Gln Thr Val Gly Asn Ser His Ala Lys Ala Gly Glu
 25 30

35

gaa gcc acc tcg agt cgc aga tat ggc cag tac act atg 208
 Glu Ala Thr Ser Ser Arg Arg Tyr Gly Gln Tyr Thr Met
 35 40 45

40

aac cag gaa agc acc acc atc aaa gtt atg gag aag cct 247
 Asn Gln Glu Ser Thr Thr Ile Lys Val Met Glu Lys Pro
 50 55

45

cca ttt gat cga tca att tcc cag gat tct ttg gat gaa 286
 Pro Phe Asp Arg Ser Ile Ser Gln Asp Ser Leu Asp Glu
 60 65 70

50

cta tct atg gaa gac tat tgg ata gaa cta gaa aac atc 325
 Leu Ser Met Glu Asp Tyr Trp Ile Glu Leu Glu Asn Ile
 75 80 85

55

aag aaa tct agt gaa aac agc caa gaa gat caa gag gtg 364
 Lys Lys Ser Ser Glu Asn Ser Gln Glu Asp Gln Glu Val
 90 95

gtt gtt gtc aaa gag cct gat gag gga gaa ttg gaa gaa 403
 Val Val Val Lys Glu Pro Asp Glu Gly Glu Leu Glu Glu
 100 105 110

60

gag tgg ctt aaa gag gcc ggt tta tcc aat ctc ttc gga 442
 Glu Trp Leu Lys Glu Ala Gly Leu Ser Asn Leu Phe Gly
 115 120

gag tct gct gga gat cca cag gaa agc att gtg ttt tta 481
 Glu Ser Ala Gly Asp Pro Gln Glu Ser Ile Val Phe Leu
 125 130 135

tca aca ttg acg cgg acc cag gca gca gca gtt cag aag 520
 Ser Thr Leu Thr Arg Thr Gln Ala Ala Val Gln Lys
 140 145 150

5 cga gta gag acg gtc tcc cag acc ttg agg aaa aaa aac 559
 Arg Val Glu Thr Val Ser Gln Thr Leu Arg Lys Lys Asn
 155 160

10 aaa cag tac cag att cct gac gtc aga gac ata ttt gct 598
 Lys Gln Tyr Gln Ile Pro Asp Val Arg Asp Ile Phe Ala
 165 170 175

15 caa cag aga gaa tca aaa gaa aca gct cca ggt ggc act 637
 Gln Gln Arg Glu Ser Lys Glu Thr Ala Pro Gly Gly Thr
 180 185

20 gaa tcg cag tca ctt aga aca aat gaa aac aaa tac caa 676
 Glu Ser Gln Ser Leu Arg Thr Asn Glu Asn Lys Tyr Gln
 190 195 200

25 gga aga gat gac gag gca tct aac ctt gtt ggt gaa gag 715
 Gly Arg Asp Asp Glu Ala Ser Asn Leu Val Gly Glu Glu
 205 210 215

30 aag ctg atc cca cct gag gag acg cct gcc cct gaa aca 754
 Lys Leu Ile Pro Pro Glu Glu Thr Pro Ala Pro Glu Thr
 220 225

35 gac atc aac ctg gag gta tca ttt gcc gag caa gca ctc 793
 Asp Ile Asn Leu Glu Val Ser Phe Ala Glu Gln Ala Leu
 230 235 240

40 aat cag aaa gag aga tcc aag gag aaa atc cag aag agc 832
 Asn Gln Lys Glu Arg Ser Lys Glu Lys Ile Gln Lys Ser
 245 250

45 aaa ggc gat gat gcc aca tta cct agt ttc aga ttg cca 871
 Lys Gly Asp Asp Ala Thr Leu Pro Ser Phe Arg Leu Pro
 255 260 265

50 aaa gac aaa acg ggt acc aca agg att ggt gac ctc gca 910
 Lys Asp Lys Thr Gly Thr Thr Arg Ile Gly Asp Leu Ala
 270 275 280

55 ccc cag gac atg aag aaa gtt tgc cat tta gcc cta att 949
 Pro Gln Asp Met Lys Lys Val Cys His Leu Ala Leu Ile
 285 290

60 gag ctg act gcc ctc tat gat gta ttg ggt att gag ctg 988
 Glu Leu Thr Ala Leu Tyr Asp Val Leu Gly Ile Glu Leu
 295 300 305

65 aaa caa caa aaa gct gtg aaa atc aaa aca aaa gat tct 1027
 Lys Gln Gln Lys Ala Val Lys Ile Lys Thr Lys Asp Ser
 310 315

70 ggt ctt ttt tgc gtt cca ttg aca gcg cta tta gaa caa 1066
 Gly Leu Phe Cys Val Pro Leu Thr Ala Leu Leu Glu Gln
 320 325 330

75 gat cag agg aaa gta cca gga atg cga ata ccc ttg atc 1105
 Asp Gln Arg Lys Val Pro Gly Met Arg Ile Pro Leu Ile
 335 340 345

ttt caa aaa ctg att tct cga att gaa gag aga ggt ttg 1144
 Phe Gln Lys Leu Ile Ser Arg Ile Glu Glu Arg Gly Leu
 350 355

5 gaa aca gaa ggc ctc tta cgg atc cct gga gct gcc att 1183
 Glu Thr Glu Gly Leu Leu Arg Ile Pro Gly Ala Ala Ile
 360 365 370

10 aga atc aag aat ctt tgc caa gaa cta gaa gca aag ttt 1222
 Arg Ile Lys Asn Leu Cys Gln Glu Leu Glu Ala Lys Phe
 375 380

15 tat gaa ggg act ttt aat tgg gaa agt gtc aaa cag cat 1261
 Tyr Glu Gly Thr Phe Asn Trp Glu Ser Val Lys Gln His
 385 390 395

20 gat gcc gcc agc ctg ctg aag ctc ttc att cgg gag ttg 1300
 Asp Ala Ala Ser Leu Leu Lys Leu Phe Ile Arg Glu Leu
 400 405 410

25 ccc cag cca ctg ctc agt gtg gag tat ctc aaa gcc ttt 1339
 Pro Gln Pro Leu Leu Ser Val Glu Tyr Leu Lys Ala Phe
 415 420

30 cag gct ttg aac ctt ctt gtc atc ctc cta cct gat gca 1417
 Gln Ala Leu Asn Leu Leu Val Ile Leu Leu Pro Asp Ala
 440 445

35 aac agg gac aca ctg aag gcc ctt ctt gaa ttt ctc caa 1456
 Asn Arg Asp Thr Leu Lys Ala Leu Leu Glu Phe Leu Gln
 450 455 460

40 aga gta ata gat aat aaa gaa aaa aat aaa atg aca gtc 1495
 Arg Val Ile Asp Asn Lys Glu Lys Asn Lys Met Thr Val
 465 470 475

45 atg aat gta gca atg gtc atg gcc ccg aat ctc ttt atg 1534
 Met Asn Val Ala Met Val Met Ala Pro Asn Leu Phe Met
 480 485

50 tgt cat gca ttg gga ttg aag tcc agt gaa cag cga gaa 1573
 Cys His Ala Leu Gly Leu Lys Ser Ser Glu Gln Arg Glu
 490 495 500

55 ttt gta atg gca gct ggg aca gca aat acc atg cac tta 1612
 Phe Val Met Ala Ala Gly Thr Ala Asn Thr Met His Leu
 505 510

60 ttg att aag tac caa aaa ctt ctg tgg aca att ccc aag 1651
 Leu Ile Lys Tyr Gln Lys Leu Leu Trp Thr Ile Pro Lys
 515 520 525

ttt att gta aac caa gtg agg aag caa aac acg gaa aat 1690
 Phe Ile Val Asn Gln Val Arg Lys Gln Asn Thr Glu Asn
 530 535 540

cat aaa aag gat aaa aga gcc atg aag aaa ttg ctg aag 1729
 His Lys Lys Asp Lys Arg Ala Met Lys Lys Leu Leu Lys
 545 550

aaa atg gct tat gac cga gaa aaa tat gaa aag caa gat 1768
 Lys Met Ala Tyr Asp Arg Glu Lys Tyr Glu Lys Gln Asp
 555 560 565

5 aag agt aca aat gat gct gac gtt cct cag gga gtg att 1807
 Lys Ser Thr Asn Asp Ala Asp Val Pro Gln Gly Val Ile
 570 575

10 cga gtg caa gct ccc cat ctt tcg aaa gtt tcc atg gca 1846
 Arg Val Gln Ala Pro His Leu Ser Lys Val Ser Met Ala
 580 585 590

15 ata cag cta act gaa gaa cta aaa gcc agt gat gta ctt 1885
 Ile Gln Leu Thr Glu Glu Leu Lys Ala Ser Asp Val Leu
 595 600 605

20 gcc agg ttt ctc agc caa gaa agt ggg gtt gcc cag act 1924
 Ala Arg Phe Leu Ser Gln Glu Ser Gly Val Ala Gln Thr
 610 615

25 ctc aag aaa gga gaa gtt ttt ttg tat gaa att gga gga 1963
 Leu Lys Lys Gly Glu Val Phe Leu Tyr Glu Ile Gly Gly
 620 625 630

30 aat att ggg gaa cgc tgc ctt gat gat gac act tac atg 2002
 Asn Ile Gly Glu Arg Cys Leu Asp Asp Asp Thr Tyr Met
 635 640

35 aag gat tta tat cag ctt aac cca aat gct gag tgg gtt 2041
 Lys Asp Leu Tyr Gln Leu Asn Pro Asn Ala Glu Trp Val
 645 650 655

40 ata aag tca aag cca ttg t agaagactta acaagctgca 2080
 Ile Lys Ser Lys Pro Leu
 660 663

45 gataaccatg tggactttctg tcataattct tgctgagtca agagtgtaaa 2130
 taaaagaaat ggcaggactc atattattca gttgtacca agtatttttaa 2180
 aaatgactct ottaagcctt aaaaagtcac agattttgtgc tgctgccaga 2230
 attatattaa attattatta atgggtattat tagaaaaaaaa aattttctgga 2280
 gtgagagtaa agaggcttaa ttagttttgtg ggcagttttc ttatgctctg 2330
 tgaaatgtgt ccagatgtga catagttttt tttttt 2366

50 <210> 14
 <211> 663
 <212> PRT
 <213> Homo sapiens

55 <400> 14
 Met Ser Trp Leu Ser Ser Ser Gln Gly Val Val Leu Thr Ala Tyr
 1 5 10 15
 His Pro Ser Gly Lys Asp Gln Thr Val Gly Asn Ser His Ala Lys
 20 25 30
 60 Ala Gly Glu Glu Ala Thr Ser Ser Arg Arg Tyr Gly Gln Tyr Thr
 35 40 45
 Met Asn Gln Glu Ser Thr Thr Ile Lys Val Met Glu Lys Pro Pro

00645-10650
 00600-10650
 00550-10650
 00500-10650
 00450-10650
 00400-10650
 00350-10650
 00300-10650
 00250-10650
 00200-10650
 00150-10650
 00100-10650
 00050-10650
 00000-10650

	50	55	60
	Phe Asp Arg Ser Ile Ser Gln Asp Ser Leu Asp Glu Leu Ser Met		
	65	70	75
5	Glu Asp Tyr Trp Ile Glu Leu Glu Asn Ile Lys Lys Ser Ser Glu		
	80	85	90
	Asn Ser Gln Glu Asp Gln Glu Val Val Val Val Lys Glu Pro Asp		
10	95	100	105
	Glu Gly Glu Leu Glu Glu Glu Trp Leu Lys Glu Ala Gly Leu Ser		
	110	115	120
15	Asn Leu Phe Gly Glu Ser Ala Gly Asp Pro Gln Glu Ser Ile Val		
	125	130	135
	Phe Leu Ser Thr Leu Thr Arg Thr Gln Ala Ala Ala Val Gln Lys		
20	140	145	150
	Arg Val Glu Thr Val Ser Gln Thr Leu Arg Lys Lys Asn Lys Gln		
	155	160	165
25	Tyr Gln Ile Pro Asp Val Arg Asp Ile Phe Ala Gln Gln Arg Glu		
	170	175	180
	Ser Lys Glu Thr Ala Pro Gly Gly Thr Glu Ser Gln Ser Leu Arg		
	185	190	195
30	Thr Asn Glu Asn Lys Tyr Gln Gly Arg Asp Asp Glu Ala Ser Asn		
	200	205	210
	Leu Val Gly Glu Glu Lys Leu Ile Pro Pro Glu Glu Thr Pro Ala		
35	215	220	225
	Pro Glu Thr Asp Ile Asn Leu Glu Val Ser Phe Ala Glu Gln Ala		
	230	235	240
40	Leu Asn Gln Lys Glu Arg Ser Lys Glu Lys Ile Gln Lys Ser Lys		
	245	250	255
	Gly Asp Asp Ala Thr Leu Pro Ser Phe Arg Leu Pro Lys Asp Lys		
	260	265	270
45	Thr Gly Thr Thr Arg Ile Gly Asp Leu Ala Pro Gln Asp Met Lys		
	275	280	285
	Lys Val Cys His Leu Ala Leu Ile Glu Leu Thr Ala Leu Tyr Asp		
50	290	295	300
	Val Leu Gly Ile Glu Leu Lys Gln Gln Lys Ala Val Lys Ile Lys		
	305	310	315
55	Thr Lys Asp Ser Gly Leu Phe Cys Val Pro Leu Thr Ala Leu Leu		
	320	325	330
	Glu Gln Asp Gln Arg Lys Val Pro Gly Met Arg Ile Pro Leu Ile		
	335	340	345
60	Phe Gln Lys Leu Ile Ser Arg Ile Glu Glu Arg Gly Leu Glu Thr		
	350	355	360
	Glu Gly Leu Leu Arg Ile Pro Gly Ala Ala Ile Arg Ile Lys Asn		
	365	370	375

60

5

10

15

20

25

30

35

40

45

50

55

60

ggg gac ctc tac ctc acc gag ggc gtg cgg cat ggc gcc 719
 Gly Asp Leu Tyr Leu Thr Glu Gly Val Pro His Gly Ala
 140 145 150

tgg cac gcg gcc gcc ctc cca ccc ctg ttg cgc cca ctg 758
 Trp His Ala Ala Ala Leu Pro Pro Leu Leu Arg Pro Leu
 155 160

5 ctg ccg cct gcc ctg cag ggt gct ctc cag cag tgg ctg 797
 Leu Pro Pro Ala Leu Gln Gly Ala Leu Gln Gln Trp Leu
 165 170 175

10 ggg cct gcg tgg cct tgg cgg gcc aag atc gcc atc ggc 836
 Gly Pro Ala Trp Pro Trp Arg Ala Lys Ile Ala Ile Gly
 180 185 190

15 ctg ctg gag ttc gtg gag gag ctc ttc cac ggc tct tac 875
 Leu Leu Glu Phe Val Glu Glu Leu Phe His Gly Ser Tyr
 195 200

20 ggg act ttc tac atg tgt gag acc aca ctg gcc aac gtg 914
 Gly Thr Phe Tyr Met Cys Glu Thr Thr Leu Ala Asn Val
 205 210 215

ggc tac aca gcc acc tac gac ttc aag atg gcc gac ctg 953
 Gly Tyr Thr Ala Thr Tyr Asp Phe Lys Met Ala Asp Leu
 220 225

25 cag cag gtg gca ccc gag gcc acc gtg cgc cgc ttc ctg 992
 Gln Gln Val Ala Pro Glu Ala Thr Val Arg Arg Phe Leu
 230 235 240

30 cag ggc cgc cgc tgc gag cac agc acc gac tgc acc tac 1031
 Gln Gly Arg Arg Cys Glu His Ser Thr Asp Cys Thr Tyr
 245 250 255

35 ggg cgc gac tgc agg gcc ccg tgt gac agg ctc atg agg 1070
 Gly Arg Asp Cys Arg Ala Pro Cys Asp Arg Leu Met Arg
 260 265

40 cag tgc aag ggc gac ctc atc cag ccc aac ctg gcc aag 1109
 Gln Cys Lys Gly Asp Leu Ile Gln Pro Asn Leu Ala Lys
 270 275 280

gtg tgc gca ctg cta cgg ggc tac ctg ctg cct ggc gcg 1148
 Val Cys Ala Leu Leu Arg Gly Tyr Leu Leu Pro Gly Ala
 285 290

45 ccc gcc gac ctc cgc gag gag ctg ggc aca cag ctg cgc 1187
 Pro Ala Asp Leu Arg Glu Glu Leu Gly Thr Gln Leu Arg
 295 300 305

50 acc tgt acc acg ctg agc ggg ctg gcc agc cag gtg gag 1226
 Thr Cys Thr Thr Leu Ser Gly Leu Ala Ser Gln Val Glu
 310 315 320

55 gcc cat cac tcg ctg gtg ctc agc cac ctc aag act ctg 1265
 Ala His His Ser Leu Val Leu Ser His Leu Lys Thr Leu
 325 330

60 ctc tgg aag aag atc tcc aac acc aag tac tct t g 1300
 Leu Trp Lys Lys Ile Ser Asn Thr Lys Tyr Ser
 335 340 344

atgggnaatg agggcttgca accttct 1327
 .
 <210> 16
 <211> 344

005001-00500

<212> PRT
 <213> Homo sapiens

<220>
 <221> unsure
 <222> 104
 <223> unknown amino acid

<400> 16

10	Met	Val	Glu	Trp	Arg	Thr	Cys	Leu	Ser	Val	Ala	Pro	Gly	Gln	Gln	1	5	10	15
	Val	Tyr	Ser	Gly	Leu	Trp	Arg	Asp	Lys	Asp	Val	Thr	Ile	Lys	Cys	20	25	30	
15	Gly	Ile	Glu	Glu	Thr	Leu	Asp	Ser	Lys	Ala	Arg	Ser	Asp	Ala	Ala	35	40	45	
20	Pro	Arg	Arg	Glu	Leu	Val	Leu	Phe	Asp	Lys	Pro	Thr	Arg	Gly	Thr	50	55	60	
	Ser	Ile	Lys	Glu	Phe	Arg	Glu	Met	Thr	Leu	Gly	Phe	Leu	Lys	Ala	65	70	75	
25	Asn	Leu	Gly	Asp	Leu	Pro	Ser	Leu	Pro	Ala	Leu	Val	Gly	Gln	Val	80	85	90	
	Leu	Leu	Met	Ala	Asp	Phe	Asn	Lys	Asp	Asn	Arg	Val	Ser	Xaa	Ala	95	100	105	
30	Glu	Ala	Lys	Ser	Val	Trp	Ala	Leu	Leu	Gln	Arg	Asn	Glu	Phe	Leu	110	115	120	
35	Leu	Leu	Leu	Ser	Leu	Gln	Glu	Lys	Glu	His	Ala	Ser	Arg	Leu	Leu	125	130	135	
	Gly	Tyr	Cys	Gly	Asp	Leu	Tyr	Leu	Thr	Glu	Gly	Val	Pro	His	Gly	140	145	150	
40	Ala	Trp	His	Ala	Ala	Ala	Leu	Pro	Pro	Leu	Leu	Arg	Pro	Leu	Leu	155	160	165	
	Pro	Pro	Ala	Leu	Gln	Gly	Ala	Leu	Gln	Gln	Trp	Leu	Gly	Pro	Ala	170	175	180	
45	Trp	Pro	Trp	Arg	Ala	Lys	Ile	Ala	Ile	Gly	Leu	Leu	Glu	Phe	Val	185	190	195	
50	Glu	Glu	Leu	Phe	His	Gly	Ser	Tyr	Gly	Thr	Phe	Tyr	Met	Cys	Glu	200	205	210	
	Thr	Thr	Leu	Ala	Asn	Val	Gly	Tyr	Thr	Ala	Thr	Tyr	Asp	Phe	Lys	215	220	225	
55	Met	Ala	Asp	Leu	Gln	Gln	Val	Ala	Pro	Glu	Ala	Thr	Val	Arg	Arg	230	235	240	
	Phe	Leu	Gln	Gly	Arg	Arg	Cys	Glu	His	Ser	Thr	Asp	Cys	Thr	Tyr	245	250	255	
60	Gly	Arg	Asp	Cys	Arg	Ala	Pro	Cys	Asp	Arg	Leu	Met	Arg	Gln	Cys	260	265	270	
	Lys	Gly	Asp	Leu	Ile	Gln	Pro	Asn	Leu	Ala	Lys	Val	Cys	Ala	Leu				

22

6

```

ctc tca cac ttg acc aag aat gtt cca ata ttt gtt tgc 1188
Leu Ser His Leu Thr Lys Asn Val Pro Ile Phe Val Cys
295 300 305

5 act atg gcc tac ccc act gtg cct tgc cct ctc cat gta 1227
Thr Met Ala Tyr Pro Thr Val Pro Cys Pro Leu His Val
310 315

10 ttt gag cca aga tac aga ttg atg att cga aga agt ata 1266
Phe Glu Pro Arg Tyr Arg Leu Met Ile Arg Arg Ser Ile
320 325 330

15 cag act gga acc aaa cag ttt ggc atg tgt gtc agt gat 1305
Gln Thr Gly Thr Lys Gln Phe Gly Met Cys Val Ser Asp
335 340 345

20 aca caa aat agt ttt gca gat tat ggt tgt atg tta caa 1344
Thr Gln Asn Ser Phe Ala Asp Tyr Gly Cys Met Leu Gln
350 355

25 att aga aac gtg cat ttc tta ccg gac gga agg tct gtg 1383
Ile Arg Asn Val His Phe Leu Pro Asp Gly Arg Ser Val
360 365 370

30 gtt gat aca gtt gga gga aag cgg ttt agg gtt tta aaa 1422
Val Asp Thr Val Gly Gly Lys Arg Phe Arg Val Leu Lys
375 380

35 aga gga atg aaa gat gga tat tgc act gcc gac att gaa 1461
Arg Gly Met Lys Asp Gly Tyr Cys Thr Ala Asp Ile Glu
385 390 395

40 tat ctg gaa gat gtt aag gtt gag aat gaa gat gag att 1500
Tyr Leu Glu Asp Val Lys Val Glu Asn Glu Asp Glu Ile
400 405 410

45 aag aat ctc aga gag ctt cat gat ttg gtt tac tct caa 1539
Lys Asn Leu Arg Glu Leu His Asp Leu Val Tyr Ser Gln
415 420

50 gcc tgc agc tgg ttt cag aat tta aga gac aga ttt cga 1578
Ala Cys Ser Trp Phe Gln Asn Leu Arg Asp Arg Phe Arg
425 430 435

55 agc caa att ctt cag cat ttc gga tca atg ccc gag ang 1617
Ser Gln Ile Leu Gln His Phe Gly Ser Met Pro Glu Xaa
440 445

60 gag gaa aac ctt cag gca gcc cct aat gga cct gca tgg 1656
Glu Glu Asn Leu Gln Ala Ala Pro Asn Gly Pro Ala Trp
450 455 460

tgt tgg tgg ctt ctt gca gtt ctc cct gta gac cca cga 1695
Cys Trp Trp Leu Leu Ala Val Leu Pro Val Asp Pro Arg
465 470 475

tac cag ctg tcg gtt ttg tca atg aag tct ttg aaa gaa 1734
Tyr Gln Leu Ser Val Leu Ser Met Lys Ser Leu Lys Glu
480 485

cgg ttg acc aag ata cag cat ata ctg acc tat ttt tct 1773
Arg Leu Thr Lys Ile Gln His Ile Leu Thr Tyr Phe Ser
490 495 500

```


[illegible]

25

<400> 18

	Met	Gln	Phe	Phe	Phe	Asn	Phe	Gln	Asp	Trp	Pro	Glu	Gly	Leu	Leu	
	1				5					10					15	
5	Gln	Glu	Xaa	Glu	Lys	Val	Leu	Cys	Asp	Ala	Gly	Phe	Leu	Gly	Asp	
					20					25					30	
	Ala	Leu	Gln	Leu	Phe	Leu	Gln	Cys	Leu	Ala	Leu	Asp	Glu	Asp	Phe	
					35					40					45	
10	Ala	Pro	Ala	Lys	Leu	Gln	Val	Gln	Lys	Ile	Leu	Cys	Asp	Leu	Leu	
					50					55					60	
	Leu	Pro	Glu	Asn	Leu	Lys	Glu	Gly	Leu	Lys	Glu	Ser	Ser	Trp	Ser	
15					65					70					75	
	Ser	Leu	Pro	Cys	Thr	Lys	Asn	Arg	Pro	Phe	Asp	Phe	His	Ser	Val	
					80					85					90	
20	Met	Glu	Glu	Ser	Gln	Ser	Leu	Asn	Glu	Pro	Ser	Pro	Lys	Gln	Ser	
					95					100					105	
	Glu	Glu	Ile	Pro	Glu	Val	Thr	Ser	Glu	Pro	Val	Lys	Gly	Ser	Leu	
25					110					115					120	
	Asn	Arg	Ala	Gln	Ser	Ala	Gln	Ser	Ile	Asn	Ser	Thr	Glu	Met	Pro	
					125					130					135	
30	Ala	Arg	Glu	Asp	Cys	Leu	Lys	Lys	Val	Ser	Ser	Glu	Pro	Val	Leu	
					140					145					150	
	Ser	Val	Gln	Glu	Lys	Gly	Val	Leu	Leu	Lys	Arg	Lys	Leu	Ser	Leu	
					155					160					165	
35	Leu	Glu	Gln	Asp	Val	Ile	Val	Asn	Glu	Asp	Gly	Arg	Asn	Lys	Leu	
					170					175					180	
	Lys	Lys	Gln	Gly	Glu	Thr	Pro	Asn	Glu	Val	Cys	Met	Phe	Ser	Leu	
40					185					190					195	
	Ala	Tyr	Gly	Asp	Ile	Pro	Glu	Glu	Leu	Ile	Asp	Val	Ser	Asp	Phe	
					200					205					210	
45	Glu	Cys	Ser	Leu	Cys	Met	Arg	Leu	Phe	Phe	Glu	Pro	Val	Thr	Thr	
					215					220					225	
	Pro	Cys	Gly	His	Ser	Phe	Cys	Lys	Asn	Cys	Leu	Glu	Arg	Cys	Leu	
					230					235					240	
50	Asp	His	Ala	Pro	Tyr	Cys	Pro	Leu	Cys	Lys	Glu	Ser	Leu	Asn	Glu	
					245					250					255	
	Tyr	Leu	Ala	Asp	Arg	Arg	Tyr	Cys	Val	Thr	Gln	Leu	Leu	Glu	Glu	
					260					265					270	
55	Leu	Ile	Val	Lys	Tyr	Leu	Pro	Asp	Glu	Leu	Ser	Glu	Arg	Lys	Lys	
					275					280					285	
	Ile	Tyr	Glu	Glu	Glu	Thr	Ala	Glu	Leu	Ser	His	Leu	Thr	Lys	Asn	
60					290					295					300	
	Val	Pro	Ile	Phe	Val	Cys	Thr	Met	Ala	Tyr	Pro	Thr	Val	Pro	Cys	
					305					310					315	

005001 00440050

40
45
50
55
60

5

10

15

20

25

30

35

40

45

50

54

6

<212> DNA
 <213> Homo sapiens

<400> 25
 5 agcagcagca ggaactcgtt acgctgcagc agggcccaca cggacttggc 50
 ttccgcccag gacacccggt tgtccttggt gaagtcagcc atgagcagga 100
 cctggccaac cagcgcgggc aggggaaggca ggtctcccag gttcgccttg 150
 10 aggaagccga gggtcattct cccgaattcc ttgatggagg tgccccgggt 200
 gggcttgtca aacagtacca gctcccgcg ggggccgcat ccgaccgggc 250
 15 cttggagtcg aggggtctct caatgccaca cttgatgggt acatccttgt 300
 cccgccagag cccgtgtac acctgctggc ccggggcaca cgagaagcag 350
 gtctccact ccaccatatg cagctcacac aggtcctggc agacggagcc 400
 20 cgagatgata cccttgcggt actggtcaca aat 433

<210> 26
 <211> 22
 <212> DNA
 <213> Homo sapiens

<400> 26
 atgcagttct ttttcaactt cc 22

<210> 27
 <211> 20
 <212> DNA
 <213> Homo sapiens

<400> 27
 ctagagacca atctaagtaa 20

<210> 28
 <211> 20
 <212> DNA
 <213> Homo sapiens


<400> 28
 atggtggagt ggaggacctg 20

<210> 29
 <211> 22
 <212> DNA
 <213> Homo sapiens

<400> 29
 ctccaacacc aagtactctt ga 22

55

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

<p>In re Application of Gerritsen <i>et al.</i></p> <p>Serial No.: not assigned</p> <p>Filed: October 5, 2000</p> <p>For: NOVEL POLYPEPTIDES, THEIR NUCLEIC ACIDS, AND METHODS FOR THEIR USE IN ANGIOGENESIS AND VASCULARIZATION</p>	<p>Group Art Unit: unassigned</p> <p>Examiner: unassigned</p> <hr/> <p>CERTIFICATE OF MAILING I hereby certify that this correspondence is being deposited with the United States Postal Service with sufficient postage as first class mail in an envelope addressed to: Director of Patents and Trademarks, Washington, DC 20231 on</p> <p>October 5, 2000</p> <p> Glory L. Tabuena</p>
---	--

CERTIFICATE RE: SEQUENCE LISTING
RESPONSE UNDER 37 CFR § 1.821(f) and (g)


Box Sequence
 Director of Patents and Trademarks
 Washington, DC 20231

Sir:

I hereby state that the Sequence Listing submitted herewith is submitted in paper copy and a computer-readable diskette, and that the information recorded in computer readable form is identical to the written sequence listing. I further state that this submission includes no new matter.

Respectfully submitted,

GENENTECH, INC.

By: 
 Timothy R. Schwartz, Ph.D.
 Reg. No. 32,171

Date: October 5, 2000

1 DNA Way
 South San Francisco, CA 94080-4990
 Phone: (650) 225-7467
 Fax: (650) 952-9881

005001 8544860

Sequence Listing

<110> Gerritsen, Mary E.
 Goddard, Audrey
 Grimaldi, J. Christopher
 Mehraban, Fuad

<120> NOVEL POLYPEPTIDES, THEIR NUCLEIC ACIDS, AND METHODS
 FOR THEIR USE IN ANGIOGENESIS AND VASCULARIZATION

<130> P1776R2US

<141> 2000-10-05

<150> US 60/158,587
 <151> 1999-10-07

<150> US 60/162,611
 <151> 1999-10-28

<160> 29

<210> 1
 <211> 2891
 <212> DNA
 <213> Homo sapiens

<400> 1
 ggcgaggagtc ggcgccctcgg gttcctacct cgcgtcccta ggcggcggcg 50

gcccgggcgtc cccggg atg gcc ttc atg gag aag ccg cca 89
 Met Ala Phe Met Glu Lys Pro Pro
 1 5

gcc ggc aag gtg ctg ctg gac gac acg gtg ccg ctg aca 128
 Ala Gly Lys Val Leu Leu Asp Asp Thr Val Pro Leu Thr
 10 15 20

gca gcc atc gag gcg agc cag agc ctg cag tcc cac acg 167
 Ala Ala Ile Glu Ala Ser Gln Ser Leu Gln Ser His Thr
 25 30

gaa tat att att cga gtg caa aga gga att tct gtg gaa 206
 Glu Tyr Ile Ile Arg Val Gln Arg Gly Ile Ser Val Glu
 35 40 45

aac agc tgg cag att gtt aga aga tac agt gac ttt gat 245
 Asn Ser Trp Gln Ile Val Arg Arg Tyr Ser Asp Phe Asp
 50 55 60

ttg ctt aac aac agc tta cag att gca ggc cta agt cta 284
 Leu Leu Asn Asn Ser Leu Gln Ile Ala Gly Leu Ser Leu
 65 70

cct ctt cct ccc aaa aaa ttg att ggt aac atg gat cgt 323
 Pro Leu Pro Pro Lys Lys Leu Ile Gly Asn Met Asp Arg
 75 80 85

gaa ttc ata gct gaa agg cag aaa ggt ctt cag aac tat 362
 Glu Phe Ile Ala Glu Arg Gln Lys Gly Leu Gln Asn Tyr
 90 95

ctc aac gtg atc aca aca aat cat atc ttg tct aat tgt 401
 Leu Asn Val Ile Thr Thr Asn His Ile Leu Ser Asn Cys
 100 105 110

gag ctg gtt aag aag ttt tta gat cca aac aac tat tcc 440
 Glu Leu Val Lys Lys Phe Leu Asp Pro Asn Asn Tyr Ser
 115 120 125

gca aac tat act gag att gcc ttg caa cag gtt tcc atg 479
 Ala Asn Tyr Thr Glu Ile Ala Leu Gln Gln Val Ser Met
 130 135

ttc ttc cga tca gaa cca aag tgg gag gtg gtg gaa cct 518
 Phe Phe Arg Ser Glu Pro Lys Trp Glu Val Val Glu Pro
 140 145 150

ttg aaa gac ata ggt tgg aga ata agg aag aaa tat ttc 557
 Leu Lys Asp Ile Gly Trp Arg Ile Arg Lys Lys Tyr Phe
 155 160

ttg atg aag att aaa aat cag cca aag gaa cgg cta gtg 596
 Leu Met Lys Ile Lys Asn Gln Pro Lys Glu Arg Leu Val
 165 170 175

tta agc tgg gct gac ctt ggc cca gac aag tat ttg tca 635
 Leu Ser Trp Ala Asp Leu Gly Pro Asp Lys Tyr Leu Ser
 180 185 190

gat aaa gat ttt cag tgt cta atc aaa ctt ctg cct tct 674
 Asp Lys Asp Phe Gln Cys Leu Ile Lys Leu Leu Pro Ser
 195 200

tgt ttg cac cct tac atc tat cgg gtt acc ttt gcc aca 713
 Cys Leu His Pro Tyr Ile Tyr Arg Val Thr Phe Ala Thr
 205 210 215

gct aat gaa tcc tca gcg ttg cta att agg atg ttt aac 752
 Ala Asn Glu Ser Ser Ala Leu Leu Ile Arg Met Phe Asn
 220 225

gaa aag gga aca ttg aag gat ctg atc tac aag gca aaa 791
 Glu Lys Gly Thr Leu Lys Asp Leu Ile Tyr Lys Ala Lys
 230 235 240

cca aaa gac cca ttt cta aag aag tac tgc aac cct aag 830
 Pro Lys Asp Pro Phe Leu Lys Lys Tyr Cys Asn Pro Lys
 245 250 255

aag att cag ggc ctg gaa ctc cag caa ata aaa aca tat 869
 Lys Ile Gln Gly Leu Glu Leu Gln Gln Ile Lys Thr Tyr
 260 265

gga cgg caa ata tta gag gta ctg aag ttt ctt cat gac 908
 Gly Arg Gln Ile Leu Glu Val Leu Lys Phe Leu His Asp
 270 275 280

aag gga ttc cct tat ggg cat ctt cac gcc tcc aat gtg 947
 Lys Gly Phe Pro Tyr Gly His Leu His Ala Ser Asn Val
 285 290

atg ctc gat ggg gac act tgt cgg ctg ctg gac ctt gag 986
Met Leu Asp Gly Asp Thr Cys Arg Leu Leu Asp Leu Glu
295 300 305

aat tcc tta ttg ggc ctg cct tcc ttc tac cga tct tat 1025
Asn Ser Leu Leu Gly Leu Pro Ser Phe Tyr Arg Ser Tyr
310 315 320

ttt tca caa ttc agg aaa atc aat aca ttg gaa agt gtg 1064
Phe Ser Gln Phe Arg Lys Ile Asn Thr Leu Glu Ser Val
325 330

gat gtc cac tgc ttt ggc cac tta ctg tat gaa atg act 1103
Asp Val His Cys Phe Gly His Leu Leu Tyr Glu Met Thr
335 340 345

tat gga cga ccg cca gac tcg gtg cct gtg gac tcc ttc 1142
Tyr Gly Arg Pro Pro Asp Ser Val Pro Val Asp Ser Phe
350 355

cct cct gcc ccg tcc atg gct gtg gtg gcc gtg ttg gag 1181
Pro Pro Ala Pro Ser Met Ala Val Val Ala Val Leu Glu
360 365 370

tct acg ctg tct tgt gaa gcc tgt aaa aat ggc atg cct 1220
Ser Thr Leu Ser Cys Glu Ala Cys Lys Asn Gly Met Pro
375 380 385

acc atc tcc cgg ctc tta cag atg cca tta ttc agc gat 1259
Thr Ile Ser Arg Leu Leu Gln Met Pro Leu Phe Ser Asp
390 395

gtt tta cta acc act tct gaa aaa cca cag ttt aag atc 1298
Val Leu Leu Thr Thr Ser Glu Lys Pro Gln Phe Lys Ile
400 405 410

cct aca aag tta aaa gag gca ttg aga att gcc aaa gaa 1337
Pro Thr Lys Leu Lys Glu Ala Leu Arg Ile Ala Lys Glu
415 420

tgt ata gag aag aga cta att gag gaa cag aaa cag att 1376
Cys Ile Glu Lys Arg Leu Ile Glu Glu Gln Lys Gln Ile
425 430 435

cac cag cat cga aga ctg aca aga gct cag tcc cac cat 1415
His Gln His Arg Arg Leu Thr Arg Ala Gln Ser His His
440 445 450

gga tct gag gag gaa aga aaa aaa aga aag att tta gct 1454
Gly Ser Glu Glu Glu Arg Lys Lys Arg Lys Ile Leu Ala
455 460

cga aag aag tca aaa cga tct gct ctt gaa aat agt gaa 1493
Arg Lys Lys Ser Lys Arg Ser Ala Leu Glu Asn Ser Glu
465 470 475

gag cat tca gcg agg tac agc aac tcc aat aat tca gga 1532
Glu His Ser Ala Arg Tyr Ser Asn Ser Asn Asn Ser Gly
480 485

tct ggg gcc agc tca cct ctc acg tcc ccg tca tcg cca 1571

Ser Gly Ala Ser Ser Pro Leu Thr Ser Pro Ser Ser Pro
490 495 500

act cca ccc tct aca tca ggg ata tct gca tta cct cca 1610
Thr Pro Pro Ser Thr Ser Gly Ile Ser Ala Leu Pro Pro
505 510 515

cct cct cca cct cca cca cca cca gca gct ccc ttg cct 1649
Pro Pro Pro Pro Pro Pro Pro Pro Ala Ala Pro Leu Pro
520 525

cct gcg agc acc gag gta cct gcc cag ctc tcg tct cag 1688
Pro Ala Ser Thr Glu Val Pro Ala Gln Leu Ser Ser Gln
530 535 540

gct gtg aat ggc atg agc cga ggg gcc ttg ctc agc tcc 1727
Ala Val Asn Gly Met Ser Arg Gly Ala Leu Leu Ser Ser
545 550

atc cag aat ttc caa aaa gga act ttg agg aaa gcc aaa 1766
Ile Gln Asn Phe Gln Lys Gly Thr Leu Arg Lys Ala Lys
555 560 565

acc tgt gat cac agt gct ccg aag atc ggc tg aa 1800
Thr Cys Asp His Ser Ala Pro Lys Ile Gly
570 575 577

gcttcctggt tacacttgga gggaaaagtt cttttttatt cctactcacc 1850
cctaccccc aaactaccct cttcctggga aagtaattgc tgagccagta 1900
cagccacaaa cagtactatt ttgcagatgc tcatgtaagc agcttttcga 1950
gagaaataat tctttaagca gaataaagtt aggctggcat tgctccctta 2000
agatottgct cctttattaa ccctgtaaag gagtcttggt tatcctctaa 2050
tgccaggct tttgggacag cagcatattg aaatattttc accaactaaa 2100
ggaaatagac agaaaaacaa tgacaatatt caatcacagc agtaaatggc 2150
ctttgtgttg caatcccttc taccatca gacagctcct agaaacattc 2200
cttacagttc atttctctaa agcattttct gattcttaga taactccaat 2250
ttttgctacc tttatcttag acattaacac tatagcccaa agcatagtta 2300
ctttgctaaa tcagaaagca actgagttct ttgttttctc ctcaaataga 2350
atggggaacg ttcacaacat tctcttaagt tctaacagga ataccattgt 2400
ggttatagaa ctcagggtg ctaaagcaac tactctagac ccatagttct 2450
tttagttag atgtattgaa acagacaaaa atattaacat cagaaaaagc 2500
tcttgccaat tagaggatct tcttaatcct cagcaattaa gtttggggtt 2550
tgaggggggc aggtcattgt tacaacagaa gtaaatgttg catctataga 2600
aatcaattat gatttttgaa agatttatct aaatatatca atatagcatc 2650

tctttaatgt tagtcattta ttagaaagat cctttatcct gatttgctta 2700
aacctttcaa taaattgcac tttaaaggat tataaataat ccatttaaaa 2750
attcaagtac acacatcagt gttggttact atgcagagaa tgtcattgtg 2800
tatagtttca tgtaatctgt tatgtcagct gtatttttta ttaaaatcat 2850
gtcaagaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a 2891

<210> 2
<211> 577
<212> PRT
<213> Homo sapiens

<400> 2
Met Ala Phe Met Glu Lys Pro Pro Ala Gly Lys Val Leu Leu Asp
1 5 10 15
Asp Thr Val Pro Leu Thr Ala Ala Ile Glu Ala Ser Gln Ser Leu
20 25 30
Gln Ser His Thr Glu Tyr Ile Ile Arg Val Gln Arg Gly Ile Ser
35 40 45
Val Glu Asn Ser Trp Gln Ile Val Arg Arg Tyr Ser Asp Phe Asp
50 55 60
Leu Leu Asn Asn Ser Leu Gln Ile Ala Gly Leu Ser Leu Pro Leu
65 70 75
Pro Pro Lys Lys Leu Ile Gly Asn Met Asp Arg Glu Phe Ile Ala
80 85 90
Glu Arg Gln Lys Gly Leu Gln Asn Tyr Leu Asn Val Ile Thr Thr
95 100 105
Asn His Ile Leu Ser Asn Cys Glu Leu Val Lys Lys Phe Leu Asp
110 115 120
Pro Asn Asn Tyr Ser Ala Asn Tyr Thr Glu Ile Ala Leu Gln Gln
125 130 135
Val Ser Met Phe Phe Arg Ser Glu Pro Lys Trp Glu Val Val Glu
140 145 150
Pro Leu Lys Asp Ile Gly Trp Arg Ile Arg Lys Lys Tyr Phe Leu
155 160 165
Met Lys Ile Lys Asn Gln Pro Lys Glu Arg Leu Val Leu Ser Trp
170 175 180
Ala Asp Leu Gly Pro Asp Lys Tyr Leu Ser Asp Lys Asp Phe Gln
185 190 195
Cys Leu Ile Lys Leu Leu Pro Ser Cys Leu His Pro Tyr Ile Tyr
200 205 210
Arg Val Thr Phe Ala Thr Ala Asn Glu Ser Ser Ala Leu Leu Ile
215 220 225

0968453 "100500"

Arg Met Phe Asn Glu Lys Gly Thr Leu Lys Asp Leu Ile Tyr Lys	230	235	240
Ala Lys Pro Lys Asp Pro Phe Leu Lys Lys Tyr Cys Asn Pro Lys	245	250	255
Lys Ile Gln Gly Leu Glu Leu Gln Gln Ile Lys Thr Tyr Gly Arg	260	265	270
Gln Ile Leu Glu Val Leu Lys Phe Leu His Asp Lys Gly Phe Pro	275	280	285
Tyr Gly His Leu His Ala Ser Asn Val Met Leu Asp Gly Asp Thr	290	295	300
Cys Arg Leu Leu Asp Leu Glu Asn Ser Leu Leu Gly Leu Pro Ser	305	310	315
Phe Tyr Arg Ser Tyr Phe Ser Gln Phe Arg Lys Ile Asn Thr Leu	320	325	330
Glu Ser Val Asp Val His Cys Phe Gly His Leu Leu Tyr Glu Met	335	340	345
Thr Tyr Gly Arg Pro Pro Asp Ser Val Pro Val Asp Ser Phe Pro	350	355	360
Pro Ala Pro Ser Met Ala Val Val Ala Val Leu Glu Ser Thr Leu	365	370	375
Ser Cys Glu Ala Cys Lys Asn Gly Met Pro Thr Ile Ser Arg Leu	380	385	390
Leu Gln Met Pro Leu Phe Ser Asp Val Leu Leu Thr Thr Ser Glu	395	400	405
Lys Pro Gln Phe Lys Ile Pro Thr Lys Leu Lys Glu Ala Leu Arg	410	415	420
Ile Ala Lys Glu Cys Ile Glu Lys Arg Leu Ile Glu Glu Gln Lys	425	430	435
Gln Ile His Gln His Arg Arg Leu Thr Arg Ala Gln Ser His His	440	445	450
Gly Ser Glu Glu Glu Arg Lys Lys Arg Lys Ile Leu Ala Arg Lys	455	460	465
Lys Ser Lys Arg Ser Ala Leu Glu Asn Ser Glu Glu His Ser Ala	470	475	480
Arg Tyr Ser Asn Ser Asn Asn Ser Gly Ser Gly Ala Ser Ser Pro	485	490	495
Leu Thr Ser Pro Ser Ser Pro Thr Pro Pro Ser Thr Ser Gly Ile	500	505	510
Ser Ala Leu Pro Pro Pro Pro Pro Pro Pro Pro Pro Ala Ala	515	520	525
Pro Leu Pro Pro Ala Ser Thr Glu Val Pro Ala Gln Leu Ser Ser			

530	535	540
Gln Ala Val Asn Gly Met Ser Arg Gly Ala Leu Leu Ser Ser Ile		
545	550	555
Gln Asn Phe Gln Lys Gly Thr Leu Arg Lys Ala Lys Thr Cys Asp		
560	565	570
His Ser Ala Pro Lys Ile Gly		
575		

<210> 3
 <211> 2119
 <212> DNA
 <213> Homo sapiens

<400> 3
 tgcacctcgg ttctatcgat tgaattcgg ccacactggc cggatcctct 50
 agagatccct cgacctcgac ccacgcgtcc gctcaaggaa gttgaatgag 100
 gtgaacatag atacagctct tcccccttcc cccccctttt aaatgtaaca 150
 aatacttttt atgttccccct tcccccttcc cctttttccc ctttccccctt 200
 ttggaaacgt gtcaggaacc aaatagttta agatgagcag ttgagggggac 250
 tgagagagtg atcgacacag aacctgggctt cttcgtgctt catcataagt 300
 cgtgctgccg gccaggttac ttaagcacc ttttaacaag gaaaccttgt 350
 gggagatcca gctggccgac tcgagttcag aaacaggacc acagaggtta 400
 cactctggga tcctggccat gaggttggat gcctcacctt actgaaagga 450
 gacactggac cttaa atg gcg cag cat gat ttt gtt cct gct 491
 Met Ala Gln His Asp Phe Val Pro Ala
 1 5
 tgg cta aat ttc tca aca cca cag tca gct aag tca cct 530
 Trp Leu Asn Phe Ser Thr Pro Gln Ser Ala Lys Ser Pro
 10 15 20
 act gcc acc ttc gaa aaa cac gga gag cac cta ccc aga 569
 Thr Ala Thr Phe Glu Lys His Gly Glu His Leu Pro Arg
 25 30 35
 gga gaa ggt aga ttt gga gta agc cgc cgt cga cat aat 608
 Gly Glu Gly Arg Phe Gly Val Ser Arg Arg Arg His Asn
 40 45
 tcc tct gat ggt ttt ttt aac aat ggt ccc cta cga act 647
 Ser Ser Asp Gly Phe Phe Asn Asn Gly Pro Leu Arg Thr
 50 55 60
 gca gga gat tct tgg cac cag ccc tcc tgt tcc gcc atg 686
 Ala Gly Asp Ser Trp His Gln Pro Ser Cys Ser Ala Met
 65 70
 att ctg tgg act ctg gtg tct cta agg gag cat atg ctg 725
 Ile Leu Trp Thr Leu Val Ser Leu Arg Glu His Met Leu

75		80		85	
gaa tca cag gga acc cat ctg gtt ggc ata gct ctt ccc 764					
Glu Ser Gln Gly Thr His Leu Val Gly Ile Ala Leu Pro					
90		95		100	
cga ggt cat gat ggc atg agc caa cgt agt gta ggt ggc 803					
Arg Gly His Asp Gly Met Ser Gln Arg Ser Val Gly Gly					
105		110			
aca ggg aac cat cgc cat tgg aat ggc agc ttc cac tcc 842					
Thr Gly Asn His Arg His Trp Asn Gly Ser Phe His Ser					
115		120		125	
cgg aaa ggg tgt gct ttt cag gaa aag cca cct atg gag 881					
Arg Lys Gly Cys Ala Phe Gln Glu Lys Pro Pro Met Glu					
130		135			
att agg gaa gaa aag aaa gaa gac aag gtg gaa aag ttg 920					
Ile Arg Glu Glu Lys Lys Glu Asp Lys Val Glu Lys Leu					
140		145		150	
cag ttt gaa gag gag gac ttt cct tcc ttg aat cca gaa 959					
Gln Phe Glu Glu Glu Asp Phe Pro Ser Leu Asn Pro Glu					
155		160		165	
gct ggc aaa cag cat cag cca tgc aga cct att ggg aca 998					
Ala Gly Lys Gln His Gln Pro Cys Arg Pro Ile Gly Thr					
170		175			
cct tct gga gta tgg gaa aac ccg cct agt gcc aag caa 1037					
Pro Ser Gly Val Trp Glu Asn Pro Pro Ser Ala Lys Gln					
180		185		190	
ccc tcc aag atg cta gtt atc aaa aaa gtt tcc aaa gag 1076					
Pro Ser Lys Met Leu Val Ile Lys Lys Val Ser Lys Glu					
195		200			
gat cct gct gct gcc ttc tct gct gca ttc acc tca cca 1115					
Asp Pro Ala Ala Ala Phe Ser Ala Ala Phe Thr Ser Pro					
205		210		215	
gga tct cac cat gca aat ggg aac aaa ttg tca tcc gtg 1154					
Gly Ser His His Ala Asn Gly Asn Lys Leu Ser Ser Val					
220		225		230	
gtt cca agt gtc tat aag aac ctg gtt cct aag cct gta 1193					
Val Pro Ser Val Tyr Lys Asn Leu Val Pro Lys Pro Val					
235		240			
cca cct cct tcc aag cct aat gca tgg aaa gct aac agg 1232					
Pro Pro Pro Ser Lys Pro Asn Ala Trp Lys Ala Asn Arg					
245		250		255	
atg gag cac aag tca gga tcc ctt tcc tct agc cgg gag 1271					
Met Glu His Lys Ser Gly Ser Leu Ser Ser Ser Arg Glu					
260		265			
tct gct ttt acc agt cca atc tcc gtt acc aaa cca gtg 1310					
Ser Ala Phe Thr Ser Pro Ile Ser Val Thr Lys Pro Val					
270		275		280	

gta ctg gct agt ggt gca gct ctg agt tct ccc aaa gag 1349
 Val Leu Ala Ser Gly Ala Ala Leu Ser Ser Pro Lys Glu
 285 290 295

agt ccc tcc agc acc acc cct cca att gag atc agc tcc 1388
 Ser Pro Ser Ser Thr Thr Pro Pro Ile Glu Ile Ser Ser
 300 305

tct cgt ctg acc aag ttg acc cgc cga acc acc gac agg 1427
 Ser Arg Leu Thr Lys Leu Thr Arg Arg Thr Thr Asp Arg
 310 315 320

aag agt gag ttc ctg aaa act ctg aag gat gac cgg aat 1466
 Lys Ser Glu Phe Leu Lys Thr Leu Lys Asp Asp Arg Asn
 325 330

gga gac ttc tca gag aat aga gac tgt gac aag ctg gaa 1505
 Gly Asp Phe Ser Glu Asn Arg Asp Cys Asp Lys Leu Glu
 335 340 345

gat ttg gag gac aac agc aca cct gaa cca aag gaa aat 1544
 Asp Leu Glu Asp Asn Ser Thr Pro Glu Pro Lys Glu Asn
 350 355 360

ggg gag gaa ggc tgt cat caa aat ggt ctt gcc ctc cct 1583
 Gly Glu Glu Gly Cys His Gln Asn Gly Leu Ala Leu Pro
 365 370

gta gtg gaa gaa ggg gag gtt ctc tca cac tct cta gaa 1622
 Val Val Glu Glu Gly Glu Val Leu Ser His Ser Leu Glu
 375 380 385

gca gag cac agg tta ttg aaa gct atg ggt tgg cag gaa 1661
 Ala Glu His Arg Leu Leu Lys Ala Met Gly Trp Gln Glu
 390 395

tat cct gaa aat gat gag aat tgc ctt ccc ctc aca gag 1700
 Tyr Pro Glu Asn Asp Glu Asn Cys Leu Pro Leu Thr Glu
 400 405 410

gat gag ctc aaa gag ttc cac atg aag aca gag cag ctg 1739
 Asp Glu Leu Lys Glu Phe His Met Lys Thr Glu Gln Leu
 415 420 425

aga aga aat ggc ttt gga aag aat ggc ttc ttg cag agc 1778
 Arg Arg Asn Gly Phe Gly Lys Asn Gly Phe Leu Gln Ser
 430 435

cgc agt tcc agt ctg ttc tcc cct tgg aga agc act tgc 1817
 Arg Ser Ser Ser Leu Phe Ser Pro Trp Arg Ser Thr Cys
 440 445 450

aaa gca gag ttt gag gac tca gac acc gaa acc agt agc 1856
 Lys Ala Glu Phe Glu Asp Ser Asp Thr Glu Thr Ser Ser
 455 460

agt gaa aca tca gat gac gat gcc tgg aag t agg 1890
 Ser Glu Thr Ser Asp Asp Asp Ala Trp Lys
 465 470 474

catataaatg ctcacagtta aatctgaccc agtaaaactct gtgtgttttag 1940
 ggagtataca aaagaaatcg ttcttttcct tttcttatgt tgttgaatac 1990
 ttcattcaca agggaaataa tcatatccca aagagagaaa aaaaaaaaaa 2040
 aaagggcggc cgcgactcta gagtcgacct gcagtaggga taacagggta 2090
 ataagcttgg cgcgcatggc ccaacttgt 2119

<210> 4
 <211> 474
 <212> PRT
 <213> Homo sapiens

<400> 4
 Met Ala Gln His Asp Phe Val Pro Ala Trp Leu Asn Phe Ser Thr
 1 5 10 15
 Pro Gln Ser Ala Lys Ser Pro Thr Ala Thr Phe Glu Lys His Gly
 20 25 30
 Glu His Leu Pro Arg Gly Glu Gly Arg Phe Gly Val Ser Arg Arg
 35 40 45
 Arg His Asn Ser Ser Asp Gly Phe Phe Asn Asn Gly Pro Leu Arg
 50 55 60
 Thr Ala Gly Asp Ser Trp His Gln Pro Ser Cys Ser Ala Met Ile
 65 70 75
 Leu Trp Thr Leu Val Ser Leu Arg Glu His Met Leu Glu Ser Gln
 80 85 90
 Gly Thr His Leu Val Gly Ile Ala Leu Pro Arg Gly His Asp Gly
 95 100 105
 Met Ser Gln Arg Ser Val Gly Gly Thr Gly Asn His Arg His Trp
 110 115 120
 Asn Gly Ser Phe His Ser Arg Lys Gly Cys Ala Phe Gln Glu Lys
 125 130 135
 Pro Pro Met Glu Ile Arg Glu Glu Lys Lys Glu Asp Lys Val Glu
 140 145 150
 Lys Leu Gln Phe Glu Glu Glu Asp Phe Pro Ser Leu Asn Pro Glu
 155 160 165
 Ala Gly Lys Gln His Gln Pro Cys Arg Pro Ile Gly Thr Pro Ser
 170 175 180
 Gly Val Trp Glu Asn Pro Pro Ser Ala Lys Gln Pro Ser Lys Met
 185 190 195
 Leu Val Ile Lys Lys Val Ser Lys Glu Asp Pro Ala Ala Ala Phe
 200 205 210
 Ser Ala Ala Phe Thr Ser Pro Gly Ser His His Ala Asn Gly Asn
 215 220 225

Lys	Leu	Ser	Ser	Val	Val	Pro	Ser	Val	Tyr	Lys	Asn	Leu	Val	Pro	230	235	240
Lys	Pro	Val	Pro	Pro	Pro	Ser	Lys	Pro	Asn	Ala	Trp	Lys	Ala	Asn	245	250	255
Arg	Met	Glu	His	Lys	Ser	Gly	Ser	Leu	Ser	Ser	Ser	Arg	Glu	Ser	260	265	270
Ala	Phe	Thr	Ser	Pro	Ile	Ser	Val	Thr	Lys	Pro	Val	Val	Leu	Ala	275	280	285
Ser	Gly	Ala	Ala	Leu	Ser	Ser	Pro	Lys	Glu	Ser	Pro	Ser	Ser	Thr	290	295	300
Thr	Pro	Pro	Ile	Glu	Ile	Ser	Ser	Ser	Arg	Leu	Thr	Lys	Leu	Thr	305	310	315
Arg	Arg	Thr	Thr	Asp	Arg	Lys	Ser	Glu	Phe	Leu	Lys	Thr	Leu	Lys	320	325	330
Asp	Asp	Arg	Asn	Gly	Asp	Phe	Ser	Glu	Asn	Arg	Asp	Cys	Asp	Lys	335	340	345
Leu	Glu	Asp	Leu	Glu	Asp	Asn	Ser	Thr	Pro	Glu	Pro	Lys	Glu	Asn	350	355	360
Gly	Glu	Glu	Gly	Cys	His	Gln	Asn	Gly	Leu	Ala	Leu	Pro	Val	Val	365	370	375
Glu	Glu	Gly	Glu	Val	Leu	Ser	His	Ser	Leu	Glu	Ala	Glu	His	Arg	380	385	390
Leu	Leu	Lys	Ala	Met	Gly	Trp	Gln	Glu	Tyr	Pro	Glu	Asn	Asp	Glu	395	400	405
Asn	Cys	Leu	Pro	Leu	Thr	Glu	Asp	Glu	Leu	Lys	Glu	Phe	His	Met	410	415	420
Lys	Thr	Glu	Gln	Leu	Arg	Arg	Asn	Gly	Phe	Gly	Lys	Asn	Gly	Phe	425	430	435
Leu	Gln	Ser	Arg	Ser	Ser	Ser	Leu	Phe	Ser	Pro	Trp	Arg	Ser	Thr	440	445	450
Cys	Lys	Ala	Glu	Phe	Glu	Asp	Ser	Asp	Thr	Glu	Thr	Ser	Ser	Ser	455	460	465
Glu	Thr	Ser	Asp	Asp	Asp	Ala	Trp	Lys							470		

<210> 5

<211> 294

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> 162, 165, 183-184, 187, 204, 207, 211, 216, 219, 222-223, 225, 237, 240, 244, 250, 253, 262, 267, 269, 275, 279, 282, 289-290

<223> unknown base

```

<400> 5
cccgggatgg ccttcatgga gaagccgcca gctggcaagg tgctgctgga 50
cgacacgggtg ccgctgacag cagccatcga ggcgagccag agcctgcagt 100
cccacacgga atatattatt cgagtgcaaa gaggaatttc tctggaaaac 150
agctggcaga tngtnagaag atacagtgac tttnatntgc ttaacaacag 200
cttncanatt ncaggncrna gnntncctct tcctccnaan aaantgattn 250
ggnaacatgg ancgtgnant tcatngctng anaggcagnn aggt 294

<210> 6
<211> 24
<212> DNA
<213> Homo sapiens

<400> 6
ggacgacacg gtgccgctga cagc 24

<210> 7
<211> 51
<212> DNA
<213> Homo sapiens

<400> 7
gccatcgagg cgagccagag cctgcagtcc cacacggaat atattattcg 50
a 51

<210> 8
<211> 31
<212> DNA
<213> Homo sapiens

<400> 8
gttttccaga gaaattcctc tttgcaactcg a 31

<210> 9
<211> 793
<212> DNA
<213> Homo sapiens

<220>
<221> unsure
<222> 379, 433, 459, 492, 517, 541, 549, 561, 575, 579, 582, 710, 742,
774, 784
<223> unknown base

<400> 9
gggaagaaaa gaaagaagac aaggtggaaa agttgcagat ttgaagagga 50
ggactttcct tccttgaatc cagaagctgg caaacagcat cagccatgca 100
gacctattgg gacaccttct ggagtatggg aaaaccogcc tagtgccaag 150
caaccctcca agatgctagt tatcaaaaaa gtttccaaag aggatcctgc 200
tgctgcyttc tctgctgcat tcacctcacc aggatctcac catgcaaata 250

```

ggaacaaatt gtcacccgtg gttccaagtg tctataagaa cctggttcct 300
aagcctgtac cacctccttc caagcctaata gcatggaaag ctaacaggat 350
ggagcacaag tcaggatccc ttctctctna gccgggagtc tgcttttacc 400
agtccaatct ctgttaccaa accagtggta ctnggctagt ggtgcagctc 450
tgagttctnc ccaaagagag tccctccagc accaccctc cnaattgaga 500
tcagctcctc tcgtctngac caagtttgac ccgccgaacc naccgacang 550
gaagragtga ngttcctgaa aactnctgna anggatgacc gggaatggga 600
agacttityc agaagaatag agactgtgac aagctggaag atttgaggga 650
caacagcaca ctgaaccaa ggaaaatggg ggaggaaggc tgttcatcaa 700
atgggtctttn ccttctctgta gttggaagga ggggaggttt tntcacattt 750
ttaggagcag agcacagggt tttnaagttt tggntggcgg att 793

<210> 10
<211> 24
<212> DNA
<213> Homo sapiens

<400> 10
gacctattgg gacaccttct ggag 24

<210> 11
<211> 43
<212> DNA
<213> Homo sapiens

<400> 11
ccgcctagt gccaaagcaac cctccaagat gctagttatc aaa 43

<210> 12
<211> 25
<212> DNA
<213> Homo sapiens

<400> 12
cttggtcaga cgagaggagc tgatc 25

<210> 13
<211> 2366
<212> DNA
<213> Homo sapiens

<220>
<221> unsure
<222> 103
<223> unknown base

<400> 13
tgcccgatc ctctagagat ccctcgacct cgaccacgc gtccgcagaa 50

agtatgtccc ttctctcacc atg agc tgg ctc tcc agt tcc 91

														Met	Ser	Trp	Leu	Ser	Ser	Ser	
														1				5			
cag	gga	gtg	gtn	cta	aca	gcc	tac	cac	ccc	agc	ggc	aag	130								
Gln	Gly	Val	Xaa	Leu	Thr	Ala	Tyr	His	Pro	Ser	Gly	Lys									
				10					15					20							
gac	cag	acc	gtc	ggg	aac	agc	cat	gca	aag	gca	ggg	gag	169								
Asp	Gln	Thr	Val	Gly	Asn	Ser	His	Ala	Lys	Ala	Gly	Glu									
				25					30												
gaa	gcc	acc	tcg	agt	cgc	aga	tat	ggc	cag	tac	act	atg	208								
Glu	Ala	Thr	Ser	Ser	Arg	Arg	Tyr	Gly	Gln	Tyr	Thr	Met									
				35					40					45							
aac	cag	gaa	agc	acc	acc	atc	aaa	gtt	atg	gag	aag	cct	247								
Asn	Gln	Glu	Ser	Thr	Thr	Ile	Lys	Val	Met	Glu	Lys	Pro									
				50					55												
cca	ttt	gat	cga	tca	att	tcc	cag	gat	tct	ttg	gat	gaa	286								
Pro	Phe	Asp	Arg	Ser	Ile	Ser	Gln	Asp	Ser	Leu	Asp	Glu									
				60					65					70							
cta	tct	atg	gaa	gac	tat	tgg	ata	gaa	cta	gaa	aac	atc	325								
Leu	Ser	Met	Glu	Asp	Tyr	Trp	Ile	Glu	Leu	Glu	Asn	Ile									
				75					80					85							
aag	aaa	tct	agt	gaa	aac	agc	caa	gaa	gat	caa	gag	gtg	364								
Lys	Lys	Ser	Ser	Glu	Asn	Ser	Gln	Glu	Asp	Gln	Glu	Val									
				90					95												
gtt	gtt	gtc	aaa	gag	cct	gat	gag	gga	gaa	ttg	gaa	gaa	403								
Val	Val	Val	Lys	Glu	Pro	Asp	Glu	Gly	Glu	Leu	Glu	Glu									
				100					105					110							
gag	tgg	ctt	aaa	gag	gcc	ggt	tta	tcc	aat	ctc	ttc	gga	442								
Glu	Trp	Leu	Lys	Glu	Ala	Gly	Leu	Ser	Asn	Leu	Phe	Gly									
				115					120												
gag	tct	gct	gga	gat	cca	cag	gaa	agc	att	gtg	ttt	tta	481								
Glu	Ser	Ala	Gly	Asp	Pro	Gln	Glu	Ser	Ile	Val	Phe	Leu									
				125					130					135							
tca	aca	ttg	acg	cgg	acc	cag	gca	gca	gca	gtt	cag	aag	520								
Ser	Thr	Leu	Thr	Arg	Thr	Gln	Ala	Ala	Ala	Val	Gln	Lys									
				140					145					150							
cga	gta	gag	acg	gtc	tcc	cag	acc	ttg	agg	aaa	aaa	aac	559								
Arg	Val	Glu	Thr	Val	Ser	Gln	Thr	Leu	Arg	Lys	Lys	Asn									
				155					160												
aaa	cag	tac	cag	att	cct	gac	gtc	aga	gac	ata	ttt	gct	598								
Lys	Gln	Tyr	Gln	Ile	Pro	Asp	Val	Arg	Asp	Ile	Phe	Ala									
				165					170					175							
caa	cag	aga	gaa	tca	aaa	gaa	aca	gct	cca	ggt	ggc	act	637								
Gln	Gln	Arg	Glu	Ser	Lys	Glu	Thr	Ala	Pro	Gly	Gly	Thr									
				180					185												
gaa	tcg	cag	tca	ctt	aga	aca	aat	gaa	aac	aaa	tac	caa	676								
Glu	Ser	Gln	Ser	Leu	Arg	Thr	Asn	Glu	Asn	Lys	Tyr	Gln									

190		195		200	
gga aga gat gac gag gca tct aac ctt gtt ggt gaa gag 715					
Gly Arg Asp Asp Glu Ala Ser Asn Leu Val Gly Glu Glu					
205			210		215
aag ctg atc cca cct gag gag acg cct gcc cct gaa aca 754					
Lys Leu Ile Pro Pro Glu Glu Thr Pro Ala Pro Glu Thr					
220			225		
gac atc aac ctg gag gta tca ttt gcc gag caa gca ctc 793					
Asp Ile Asn Leu Glu Val Ser Phe Ala Glu Gln Ala Leu					
230			235		240
aat cag aaa gag aga tcc aag gag aaa atc cag aag agc 832					
Asn Gln Lys Glu Arg Ser Lys Glu Lys Ile Gln Lys Ser					
245			250		
aaa ggc gat gat gcc aca tta cct agt ttc aga ttg cca 871					
Lys Gly Asp Asp Ala Thr Leu Pro Ser Phe Arg Leu Pro					
255			260		265
aaa gac aaa acg ggt acc aca agg att ggt gac ctc gca 910					
Lys Asp Lys Thr Gly Thr Thr Arg Ile Gly Asp Leu Ala					
270			275		280
ccc cag gac atg aag aaa gtt tgc cat tta gcc cta att 949					
Pro Gln Asp Met Lys Lys Val Cys His Leu Ala Leu Ile					
285			290		
gag ctg act gcc ctc tat gat gta ttg ggt att gag ctg 988					
Glu Leu Thr Ala Leu Tyr Asp Val Leu Gly Ile Glu Leu					
295			300		305
aaa caa caa aaa gct gtg aaa atc aaa aca aaa gat tct 1027					
Lys Gln Gln Lys Ala Val Lys Ile Lys Thr Lys Asp Ser					
310			315		
ggt ctt ttt tgc gtt cca ttg aca gcg cta tta gaa caa 1066					
Gly Leu Phe Cys Val Pro Leu Thr Ala Leu Leu Glu Gln					
320			325		330
gat cag agg aaa gta cca gga atg cga ata ccc ttg atc 1105					
Asp Gln Arg Lys Val Pro Gly Met Arg Ile Pro Leu Ile					
335			340		345
ttt caa aaa ctg att tct cga att gaa gag aga ggt ttg 1144					
Phe Gln Lys Leu Ile Ser Arg Ile Glu Glu Arg Gly Leu					
350			355		
gaa aca gaa ggc ctc tta cgg atc cct gga gct gcc att 1183					
Glu Thr Glu Gly Leu Leu Arg Ile Pro Gly Ala Ala Ile					
360			365		370
aga atc aag aat ctt tgc caa gaa cta gaa gca aag ttt 1222					
Arg Ile Lys Asn Leu Cys Gln Glu Leu Glu Ala Lys Phe					
375			380		
tat gaa ggg act ttt aat tgg gaa agt gtc aaa cag cat 1261					
Tyr Glu Gly Thr Phe Asn Trp Glu Ser Val Lys Gln His					
385			390		395

```

gat gcc gcc agc ctg ctg aag ctc ttc att cgg gag ttg 1300
Asp Ala Ala Ser Leu Leu Lys Leu Phe Ile Arg Glu Leu
      400              405              410

ccc cag cca ctg ctc agt gtg gag tat ctc aaa gcc ttt 1339
Pro Gln Pro Leu Leu Ser Val Glu Tyr Leu Lys Ala Phe
      415              420

cag gct gtc cag aat ctt cca acc aag aag cag caa cta 1378
Gln Ala Val Gln Asn Leu Pro Thr Lys Lys Gln Gln Leu
      425              430              435

cag gct ttg aac ctt ctt gtc atc ctc cta cct gat gca 1417
Gln Ala Leu Asn Leu Leu Val Ile Leu Leu Pro Asp Ala
      440              445

aac agg gac aca ctg aag gcc ctt ctt gaa ttt ctc caa 1456
Asn Arg Asp Thr Leu Lys Lys Ala Leu Leu Glu Phe Leu Gln
      450              455              460

aga gta ata gat aat aaa gaa aaa aat aaa atg aca gtc 1495
Arg Val Ile Asp Asn Lys Glu Lys Asn Lys Met Thr Val
      465              470              475

atg aat gta gca atg gtc atg gcc ccg aat ctc ttt atg 1534
Met Asn Val Ala Met Val Met Ala Pro Asn Leu Phe Met
      480              485

tgt cat gca ttg gga ttg aag tcc agt gaa cag cga gaa 1573
Cys His Ala Leu Gly Leu Lys Ser Ser Glu Gln Arg Glu
      490              495              500

ttt gta atg gca gct ggg aca gca aat acc atg cac tta 1612
Phe Val Met Ala Ala Gly Thr Ala Asn Thr Met His Leu
      505              510

ttg att aag tac caa aaa ctt ctg tgg aca att ccc aag 1651
Leu Ile Lys Tyr Gln Lys Leu Leu Trp Thr Ile Pro Lys
      515              520              525

ttt att gta aac caa gtg agg aag caa aac acg gaa aat 1690
Phe Ile Val Asn Gln Val Arg Lys Gln Asn Thr Glu Asn
      530              535              540

cat aaa aag gat aaa aga gcc atg aag aaa ttg ctg aag 1729
His Lys Lys Asp Lys Arg Ala Met Lys Lys Leu Leu Lys
      545              550

aaa atg gct tat gac cga gaa aaa tat gaa aag caa gat 1768
Lys Met Ala Tyr Asp Arg Glu Lys Tyr Glu Lys Gln Asp
      555              560              565

aag agt aca aat gat gct gac gtt cct cag gga gtg att 1807
Lys Ser Thr Asn Asp Ala Asp Val Pro Gln Gly Val Ile
      570              575

cga gtg caa gct ccc cat ctt tcg aaa gtt tcc atg gca 1846
Arg Val Gln Ala Pro His Leu Ser Lys Val Ser Met Ala
      580              585              590

```

ata cag cta act gaa gaa cta aaa gcc agt gat gta ctt 1885
 Ile Gln Leu Thr Glu Glu Leu Lys Ala Ser Asp Val Leu
 595 600 605

gcc agg ttt ctc agc caa gaa agt ggg gtt gcc cag act 1924
 Ala Arg Phe Leu Ser Gln Glu Ser Gly Val Ala Gln Thr
 610 615

ctc aag aaa gga gaa gtt ttt ttg tat gaa att gga gga 1963
 Leu Lys Lys Gly Glu Val Phe Leu Tyr Glu Ile Gly Gly
 620 625 630

aat att ggg gaa cgc tgc ctt gat gat gac act tac atg 2002
 Asn Ile Gly Glu Arg Cys Leu Asp Asp Asp Thr Tyr Met
 635 640

aag gat tta tat cag ctt aac cca aat gct gag tgg gtt 2041
 Lys Asp Leu Tyr Gln Leu Asn Pro Asn Ala Glu Trp Val
 645 650 655

ata aag tca aag cca ttg t agaagactta acaagctgca 2080
 Ile Lys Ser Lys Pro Leu
 660 663

gataaccatg tggacttctg tcataattct tgctgagtca agagtgtaaa 2130

taaaagaaat ggcaggactc atattattca gttgtaccca agtatttttaa 2180

aaatgactct cttaagcctt aaaaagtcac agatttgtgc tgctgccaga 2230

attatattaa attattatta atggtattat tagaapaaaaa aatttctgga 2280

gtgagagtaa agaggcttaa ttagtttgtg ggcagttttc ttatgctctg 2330

tgaaatgtgt ccagatgtga catagttttt tttttt 2366

<210> 14
 <211> 663
 <212> PRT
 <213> Homo sapiens

<400> 14
 Met Ser Trp Leu Ser Ser Ser Gln Gly Val Val Leu Thr Ala Tyr
 1 5 10 15
 His Pro Ser Gly Lys Asp Gln Thr Val Gly Asn Ser His Ala Lys
 20 25 30
 Ala Gly Glu Glu Ala Thr Ser Ser Arg Arg Tyr Gly Gln Tyr Thr
 35 40 45
 Met Asn Gln Glu Ser Thr Thr Ile Lys Val Met Glu Lys Pro Pro
 50 55 60
 Phe Asp Arg Ser Ile Ser Gln Asp Ser Leu Asp Glu Leu Ser Met
 65 70 75
 Glu Asp Tyr Trp Ile Glu Leu Glu Asn Ile Lys Lys Ser Ser Glu
 80 85 90
 Asn Ser Gln Glu Asp Gln Glu Val Val Val Val Lys Glu Pro Asp

	95		100		105
Glu Gly Glu Leu	Glu Glu Glu Trp Leu	Lys Glu Ala Gly Leu Ser			
	110	115		120	
Asn Leu Phe Gly	Glu Ser Ala Gly Asp	Pro Gln Glu Ser Ile Val			
	125	130		135	
Phe Leu Ser Thr	Leu Thr Arg Thr Gln	Ala Ala Ala Val Gln Lys			
	140	145		150	
Arg Val Glu Thr	Val Ser Gln Thr Leu	Arg Lys Lys Asn Lys Gln			
	155	160		165	
Tyr Gln Ile Pro	Asp Val Arg Asp Ile	Phe Ala Gln Gln Arg Glu			
	170	175		180	
Ser Lys Glu Thr	Ala Pro Gly Gly Thr	Glu Ser Gln Ser Leu Arg			
	185	190		195	
Thr Asn Glu Asn	Lys Tyr Gln Gly Arg	Asp Asp Glu Ala Ser Asn			
	200	205		210	
Leu Val Gly Glu	Glu Lys Leu Ile Pro	Pro Glu Glu Thr Pro Ala			
	215	220		225	
Pro Glu Thr Asp	Ile Asn Leu Glu Val	Ser Phe Ala Glu Gln Ala			
	230	235		240	
Leu Asn Gln Lys	Glu Arg Ser Lys Glu	Lys Ile Gln Lys Ser Lys			
	245	250		255	
Gly Asp Asp Ala	Thr Leu Pro Ser Phe	Arg Leu Pro Lys Asp Lys			
	260	265		270	
Thr Gly Thr Thr	Arg Ile Gly Asp Leu	Ala Pro Gln Asp Met Lys			
	275	280		285	
Lys Val Cys His	Leu Ala Leu Ile Glu	Leu Thr Ala Leu Tyr Asp			
	290	295		300	
Val Leu Gly Ile	Glu Leu Lys Gln Gln	Lys Ala Val Lys Ile Lys			
	305	310		315	
Thr Lys Asp Ser	Gly Leu Phe Cys Val	Pro Leu Thr Ala Leu Leu			
	320	325		330	
Glu Gln Asp Gln	Arg Lys Val Pro Gly	Met Arg Ile Pro Leu Ile			
	335	340		345	
Phe Gln Lys Leu	Ile Ser Arg Ile Glu	Glu Arg Gly Leu Glu Thr			
	350	355		360	
Glu Gly Leu Leu	Arg Ile Pro Gly Ala	Ala Ile Arg Ile Lys Asn			
	365	370		375	
Leu Cys Gln Glu	Leu Glu Ala Lys Phe	Tyr Glu Gly Thr Phe Asn			
	380	385		390	
Trp Glu Ser Val	Lys Gln His Asp Ala	Ala Ser Leu Leu Lys Leu			
	395	400		405	

Phe Ile Arg Glu Leu Pro Gln Pro Leu Leu Ser Val Glu Tyr Leu
 410 415 420
 Lys Ala Phe Gln Ala Val Gln Asn Leu Pro Thr Lys Lys Gln Gln
 425 430 435
 Leu Gln Ala Leu Asn Leu Leu Val Ile Leu Leu Pro Asp Ala Asn
 440 445 450
 Arg Asp Thr Leu Lys Ala Leu Leu Glu Phe Leu Gln Arg Val Ile
 455 460 465
 Asp Asn Lys Glu Lys Asn Lys Met Thr Val Met Asn Val Ala Met
 470 475 480
 Val Met Ala Pro Asn Leu Phe Met Cys His Ala Leu Gly Leu Lys
 485 490 495
 Ser Ser Glu Gln Arg Glu Phe Val Met Ala Ala Gly Thr Ala Asn
 500 505 510
 Thr Met His Leu Leu Ile Lys Tyr Gln Lys Leu Leu Trp Thr Ile
 515 520 525
 Pro Lys Phe Ile Val Asn Gln Val Arg Lys Gln Asn Thr Glu Asn
 530 535 540
 His Lys Lys Asp Lys Arg Ala Met Lys Lys Leu Leu Lys Lys Met
 545 550 555
 Ala Tyr Asp Arg Glu Lys Tyr Glu Lys Gln Asp Lys Ser Thr Asn
 560 565 570
 Asp Ala Asp Val Pro Gln Gly Val Ile Arg Val Gln Ala Pro His
 575 580 585
 Leu Ser Lys Val Ser Met Ala Ile Gln Leu Thr Glu Glu Leu Lys
 590 595 600
 Ala Ser Asp Val Leu Ala Arg Phe Leu Ser Gln Glu Ser Gly Val
 605 610 615
 Ala Gln Thr Leu Lys Lys Gly Glu Val Phe Leu Tyr Glu Ile Gly
 620 625 630
 Gly Asn Ile Gly Glu Arg Cys Leu Asp Asp Asp Thr Tyr Met Lys
 635 640 645
 Asp Leu Tyr Gln Leu Asn Pro Asn Ala Glu Trp Val Ile Lys Ser
 650 655 660
 Lys Pro Leu

<210> 15

<211> 1327

<212> DNA

<213> Homo sapiens

<220>

<221> unsure
 <222> 74, 1306
 <223> unknown base

<400> 15
 gcggggctgc cctgcagcgc acagcccat cctgcagcca ccaatggcat 50
 gacccagggc cccggcactg cctngtgtga ggggctggca gctttccaac 100
 tgcagcaagt ggaggcccct gccagcttcg ggcctgtggg caggggctca 150
 gtggggcagg ggtgtggctg ccccgcccgg cagcctgca cctgtctcct 200
 ctttgtgacc agtaccgcaa ggggatcatc tcgggctccg tctgccagga 250
 cctgtgtgag ctgcat atg gtg gag tgg agg acc tgc ctc 290
 Met Val Glu Trp Arg Thr Cys Leu
 1 5
 tcg gtg gcc ccg ggc cag cag gtg tac agc ggg ctc tgg 329
 Ser Val Ala Pro Gly Gln Gln Val Tyr Ser Gly Leu Trp
 10 15 20
 cgg gac aag gat gta acc atc aag tgt ggc att gag gag 368
 Arg Asp Lys Asp Val Thr Ile Lys Cys Gly Ile Glu Glu
 25 30
 acc ctc gac tcc aag gcc cgg tcg gat gcg gcc ccc cgg 407
 Thr Leu Asp Ser Lys Ala Arg Ser Asp Ala Ala Pro Arg
 35 40 45
 cgg gag ctg gta ctg ttt gac aag ccc acc cgg ggc acc 446
 Arg Glu Leu Val Leu Phe Asp Lys Pro Thr Arg Gly Thr
 50 55 60
 tcc atc aag gaa ttc cgg gag atg acc ctc ggc ttc ctc 485
 Ser Ile Lys Glu Phe Arg Glu Met Thr Leu Gly Phe Leu
 65 70
 aag gcg aac ctg gga gac ctg cct tcc ctg ccg gcg ctg 524
 Lys Ala Asn Leu Gly Asp Leu Pro Ser Leu Pro Ala Leu
 75 80 85
 gtt ggc cag gtc ctg ctc atg gct gac ttc aac aag gac 563
 Val Gly Gln Val Leu Leu Met Ala Asp Phe Asn Lys Asp
 90 95
 aac cgg gtg tcc ctg gcg gaa gcc aag tcc gtg tgg gcc 602
 Asn Arg Val Ser Leu Ala Glu Ala Lys Ser Val Trp Ala
 100 105 110
 ctg ctg cag cgt aac gag ttc ctg ctg ctg ctg tcc ctg 641
 Leu Leu Gln Arg Asn Glu Phe Leu Leu Leu Leu Ser Leu
 115 120 125
 cag gag aag gag cac gcc tcc aga ctg ctg ggc tac tgt 680
 Gln Glu Lys Glu His Ala Ser Arg Leu Leu Gly Tyr Cys
 130 135
 ggg gac ctc tac ctc acc gag ggc gtg ccg cat ggc gcc 719
 Gly Asp Leu Tyr Leu Thr Glu Gly Val Pro His Gly Ala

005001 "0544850

140	145	150	
tgg cac gcg gcc gcc ctc cca ccc ctg ttg cgc cca ctg			758
Trp His Ala Ala Ala Leu Pro Pro Leu Leu Arg Pro Leu			
155	160		
ctg ccg cct gcc ctg cag ggt gct ctc cag cag tgg ctg			797
Leu Pro Pro Ala Leu Gln Gly Ala Leu Gln Gln Trp Leu			
165	170	175	
ggg cct gcg tgg cct tgg cgg gcc aag atc gcc atc ggc			836
Gly Pro Ala Trp Pro Trp Arg Ala Lys Ile Ala Ile Gly			
180	185	190	
ctg ctg gag ttc gtg gag gag ctc ttc cac ggc tct tac			875
Leu Leu Glu Phe Val Glu Glu Leu Phe His Gly Ser Tyr			
195	200		
ggg act ttc tac atg tgt gag acc aca ctg gcc aac gtg			914
Gly Thr Phe Tyr Met Cys Glu Thr Thr Leu Ala Asn Val			
205	210	215	
ggc tac aca gcc acc tac gag ttc aag atg gcc gag ctg			953
Gly Tyr Thr Ala Thr Tyr Asp Phe Lys Met Ala Asp Leu			
220	225		
cag cag gtg gca ccc gag gcc acc gtg cgc cgc ttc ctg			992
Gln Gln Val Ala Pro Glu Ala Thr Val Arg Arg Phe Leu			
230	235	240	
cag ggc cgc cgc tgc gag cac agc acc gac tgc acc tac			1031
Gln Gly Arg Arg Cys Glu His Ser Thr Asp Cys Thr Tyr			
245	250	255	
ggg cgc gac tgc agg gcc ccg tgt gac agg ctc atg agg			1070
Gly Arg Asp Cys Arg Ala Pro Cys Asp Arg Leu Met Arg			
260	265		
cag tgc aag ggc gac ctc atc cag ccc aac ctg gcc aag			1109
Gln Cys Lys Gly Asp Leu Ile Gln Pro Asn Leu Ala Lys			
270	275	280	
gtg tgc gca ctg cta cgg ggc tac ctg ctg cct ggc gcg			1148
Val Cys Ala Leu Leu Arg Gly Tyr Leu Leu Pro Gly Ala			
285	290		
ccc gcc gac ctc cgc gag gag ctg ggc aca cag ctg cgc			1187
Pro Ala Asp Leu Arg Glu Glu Leu Gly Thr Gln Leu Arg			
295	300	305	
acc tgt acc acg ctg agc ggg ctg gcc agc cag gtg gag			1226
Thr Cys Thr Thr Leu Ser Gly Leu Ala Ser Gln Val Glu			
310	315	320	
gcc cat cac tcg ctg gtg ctc agc cac ctc aag act ctg			1265
Ala His His Ser Leu Val Leu Ser His Leu Lys Thr Leu			
325	330		
ctc tgg aag aag atc tcc aac acc aag tac tct t g			1300
Leu Trp Lys Lys Ile Ser Asn Thr Lys Tyr Ser			
335	340	344	

atgggnaatg agggcttgca accttct 1327

<210> 16
<211> 344
<212> PRT
<213> Homo sapiens

<220>
<221> unsure
<222> 104
<223> unknown amino acid

<400> 16
Met Val Glu Trp Arg Thr Cys Leu Ser Val Ala Pro Gly Gln Gln
1 5 10 15
Val Tyr Ser Gly Leu Trp Arg Asp Lys Asp Val Thr Ile Lys Cys
20 25 30
Gly Ile Glu Glu Thr Leu Asp Ser Lys Ala Arg Ser Asp Ala Ala
35 40 45
Pro Arg Arg Glu Leu Val Leu Phe Asp Lys Pro Thr Arg Gly Thr
50 55 60
Ser Ile Lys Glu Phe Arg Glu Met Thr Leu Gly Phe Leu Lys Ala
65 70 75
Asn Leu Gly Asp Leu Pro Ser Leu Pro Ala Leu Val Gly Gln Val
80 85 90
Leu Leu Met Ala Asp Phe Asn Lys Asp Asn Arg Val Ser Xaa Ala
95 100 105
Glu Ala Lys Ser Val Trp Ala Leu Leu Gln Arg Asn Glu Phe Leu
110 115 120
Leu Leu Leu Ser Leu Gln Glu Lys Glu His Ala Ser Arg Leu Leu
125 130 135
Gly Tyr Cys Gly Asp Leu Tyr Leu Thr Glu Gly Val Pro His Gly
140 145 150
Ala Trp His Ala Ala Ala Leu Pro Pro Leu Leu Arg Pro Leu Leu
155 160 165
Pro Pro Ala Leu Gln Gly Ala Leu Gln Gln Trp Leu Gly Pro Ala
170 175 180
Trp Pro Trp Arg Ala Lys Ile Ala Ile Gly Leu Leu Glu Phe Val
185 190 195
Glu Glu Leu Phe His Gly Ser Tyr Gly Thr Phe Tyr Met Cys Glu
200 205 210
Thr Thr Leu Ala Asn Val Gly Tyr Thr Ala Thr Tyr Asp Phe Lys
215 220 225
Met Ala Asp Leu Gln Gln Val Ala Pro Glu Ala Thr Val Arg Arg
230 235 240

Phe Leu Gln Gly Arg Arg Cys Glu His Ser Thr Asp Cys Thr Tyr
 245 250 255
 Gly Arg Asp Cys Arg Ala Pro Cys Asp Arg Leu Met Arg Gln Cys
 260 265 270
 Lys Gly Asp Leu Ile Gln Pro Asn Leu Ala Lys Val Cys Ala Leu
 275 280 285
 Leu Arg Gly Tyr Leu Leu Pro Gly Ala Pro Ala Asp Leu Arg Glu
 290 295 300
 Glu Leu Gly Thr Gln Leu Arg Thr Cys Thr Thr Leu Ser Gly Leu
 305 310 315
 Ala Ser Gln Val Glu Ala His His Ser Leu Val Leu Ser His Leu
 320 325 330
 Lys Thr Leu Leu Trp Lys Lys Ile Ser Asn Thr Lys Tyr Ser
 335 340

<210> 17
 <211> 3020
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> unsure
 <222> 4, 107, 109, 112, 119, 122, 176, 323, 1616, 1878, 1881, 1911,
 1924, 1939, 1953, 1982, 1991-1992, 2000, 2443
 <223> unknown base

<400> 17
 gcgncgcgcg cctctggccg ccgccatcgc cgcttcagac ttcagaacca 50
 gcgtcgtcct caaccacctg gccgagaagt ggtttccggg ccagcgcgag 100
 tgctctngng cncggcaant cnggggagct actgcaccag gggcgctacc 150
 gggagccctg gcgccgcctg cgaggntgc gagcagaacc cagtgacttg 200
 attgtaaaaa ttacagagc ggaatcatat gctgggtctcc aagagttaa 250
 agcagcctta gaagatttaa atg cag ttc ttt ttc aac ttc 291
 Met Gln Phe Phe Phe Asn Phe
 1 5

caa gat tgg cct gaa ggt cta ctt cag gaa ang gaa aaa 330
 Gln Asp Trp Pro Glu Gly Leu Leu Gln Glu Xaa Glu Lys
 10 15 20
 gta ctc tgc gat gct ggt ttt tta ggt gat gcc tta caa 369
 Val Leu Cys Asp Ala Gly Phe Leu Gly Asp Ala Leu Gln
 25 30
 ctc ttt ctt cag tgc tta gcc ctt gat gaa gat ttt gca 408
 Leu Phe Leu Gln Cys Leu Ala Leu Asp Glu Asp Phe Ala
 35 40 45

cct gca aag ctg caa gta caa aag att tta tgt gat tta 447

```

Pro Ala Lys Leu Gln Val Gln Lys Ile Leu Cys Asp Leu
      50                      55

tta tta cct gaa aac tta aaa gaa ggc ctg aag gaa tct 486
Leu Leu Pro Glu Asn Leu Lys Glu Gly Leu Lys Glu Ser
  60                      65                      70

tcc tgg agt tca tta cca tgt act aaa aac aga cct ttt 525
Ser Trp Ser Ser Leu Pro Cys Thr Lys Asn Arg Pro Phe
      75                      80                      85

gat ttt cat tca gtg atg gaa gag tct cag tct ctc aat 564
Asp Phe His Ser Val Met Glu Glu Ser Gln Ser Leu Asn
      90                      95

gaa cct agc cca aag cag agt gaa gaa ata cca gag gtc 603
Glu Pro Ser Pro Lys Gln Ser Glu Glu Ile Pro Glu Val
  100                      105                      110

act tca gag cct gtc aaa gga agc tta aac cgt gct cag 642
Thr Ser Glu Pro Val Lys Gly Ser Leu Asn Arg Ala Gln
      115                      120

tca gca cag tct ata aat tca aca gaa atg cct gcc aga 681
Ser Ala Gln Ser Ile Asn Ser Thr Glu Met Pro Ala Arg
  125                      130                      135

gag gac tgt tta aaa aaa gtg tcc tca gaa cct gtt ctg 720
Glu Asp Cys Leu Lys Lys Val Ser Ser Glu Pro Val Leu
      140                      145                      150

tca gtt caa gaa aaa ggt gtt ctg ctg aaa aga aag ttg 759
Ser Val Gln Glu Lys Gly Val Leu Leu Lys Arg Lys Leu
      155                      160

tct ctt tta gaa cag gat gtg att gta aat gaa gat gga 798
Ser Leu Leu Glu Gln Asp Val Ile Val Asn Glu Asp Gly
  165                      170                      175

aga aat aag ctg aaa aaa caa gga gaa act ccc aat gaa 837
Arg Asn Lys Leu Lys Lys Gln Gly Glu Thr Pro Asn Glu
      180                      185

gtc tgt atg ttt tcc tta gct tat ggt gat att cca gaa 876
Val Cys Met Phe Ser Leu Ala Tyr Gly Asp Ile Pro Glu
  190                      195                      200

gaa tta atc gat gtc tca gat ttc gag tgt tct ctc tgc 915
Glu Leu Ile Asp Val Ser Asp Phe Glu Cys Ser Leu Cys
  205                      210                      215

atg agg ttg ttt ttt gag cca gta aca acc cct tgc gga 954
Met Arg Leu Phe Phe Glu Pro Val Thr Thr Pro Cys Gly
      220                      225

cat tcg ttc tgt aag aat tgt ctt gag cgt tgt tta gat 993
His Ser Phe Cys Lys Asn Cys Leu Glu Arg Cys Leu Asp
  230                      235                      240

cat gca cca tat tgt cct ctt tgc aaa gaa agc tta aat 1032
His Ala Pro Tyr Cys Pro Leu Cys Lys Glu Ser Leu Asn

```

0958458 100500

245

250

gag tat cta gca gat agg agg tac tgt gtc aca cag ctg 1071
Glu Tyr Leu Ala Asp Arg Arg Tyr Cys Val Thr Gln Leu
255 260 265

ttg gaa gaa tta ata gtg aag tat ctg cct gat gaa ctg 1110
Leu Glu Glu Leu Ile Val Lys Tyr Leu Pro Asp Glu Leu
270 275 280

tct gag aga aaa aaa ata tat gaa gaa gaa act gct gaa 1149
Ser Glu Arg Lys Lys Ile Tyr Glu Glu Glu Thr Ala Glu
285 290

ctc tca cac ttg acc aag aat gtt cca ata ttt gtt tgc 1188
Leu Ser His Leu Thr Lys Asn Val Pro Ile Phe Val Cys
295 300 305

act atg gcc tac ccc act gtg cct tgc cct ctc cat gta 1227
Thr Met Ala Tyr Pro Thr Val Pro Cys Pro Leu His Val
310 315

ttt gag cca aga tac aga ttg atg att cga aga agt ata 1266
Phe Glu Pro Arg Tyr Arg Leu Met Ile Arg Arg Ser Ile
320 325 330

cag act gga acc aaa cag ttt ggc atg tgt gtc agt gat 1305
Gln Thr Gly Thr Lys Gln Phe Gly Met Cys Val Ser Asp
335 340 345

aca caa aat agt ttt gca gat tat ggt tgt atg tta caa 1344
Thr Gln Asn Ser Phe Ala Asp Tyr Gly Cys Met Leu Gln
350 355

att aga aac gtg cat ttc tta ccg gac gga agg tct gtg 1383
Ile Arg Asn Val His Phe Leu Pro Asp Gly Arg Ser Val
360 365 370

gtt gat aca gtt gga gga aag cgg ttt agg gtt tta aaa 1422
Val Asp Thr Val Gly Gly Lys Arg Phe Arg Val Leu Lys
375 380

aga gga atg aaa gat gga tat tgc act gcc gac att gaa 1461
Arg Gly Met Lys Asp Gly Tyr Cys Thr Ala Asp Ile Glu
385 390 395

tat ctg gaa gat gtt aag gtt gag aat gaa gat gag att 1500
Tyr Leu Glu Asp Val Lys Val Glu Asn Glu Asp Glu Ile
400 405 410

aag aat ctc aga gag ctt cat gat ttg gtt tac tct caa 1539
Lys Asn Leu Arg Glu Leu His Asp Leu Val Tyr Ser Gln
415 420

gcc tgc agc tgg ttt cag aat tta aga gac aga ttt cga 1578
Ala Cys Ser Trp Phe Gln Asn Leu Arg Asp Arg Phe Arg
425 430 435

agc caa att ctt cag cat ttc gga tca atg ccc gag ang 1617
Ser Gln Ile Leu Gln His Phe Gly Ser Met Pro Glu Xaa
440 445

gag gaa aac ctt cag gca gcc cct aat gga cct gca tgg 1656
 Glu Glu Asn Leu Gln Ala Ala Pro Asn Gly Pro Ala Trp
 450 455 460

tgt tgg tgg ctt ctt gca gtt ctc cct gta gac cca cga 1695
 Cys Trp Trp Leu Leu Ala Val Leu Pro Val Asp Pro Arg
 465 470 475

tac cag ctg tgc gtt ttg tca atg aag tct ttg aaa gaa 1734
 Tyr Gln Leu Ser Val Leu Ser Met Lys Ser Leu Lys Glu
 480 485

cgg ttg acc aag ata cag cat ata ctg acc tat ttt tct 1773
 Arg Leu Thr Lys Ile Gln His Ile Leu Thr Tyr Phe Ser
 490 495 500

aga gac caa tct aag t a actaactctt tggatctccc 1810
 Arg Asp Gln Ser Lys
 505 506

tttaaagtga ccctaactctg gctgcattga tggccagatt gtctgctgcc 1860
 tttgcacatc tagtgctnng ntttcagaaa tttaatgaaa cttttctttt 1910
 ntccctcgac ctcnctgaat catgtggtnt ctgcaaatga atnaccttca 1960
 actaggattt agaccactaa gnaacttgca nncagaaaaan acacgcattg 2010
 aatgtgtgtc gaacctctac attgtgaagt tgcactatgt accatactct 2060
 aaaatgaaat aagaactctt tatgtctgtg agagagtgtg tgtgtgtgtg 2110
 tgcgtgctgtg tgtgcttgtg ggggttgggt agtgtgtgtg tattttctct 2160
 ggcttttaaaa ttttaaaaca aacaaaccaa aaagccatag agagcagaac 2210
 ttgccgaggg tcattttattg cccaagttta caagagtagc gatacaagtt 2260
 tttgcaaatt gaatttgctt cagatatatc tgtcctaattg cttatatttg 2310
 cacaagtatg taaaatatcg tgttgaggat cattctttgt tggaaatact 2360
 gctcttgcgtg aactgtcttg accattgact atgacacagt ttcttattta 2410
 tgtaaatact tgcatcacag tggccgacag atntgggatg cagaacctag 2460
 agccagtttt caggaacaat tgtaaacctg acatgggtact gtgcatctat 2510
 tcataaaaca ctcaaaactg tgaaaatatg gtttacattt aattgtacat 2560
 aaaggtaaag ggagaactca attcagtacc agttagtttg tacattttag 2610
 ggggcttttc acattaactg ccattttgtg taatttatag tttgacatga 2660
 tgtgtttgtt ttaaaaaaaaa atgcatagta taaaccatt aaggatctgg 2710
 gaaaagagaa gaagtttaat atagaactaa gcttttaaaag tttgtttttg 2760
 tttttaattc tgggtctcggg gcaaatgtta gttatgcctt attcatatca 2810

cagttagatc accatgctgc aacatgggtt atattcatgc tgcctagaa 2860
 acttttgtaa ttatttggtg caaatttggt actgtcctta ttaactttct 2910
 tttatgtaag taatttgtaa aagtttctta aaatttttgc ttttgcttat 2960
 ttaattttga ataaaagcta aattcctaataaaaaaaaaa 3010
 aaaaaaaaaa 3020

<210> 18
 <211> 506
 <212> PRT
 <213> Homo sapiens

<220>
 <221> unsure
 <222> 18, 448
 <223> unknown amino acid

<400> 18
 Met Gln Phe Phe Phe Asn Phe Gln Asp Trp Pro Glu Gly Leu Leu
 1 5 10 15
 Gln Glu Xaa Glu Lys Val Leu Cys Asp Ala Gly Phe Leu Gly Asp
 20 25 30
 Ala Leu Gln Leu Phe Leu Gln Cys Leu Ala Leu Asp Glu Asp Phe
 35 40 45
 Ala Pro Ala Lys Leu Gln Val Gln Lys Ile Leu Cys Asp Leu Leu
 50 55 60
 Leu Pro Glu Asn Leu Lys Glu Gly Leu Lys Glu Ser Ser Trp Ser
 65 70 75
 Ser Leu Pro Cys Thr Lys Asn Arg Pro Phe Asp Phe His Ser Val
 80 85 90
 Met Glu Glu Ser Gln Ser Leu Asn Glu Pro Ser Pro Lys Gln Ser
 95 100 105
 Glu Glu Ile Pro Glu Val Thr Ser Glu Pro Val Lys Gly Ser Leu
 110 115 120
 Asn Arg Ala Gln Ser Ala Gln Ser Ile Asn Ser Thr Glu Met Pro
 125 130 135
 Ala Arg Glu Asp Cys Leu Lys Lys Val Ser Ser Glu Pro Val Leu
 140 145 150
 Ser Val Gln Glu Lys Gly Val Leu Leu Lys Arg Lys Leu Ser Leu
 155 160 165
 Leu Glu Gln Asp Val Ile Val Asn Glu Asp Gly Arg Asn Lys Leu
 170 175 180
 Lys Lys Gln Gly Glu Thr Pro Asn Glu Val Cys Met Phe Ser Leu
 185 190 195
 Ala Tyr Gly Asp Ile Pro Glu Glu Leu Ile Asp Val Ser Asp Phe

200										205					210				
Glu	Cys	Ser	Leu	Cys	Met	Arg	Leu	Phe	Phe	Glu	Pro	Val	Thr	Thr					
				215					220					225					
Pro	Cys	Gly	His	Ser	Phe	Cys	Lys	Asn	Cys	Leu	Glu	Arg	Cys	Leu					
				230					235					240					
Asp	His	Ala	Pro	Tyr	Cys	Pro	Leu	Cys	Lys	Glu	Ser	Leu	Asn	Glu					
				245					250					255					
Tyr	Leu	Ala	Asp	Arg	Arg	Tyr	Cys	Val	Thr	Gln	Leu	Leu	Glu	Glu					
				260					265					270					
Leu	Ile	Val	Lys	Tyr	Leu	Pro	Asp	Glu	Leu	Ser	Glu	Arg	Lys	Lys					
				275					280					285					
Ile	Tyr	Glu	Glu	Glu	Thr	Ala	Glu	Leu	Ser	His	Leu	Thr	Lys	Asn					
				290					295					300					
Val	Pro	Ile	Phe	Val	Cys	Thr	Met	Ala	Tyr	Pro	Thr	Val	Pro	Cys					
				305					310					315					
Pro	Leu	His	Val	Phe	Glu	Pro	Arg	Tyr	Arg	Leu	Met	Ile	Arg	Arg					
				320					325					330					
Ser	Ile	Gln	Thr	Gly	Thr	Lys	Gln	Phe	Gly	Met	Cys	Val	Ser	Asp					
				335					340					345					
Thr	Gln	Asn	Ser	Phe	Ala	Asp	Tyr	Gly	Cys	Met	Leu	Gln	Ile	Arg					
				350					355					360					
Asn	Val	His	Phe	Leu	Pro	Asp	Gly	Arg	Ser	Val	Val	Asp	Thr	Val					
				365					370					375					
Gly	Gly	Lys	Arg	Phe	Arg	Val	Leu	Lys	Arg	Gly	Met	Lys	Asp	Gly					
				380					385					390					
Tyr	Cys	Thr	Ala	Asp	Ile	Glu	Tyr	Leu	Glu	Asp	Val	Lys	Val	Glu					
				395					400					405					
Asn	Glu	Asp	Glu	Ile	Lys	Asn	Leu	Arg	Glu	Leu	His	Asp	Leu	Val					
				410					415					420					
Tyr	Ser	Gln	Ala	Cys	Ser	Trp	Phe	Gln	Asn	Leu	Arg	Asp	Arg	Phe					
				425					430					435					
Arg	Ser	Gln	Ile	Leu	Gln	His	Phe	Gly	Ser	Met	Pro	Xaa	Arg	Glu					
				440					445					450					
Glu	Asn	Leu	Gln	Ala	Ala	Pro	Asn	Gly	Pro	Ala	Trp	Cys	Trp	Trp					
				455					460					465					
Leu	Leu	Ala	Val	Leu	Pro	Val	Asp	Pro	Arg	Tyr	Gln	Leu	Ser	Val					
				470					475					480					
Leu	Ser	Met	Lys	Ser	Leu	Lys	Glu	Arg	Leu	Thr	Lys	Ile	Gln	His					
				485					490					495					
Ile	Leu	Thr	Tyr	Phe	Ser	Arg	Asp	Gln	Ser	Lys									
				500					505										

<210> 19
<211> 310
<212> DNA
<213> Homo sapiens

<220>
<221> unsure
<222> 300
<223> unknown base

<400> 19
ggatccctgg agctgccatt agaatcaaga atctttgccca agaactagaa 50

gcaaagtttt atgaagggac ttttaattgg gaaagtgtca aacagcatga 100

tgccgccagc ctgctgaagc ttttcattcg ggagttgccc cagccactgc 150

tcagtgtgga gtatctcaaa gcctttcagg ctgtccagaa tttccaacc 200

aagaagcagc aactacaggg cttttgaacc ctttctttta ctcatctctg 250

atgtttgata cccctgggtg aaaacaattc agtaaagcat cctgcctcan 300

accccccccc 310

<210> 20
<211> 26
<212> DNA
<213> Homo sapiens

<400> 20
gctgcttctt ggttggaaga ttctgg 26

<210> 21
<211> 26
<212> DNA
<213> Homo sapiens

<400> 21
ccagaatctt ccaaccaaga agcagc 26

<210> 22
<211> 53
<212> DNA
<213> Homo sapiens

<400> 22
gcacatgct gtttgacact ttccaatta aaagtcctt cataaaactt 50

tgc 53

<210> 23
<211> 30
<212> DNA
<213> Homo sapiens

<400> 23
ggagctgccca ttagaatcaa gaatctttgc 30

<210> 24

<211> 533
 <212> DNA
 <213> Homo sapiens

 <400> 24
 ttttattagg aatttagctt ttattcaaaa ttaaataagc aaaagcaaaa 50
 attttaagaa acttttacaa attacttaca taaaagaaag ttaataagga 100
 cagtcacaaa ttgcaacaa ataattacaa aagtttctag ggcagcatga 150
 atataaacca tgttgacgca tgggtgatcta actgtgatat gaataaggca 200
 taactaacat ttgcaccgag accagaatta aaaacaaaaa caaacttta 250
 aagcttagtt ctatattaaa cttcttctct tttcccagat ccttaatggg 300
 tttatactat gcattttttt ttaaaacaaa cacatcatgt caaactataa 350
 attacacaaa tgggcagtta atgtgaaaag ccccttaaaa tgtacaaact 400
 aactgggtact gaattgagtt ctccctttac ctttatgtac aattaaatgt 450
 aaaccatatt ttcacagttt tgagtgtttt atgaatagat gcacagtacc 500
 atgtcagggt tacaattggt cctgaaaact ggc 533

<210> 25
 <211> 433
 <212> DNA
 <213> Homo sapiens

 <400> 25
 agcagcagca ggaactcggt acgctgcagc agggcccaca cggacttggc 50
 ttccgccag gacaccgggt tgtccttggt gaagtcagcc atgagcagga 100
 cctggccaac cagcgccggc agggaaggca ggtctcccag gttcgccttg 150
 aggaagccga gggatcatct ccggaattcc ttgatggagg tgccccgggt 200
 gggcttgtea aacagtacca gctcccgccg ggggcgcgat ccgaccgggc 250
 cttggagtcg agggctctct caatgccaca cttgatgggt acatccttgt 300
 cccgccagag ccogctgtac acctgctggc ccggggcaca cgagaagcag 350
 gtcctccact ccaccatag cagctcacac aggtcctggc agacggagcc 400
 cgagatgac cccttgcgggt actggtcaca aat 433

<210> 26
 <211> 22
 <212> DNA
 <213> Homo sapiens

<400> 26
 atgcagttct ttttcaactt cc 22

<210> 27
 <211> 20

```

<212> DNA
<213> Homo sapiens

<400> 27
ctagagacca atctaagtaa 20

<210> 28
<211> 20
<212> DNA
<213> Homo sapiens

<400> 28
atggtggagt ggaggacctg 20

<210> 29
<211> 22
<212> DNA
<213> Homo sapiens

<400> 29
ctccaacacc aagtactctt ga 22

```